

A LONG CHAIN OF P-POINTS

BORISA KUZELJEVIC AND DILIP RAGHAVAN

ABSTRACT. The notion of a δ -generic sequence of P-points is introduced in this paper. It is proved assuming the Continuum Hypothesis that for each $\delta < \omega_2$, any δ -generic sequence of P-points can be extended to an ω_2 -generic sequence. This shows that the Continuum Hypothesis implies that there is a chain of P-points of length \mathfrak{c}^+ with respect to both Rudin-Keisler and Tukey reducibility. These results answer an old question of Andreas Blass.

1. INTRODUCTION

In his 1973 paper on the structure of P-points [1], Blass posed the following question:

Question 1.1 (Question 4 of [1]). What ordinals can be embedded into the class of P-points when equipped with the ordering of Rudin-Keisler reducibility assuming Martin's Axiom?

Recall that an ultrafilter \mathcal{U} on ω is called a *P-point* if for any $\{a_n : n < \omega\} \subset \mathcal{U}$ there is $a \in \mathcal{U}$ such that $a \subset^* a_n$, for every $n < \omega$. All filters \mathcal{U} occurring in this paper are assumed to be *proper* – meaning that $0 \notin \mathcal{U}$ – and *non-principal* – meaning that \mathcal{U} extends the filter of co-finite sets. It is not hard to see that an ultrafilter \mathcal{U} is a P-point if and only if every $f \in \omega^\omega$ becomes either constant or finite-to-one on a set in \mathcal{U} . Recall also the well-known Rudin-Keisler ordering on P-points.

Definition 1.2. Let \mathcal{U} and \mathcal{V} be ultrafilters on ω . We say that $\mathcal{U} \leq_{RK} \mathcal{V}$, i.e. \mathcal{U} is *Rudin-Keisler (RK) reducible* to \mathcal{V} or \mathcal{U} is *Rudin-Keisler (RK) below* \mathcal{V} , if there is $f \in \omega^\omega$ such that $A \in \mathcal{U} \Leftrightarrow f^{-1}(A) \in \mathcal{V}$ for every $A \subset \omega$. We say that $\mathcal{U} \equiv_{RK} \mathcal{V}$, i.e. \mathcal{U} is *RK equivalent* to \mathcal{V} , if $\mathcal{U} \leq_{RK} \mathcal{V}$ and $\mathcal{V} \leq_{RK} \mathcal{U}$.

It is worth noting here that the class of P-points is downwards closed with respect to this order. In other words, if \mathcal{U} is a P-point, then every ultrafilter that is RK below \mathcal{U} is also a P-point. It should also be noted that the existence of P-points cannot be proved in ZFC by a celebrated result of Shelah (see [13]). Hence it is natural to assume some principle that guarantees the existence of “many” P-points when studying their properties under the RK or other similar orderings. Common examples of such principles include the Continuum Hypothesis (CH), Martin's axiom (MA), and Martin's Axiom for σ -centered posets (MA(σ – centered)).

Blass showed in [1] that the ordinal ω_1 can be embedded into the P-points with respect to the RK ordering, if MA(σ – centered) holds. In particular under CH, the ordinal $\mathfrak{c} = 2^{\aleph_0}$ embeds into the P-points with respect to RK reducibility. Note that no ultrafilter \mathcal{V} can have more than \mathfrak{c} predecessors in the RK order. This is because for each $f \in \omega^\omega$, there can be at most one ultrafilter \mathcal{U} for which f witnesses the relation $\mathcal{U} \leq_{RK} \mathcal{V}$. Therefore there can be no RK-chain of P-points of length $\mathfrak{c}^+ + 1$. Thus the strongest possible positive answer to Question 1.1 is that the ordinal \mathfrak{c}^+ embeds into the P-points under the RK ordering.

Date: March 5, 2018.

2010 Mathematics Subject Classification. 03E50, 03E05, 03E35, 54D80.

Key words and phrases. ultrafilter, Rudin-Keisler order, Tukey reducibility, P-point.

Both authors were partially supported by NUS research grant number R-146-000-161-133. The first author was partially supported by the MPNTR grant ON174006.

Though there have not been many advances directly pertaining to Question 1.1 after [1], several results have dealt with closely related issues. Rosen [11] showed assuming CH that the ordinal ω_1 occurs as an RK initial segment of the P-points. In other words, he produced a strictly increasing RK chain of P-points of length ω_1 that is downwards closed under the relation \leq_{RK} up to RK equivalence. Laflamme [5] further investigated well-ordered initial segments of the P-points under the RK ordering. For each countable ordinal α , he produced a forcing notion \mathbb{P}_α that generically adds an RK initial segment of the P-points of order type α . He also gave combinatorial characterizations of the generics added by these forcing notions.

Interest in the Tukey types of ultrafilters on ω was recently revived by Milovich [7]. Dobrinen and Todorćevic [2] considered the Tukey ordering on P-points. Recall that for any $\mathcal{X}, \mathcal{Y} \subset \mathcal{P}(\omega)$, a map $\phi : \mathcal{X} \rightarrow \mathcal{Y}$ is said to be *monotone* if for every $a, b \in \mathcal{X}$, $a \subset b$ implies $\phi(a) \subset \phi(b)$, while ϕ is said to be *cofinal* in \mathcal{Y} if for every $b \in \mathcal{Y}$ there is $a \in \mathcal{X}$ so that $\phi(a) \subset b$.

Definition 1.3. We say that $\mathcal{U} \leq_T \mathcal{V}$, i.e. \mathcal{U} is *Tukey reducible* to \mathcal{V} or \mathcal{U} is *Tukey below* \mathcal{V} , if there is a monotone $\phi : \mathcal{V} \rightarrow \mathcal{U}$ which is cofinal in \mathcal{U} . We say that $\mathcal{U} \equiv_T \mathcal{V}$, i.e. \mathcal{U} is *Tukey equivalent* to \mathcal{V} , if $\mathcal{U} \leq_T \mathcal{V}$ and $\mathcal{V} \leq_T \mathcal{U}$.

It is not hard to see that $\mathcal{U} \leq_{RK} \mathcal{V}$ implies $\mathcal{U} \leq_T \mathcal{V}$, and it was proved by Raghavan and Todorćevic in [9] that CH implies the existence of P-points \mathcal{U} and \mathcal{V} such that $\mathcal{V} <_{RK} \mathcal{U}$, but $\mathcal{V} \equiv_T \mathcal{U}$. Their result showed that the orders \leq_T and \leq_{RK} can diverge in a strong sense even within the realm of P-points, although by another result from [9], the two orders coincide within the realm of selective ultrafilters. In [2], Dobrinen and Todorćevic showed that every P-point has only \mathfrak{c} Tukey predecessors by establishing the following useful fact.

Theorem 1.4 (Dobrinen and Todorćevic [2]). *If \mathcal{V} is a P-point and \mathcal{U} is any ultrafilter with $\mathcal{U} \leq_T \mathcal{V}$, then there is a continuous monotone $\phi : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$ such that $\phi \upharpoonright \mathcal{V} : \mathcal{V} \rightarrow \mathcal{U}$ is a monotone map that is cofinal in \mathcal{U} .*

They used this in [2] to embed the ordinal ω_1 into the class of P-points equipped with the ordering of Tukey reducibility assuming $\text{MA}(\sigma\text{-centered})$. Question 54 of [2] asks whether there is a strictly increasing Tukey chain of P-points of length \mathfrak{c}^+ . In [3] and [4], Dobrinen and Todorćevic proved some analogues of Laflamme's results mentioned above for the Tukey order. In particular, they showed that each countable ordinal occurs as a Tukey initial segment of the class of P-points, assuming $\text{MA}(\sigma\text{-centered})$. Raghavan and Shelah proved in [8] that $\text{MA}(\sigma\text{-centered})$ implies that the Boolean algebra $\mathcal{P}(\omega)/\text{FIN}$ equipped with its natural ordering embeds into the P-points with respect to both the RK and Tukey orders. In particular, for each $\alpha < \mathfrak{c}^+$, the ordinal α embeds into the P-points with respect to both of these orders.

In this paper, we give a complete answer to Question 1.1 by showing that the ordinal \mathfrak{c}^+ can be embedded into the P-points under RK reducibility. Our chain of P-points of length \mathfrak{c}^+ will also be strictly increasing with respect to Tukey reducibility, so we get a positive answer to Question 54 of [2] as well. The construction will be presented assuming CH for simplicity. However the same construction can be run under MA with some fairly straightforward modifications. We will try to point out these necessary modifications at the appropriate places in the proofs below. We will make use of Theorem 1.4 in our construction to ensure that our chain is also strictly increasing in the sense of Tukey reducibility. However the continuity of the monotone maps will not be important for us. Rather any other fixed collection of \mathfrak{c} many monotone maps from $\mathcal{P}(\omega)$ to itself which is large enough to catch all Tukey reductions from any P-point will suffice. For instance, it was proved in [9] that the collection of monotone maps of the first Baire class suffice to catch all Tukey reductions from any basically generated ultrafilter, which form a larger class of ultrafilters than the P-points. So we could equally well have used monotone maps of the first Baire class in our construction.

A powerful machinery for constructing objects of size \aleph_2 under \diamond was introduced by Shelah, Laflamme, and Hart in [15]. This machinery can be used to build a chain of P-points of length ω_2 that is strictly increasing with respect to both RK and Tukey reducibility assuming \diamond . More generally, the methods in [15] allow for the construction of certain types of objects of size λ^+ from a principle called DI_λ , which is closely related to \diamond_λ . Shelah's results in [14] imply that DI_\aleph_1 follows from MA when $\aleph_1 < \aleph_2$ and is a successor cardinal. Thus the methods of [15], when combined with the results of [14], can also be used to get a chain of P-points of length \aleph_1^+ when $\aleph_1 < \aleph_2$, \aleph_1 is a successor cardinal, and MA holds. However the techniques from [15] are inadequate to treat the case when only CH holds¹.

Some of the ideas in this paper have been elaborated on by Raghavan and Verner [10] recently to prove that CH implies that the class of rapid P-points is \aleph_1^+ -closed, meaning that any RK-increasing chain of rapid P-points of length at most \aleph_1 has an RK-upper bound which is a rapid P-point. Therefore every chain of rapid P-points of length at most \aleph_1 can be extended to a chain of rapid P-points of length \aleph_1^+ so that long chains of rapid P-points abound. However the construction from [10] does not produce any long chains that are increasing with respect to \leq_{RB}^+ . Hence the existence result in this paper is stronger in a certain sense than the existence result that can be deduced using the methods in [10] (see Definition 2.7 and the following discussion).

2. PRELIMINARIES

We use standard notation. “ $\forall^\infty x \dots$ ” abbreviates the quantifier “for all but finitely many $x \dots$ ” and “ $\exists^\infty x \dots$ ” stands for “there exist infinitely many x such that \dots ”. $[X]^{\aleph_0}$ refers to the collection of all subsets of X of cardinality \aleph_0 , and $[X]^{<\aleph_0}$ is the collection of all finite subsets of X . The symbol \subset^* denotes the relation of containment modulo a finite set: $a \subset^* b$ iff $a \setminus b$ is finite.

Even though the final construction uses CH, none of the preliminary lemmas rely on it. In fact, CH will only be used in Section 5. The results in Sections 2–4 are all in ZFC, and CH is needed later on to ensure that these results are sufficient to carry out the final construction and that they are applicable to it. So we do not need to make any assumptions about cardinal arithmetic at the moment.

One of the difficulties in embedding various partially ordered structures into the P-points is that, unlike the class of all ultrafilters on ω , this class is not \aleph_1 -directed under \leq_{RK} . It is not hard to prove in ZFC that if $\{\mathcal{U}_i : i < \aleph_1\}$ is an arbitrary collection of ultrafilters on ω , then there is an ultrafilter \mathcal{U} on ω such that $\forall i < \aleph_1 [\mathcal{U}_i \leq_{RK} \mathcal{U}]$. However it is well-known that there are two P-points \mathcal{U} and \mathcal{V} with no RK upper bound that is a P-point under CH (see [1]). Even if we restrict ourselves to chains, it is easy to construct, assuming CH, an RK chain of P-points $\langle \mathcal{U}_i : i < \omega_1 \rangle$ which has no P-point upper bound. The strategy for ensuring that our chains are always extensible is to make each ultrafilter “very generic” (with respect to some partial order to be defined in Section 4). The same strategy was used in [8], but with one crucial difference. Only \aleph_1 many ultrafilters were constructed in [8] and so all of the ultrafilters in question could be built simultaneously in \aleph_1 steps. But by the time we get to, for example, the ultrafilter \mathcal{U}_{ω_1} in our present construction, all of the ultrafilters \mathcal{U}_i , for $i < \omega_1$, will have been fully determined with no room for further improvements. Thus the ultrafilters that were built before should have already predicted and satisfied the requirements imposed by \mathcal{U}_{ω_1} , and indeed by all of the ultrafilters to come in future. This is possible because there are only ω_1 possible initial segments of ultrafilters. More precisely, each of the ω_2 many P-points is generated by a \subset^* -descending tower $A_\alpha = \langle C_i^\alpha : i < \omega_1 \rangle$. For each $j < \omega_1$, the collection $\{A_\alpha \upharpoonright j : \alpha < \omega_2\}$ just has size ω_1 . This leads to the notion of a δ -generic sequence, which is essentially an RK-chain of P-points of length δ where every ultrafilter in the sequence has predicted and met certain requirements involving such

¹Personal communication with Shelah. The second author thanks Shelah for discussing these issues with him.

initial segments of potential future ultrafilters and potential RK maps going from such initial segments into it. The precise definition is given in Definition 2.10. Our main result is that such generic sequences can always be extended.

Remark 2.1. In the rest of the paper we will use the following notation:

- For $A \subset \omega$, define $A^0 = A$ and $A^1 = \omega \setminus A$.
- For $A \in [\omega]^{\aleph_0}$, $A(m)$ is the m th element of A in its increasing enumeration.
- For $A \in [\omega]^{\aleph_0}$ and $k, m < \omega$ let $A[k, m] = \{A(l) : k \leq l < m\}$.
- For $A \in [\omega]^{\aleph_0}$ and $m < \omega$, let $A[m] = \{A(l) : l \geq m\}$.
- For a sequence $c = \langle c(\xi) : \xi < \alpha \rangle$, let $\text{set}(c) = \bigcup_{\xi < \alpha} c(\xi)$.
- For a sequence $c = \langle c(\xi) : \xi < \alpha \rangle$ and $\eta < \alpha$, let $\text{set}(c) \upharpoonright \eta = \bigcup_{\eta \leq \xi < \alpha} c(\xi)$.
- For $m \in \omega$, let $s(m) = m(m+1)/2$ and $t(m) = \sum_{s(m) \leq k < s(m+1)} (k+1)$.
- A function $f \in \omega^\omega$ is *increasing* if $\forall n \in \omega [f(n) \leq f(n+1)]$.

We also consider, for an ordinal α , triples $\rho = \langle \bar{D}, \bar{K}, \bar{\pi} \rangle$ where $\bar{D} = \langle D_n : n < \alpha \rangle$ is a sequence of sets in $\mathcal{P}(\omega)$, $\bar{K} = \langle K_{m,n} : m \leq n < \alpha \rangle$ is a sequence in ω and $\bar{\pi} = \langle \pi_{n,m} : m \leq n < \alpha \rangle$ is a sequence in ω^ω . Then, for $n < \alpha$, denote:

- $\Delta_n^\rho = \bigcup_{m \leq n} H_{m,n}^\rho$ where $H_{m,n}^\rho$ is the set of all $k \in D_m \setminus [K_{m,n}, K_{m,n+1})$ such that $\pi_{m',m}^{-1}(\{k\}) \cap D_{m'} \setminus [K_{m',n}, K_{m',n+1}) = \emptyset$ for all $m < m' \leq n$;
- $L_0^\rho = \emptyset$ and for each $n \in \omega$, $L_{n+1}^\rho = L_n^\rho \cup \Delta_n^\rho$.

Regarding the definition of Δ_n^ρ , the reader should think of the sequence \bar{K} as defining an interval partition of D_m , for each m . Then for any $m \leq n$, $H_{m,n}^\rho$ consists of the points in the n th interval of D_m that do not have a preimage in the n th interval of $D_{m'}$ for any $m < m' \leq n$.

Next we recall the notion of a rapid ultrafilter. All the P-points in our construction will be rapid. This happens because the requirement of genericity forces our ultrafilters to contain some ‘‘very thin’’ sets. However they cannot be too thin, lest we end up with a Q-point. Rapidity turns out to be a good compromise.

Definition 2.2. We say that the ultrafilter \mathcal{U} is *rapid* if for every $f \in \omega^\omega$ there is $X \in \mathcal{U}$ such that $X(n) \geq f(n)$ for every $n < \omega$.

Lemma 2.3. If \mathcal{U} is a rapid ultrafilter, then for every $f \in \omega^\omega$ and every $X \in \mathcal{U}$ there is $Y \in \mathcal{U}$ such that $Y \subset X$ and $Y(n) \geq f(n)$ for every $n < \omega$.

Proof. Let $f \in \omega^\omega$ and $X \in \mathcal{U}$. Let Z be as in Definition 2.2, i.e. $Z \in \mathcal{U}$ and $Z(n) \geq f(n)$ for $n < \omega$. Let $Y = X \cap Z$ and note $Y \in \mathcal{U}$. Since $Y \subset Z$ we have $Y(n) \geq Z(n) \geq f(n)$ for every $n < \omega$. \square

Claim 2.4. Let $\langle \mathcal{U}_n : n < \omega \rangle$ be a sequence of distinct P-points and M a countable elementary submodel of $H_{(2^c)^+}$ containing $\langle \mathcal{U}_n : n < \omega \rangle$. For $n < \omega$ let $A_n \in \mathcal{U}_n$ be such that $A_n \subset^* A$ for each $A \in \mathcal{U}_n \cap M$. Then $|A_n \cap A_m| < \omega$ for $m < n < \omega$.

Proof. Fix $m < n < \omega$ and pick a set $A_{m,n}$ so that $A_{m,n} \in \mathcal{U}_m$ and $\omega \setminus A_{m,n} \in \mathcal{U}_n$ (this can be done because $\mathcal{U}_m \neq \mathcal{U}_n$ for $m \neq n$). By elementarity of M , since $\mathcal{U}_m, \mathcal{U}_n \in M$ we can assume that $A_{m,n} \in M$. Now we have that $A_m \subset^* A_{m,n}$ and that $A_n \subset^* \omega \setminus A_{m,n}$. So $A_m \cap A_n \subset^* \emptyset$ implying $|A_m \cap A_n| < \omega$. \square

Now we introduce one of the basic partial orders used in the construction. The definition of (\mathbb{P}, \leq) is inspired by the many examples of creature forcing in the literature, for example see [12]. However there is no notion of norm in this case, or rather the norm is simply the cardinality.

Definition 2.5. Define \mathbb{P} as the set of all functions $c : \omega \rightarrow [\omega]^{< \aleph_0} \setminus \{0\}$ such that $\forall n \in \omega [|c(n)| < |c(n+1)| \wedge \max(c(n)) < \min(c(n+1))]$. If $c, d \in \mathbb{P}$, then $c \leq d$ if there is $l < \omega$ such that $c \leq_l d$, where $c \leq_l d \Leftrightarrow \forall m \geq l \exists n \geq m [c(m) \subset d(n)]$.

Note that if $d \leq c$, then $\text{set}(d) \subset^* \text{set}(c)$. Each of our ultrafilters will be generated by a tower of the form $\langle \text{set}(c_i) : i < \omega_1 \rangle$ where $\langle c_i : i < \omega_1 \rangle$ is some decreasing sequence of conditions in \mathbb{P} . This guarantees that each ultrafilter is a P-point.

Remark 2.6. Note that $\langle \mathbb{P}, \leq \rangle$ is a partial order and has the following properties:

- (1) For any $c \in \mathbb{P}$ we have $\min(c(n)) \geq n$;
- (2) If $a \leq_n b$ and $b \leq_m c$, then $a \leq_l c$ for $l = \max\{m, n\}$. To see this take any $k \geq l$. There is $k' \geq k \geq l$ such that $a(k) \subset b(k')$. There is also $k'' \geq k' \geq k \geq l$ so that $a(k) \subset b(k') \subset c(k'')$ as required.
- (3) Let $\{d_n : n < \omega\} \subset \mathbb{P}$ be such that $\forall n < \omega [d_{n+1} \leq_{m_n} d_n]$. Then $d_{n+1} \leq_l d_0$ for $n < \omega$ and $l = \max\{m_k : k \leq n\}$. The proof is by induction on n . For $n = 0$, $d_1 \leq_{m_0} d_0$. Let $n > 0$ and assume the statement is true for all $m \leq n$. Then $d_{n+2} \leq_{m_{n+1}} d_{n+1}$ and $d_{n+1} \leq_{l_1} d_0$, where $l_1 = \max\{m_k : k \leq n\}$. So, by (2), $d_{n+2} \leq_l d_0$ for $l = \max\{m_k : k \leq n+1\}$.

Definition 2.7. A triple $\langle \pi, \psi, c \rangle$ is called a *normal triple* if $\pi, \psi \in \omega^\omega$, for every $l \leq l' < \omega$ we have that $\psi(l) \leq \psi(l')$, if $\text{ran}(\psi)$ is infinite, and if $c \in \mathbb{P}$ is such that for $l < \omega$ we have $\pi''c(l) = \{\psi(l)\}$ and for $n \in \omega \setminus \text{set}(c)$ we have $\pi(n) = 0$.

The notion of a normal triple will help us ensure that when $\alpha < \beta < \omega_2$, the RK reduction from \mathcal{U}_β to \mathcal{U}_α is witnessed by a function that is increasing on a set in \mathcal{U}_β . Thus our sequence of P-points will actually even be a chain with respect to the order \leq_{RB}^+ . Recall that for ultrafilters \mathcal{U} and \mathcal{V} on ω , $\mathcal{U} \leq_{RB}^+ \mathcal{V}$ if there is an increasing $f \in \omega^\omega$ such that $A \in \mathcal{U} \iff f^{-1}(A) \in \mathcal{V}$, for every $A \subset \omega$. More information about the order \leq_{RB}^+ can be found in [6].

Remark 2.8. Suppose that $d \leq c$ and $\langle \pi, \psi, c \rangle$ is a normal triple. There is $N < \omega$ such that for every $k, l \in \text{set}(d) \setminus N$ if $k \leq l$, then $\pi(k) \leq \pi(l)$.

Lemma 2.9. Suppose that $\langle \pi, \psi, b \rangle$ is a normal triple, that $a \subset \pi''\text{set}(c) \llbracket n_0 \rrbracket$, and that $c \leq_{n_0} b$. For $n < \omega$ denote $F_n = \{m < \omega : \pi''c(m) = \{a(n)\}\}$. Then for $n < \omega$: $F_n \setminus n_0 \neq \emptyset$, $|F_n| < \omega$ and $\max(F_n) < \min(F_{n+1} \setminus n_0) \leq \max(F_{n+1})$.

Proof. Fix $n < \omega$. By the choice of the set a there are $k \geq n_0$ and $x \in c(k)$ so that $\pi(x) = a(n)$. Since $c \leq_{n_0} b$ there is $m \geq k$ so that $c(k) \subset b(m)$ and because $\langle \pi, \psi, b \rangle$ is a normal triple we know $\pi''b(m) = \{a(n)\}$. So $k \in F_n$ is such that $k \geq n_0$, implying $F_n \setminus n_0 \neq \emptyset$. To show that each F_n is finite take any $k \in F_n$. As above, $F_{n+1} \setminus n_0 \neq \emptyset$. Let $k' = \min(F_{n+1} \setminus n_0)$. We will show $k < k'$. If $k < n_0$ the statement follows. If $k \geq n_0$ then there are m_1, m_2 so that $c(k) \subset b(m_1)$ and $c(k') \subset b(m_2)$. Since $\pi''b(m_1) = \{a(n)\}$, $\pi''b(m_2) = \{a(n+1)\}$ and $\langle \pi, \psi, b \rangle$ is a normal triple we have $m_1 < m_2$ and consequently $k < k'$. So $\max(F_n) < k'$ implying both $|F_n| < \omega$ and $\max(F_n) < \min(F_{n+1} \setminus n_0) \leq \max(F_{n+1})$. \square

Now comes the central definition of the construction. We will briefly try to explain the intuition behind each of the clauses below. Clauses (1), (2), and (4) are self explanatory and were commented on earlier. Clause (5a) guarantees that $\pi_{\beta, \alpha}$ is an RK map from \mathcal{U}_β to \mathcal{U}_α whenever $\alpha \leq \beta$. This is because if \mathcal{U}, \mathcal{V} are ultrafilters on ω and $f \in \omega^\omega$ is such that $f''b \in \mathcal{U}$ for every $b \in \mathcal{V}$, then f witnesses that $\mathcal{U} \leq_{RK} \mathcal{V}$. Clause (5b) says that if $\alpha \leq \beta \leq \gamma$, then $\pi_{\gamma, \alpha} = \pi_{\beta, \alpha} \circ \pi_{\gamma, \beta}$ modulo a set in \mathcal{U}_γ . This type of commuting of RK maps is unavoidable in a chain. Clause (5c) makes the map $\pi_{\beta, \alpha}$ increasing on a set in \mathcal{U}_β ; this makes $\mathcal{U}_\alpha \leq_{RB}^+ \mathcal{U}_\beta$. The fact that $\pi_{\beta, \alpha}$ is constant on $c_i^\beta(n)$ for almost all n is helpful for killing unwanted Tukey maps.

Clauses (3) and (6) deal with the prediction of requirements imposed by future ultrafilters. To understand (3), suppose for simplicity that $\langle \mathcal{U}_n : n < \omega \rangle$ has already been built and that \mathcal{U}_ω is being built. At a certain stage you have decided to put $\text{set}(d) \in \mathcal{U}_\omega$, for some $d \in \mathbb{P}$, and you have also decided the sequence of RK maps $\langle \pi_{\omega, i} : i \leq n \rangle$,

for some $n \in \omega$. In particular $\pi''_{\omega,n} \text{set}(d) \in \mathcal{U}_n$. Now you wish to decide $\pi_{\omega,n+1}$ and you are permitted to extend d to some $d^* \leq d$ in the process. But you must ensure that $\pi''_{\omega,n+1} \text{set}(d^*) \in \mathcal{U}_{n+1}$ and that $\pi_{\omega,n}$ commutes through $\pi_{\omega,n+1}$. Clause (3) says that \mathcal{U}_{n+1} anticipated this requirement and that there is a $b \in \mathcal{U}_{n+1}$ (in fact cofinally many b) that allows this requirement to be fulfilled. Next to understand (6), suppose that $\langle \mathcal{U}_\alpha : \alpha < \omega_1 \rangle$ has been built and that you are building \mathcal{U}_{ω_1} . At some stage you have determined that $\langle \text{set}(d_n) : n < \omega \rangle \subset \mathcal{U}_{\omega_1}$, for some decreasing sequence of conditions $\langle d_n : n < \omega \rangle \subset \mathbb{P}$. You have also determined the sequence $\langle \pi_{\omega_1,n} : n < \omega \rangle$. In particular $\forall n, m < \omega [\pi''_{\omega_1,n} \text{set}(d_m) \in \mathcal{U}_n]$, and each $\pi_{\omega_1,n}$ has the right form on some d_m . Now you would like to find a $d^* \in \mathbb{P}$ that is below all of the d_n . You would also like to determine $\pi_{\omega_1,\omega}$. But you must ensure that $\pi''_{\omega_1,\omega} \text{set}(d^*) \in \mathcal{U}_\omega$, that $\pi_{\omega_1,\omega}$ has the appropriate form on d^* , and that all of the $\pi_{\omega_1,n}$ commute through $\pi_{\omega_1,\omega}$. Clause (6) says that \mathcal{U}_ω anticipated this requirement and that there is a $b \in \mathcal{U}_\omega$ (in fact cofinally many b) enabling you to find such a d^* and $\pi_{\omega_1,\omega}$.

Definition 2.10. Let $\delta \leq \omega_2$. We call $\langle \langle c_i^\alpha : i < \mathfrak{c} \wedge \alpha < \delta \rangle, \langle \pi_{\beta,\alpha} : \alpha \leq \beta < \delta \rangle \rangle$ δ -generic if and only if²:

- (1) for every $\alpha < \delta$, $\langle c_i^\alpha : i < \mathfrak{c} \rangle$ is a decreasing sequence in \mathbb{P} ; for every $\alpha \leq \beta < \delta$, $\pi_{\beta,\alpha} \in \omega^\omega$;
- (2) for every $\alpha < \delta$, $\mathcal{U}_\alpha = \{a \in \mathcal{P}(\omega) : \exists i < \mathfrak{c} [\text{set}(c_i^\alpha) \subset^* a]\}$ is an ultrafilter on ω and it is a rapid P-point (we say that \mathcal{U}_α is *generated by* $\langle c_i^\alpha : i < \mathfrak{c} \rangle$);
- (3) for every $\alpha < \beta < \delta$, every normal triple $\langle \pi_1, \psi_1, b_1 \rangle$ and every $d \leq b_1$ if $\pi''_1 \text{set}(d) \in \mathcal{U}_\alpha$, then for every $a \in \mathcal{U}_\beta$ there is $b \in \mathcal{U}_\beta$ such that $b \subset^* a$ and that there are $\pi, \psi \in \omega^\omega$ and $d^* \leq_0 d$ so that $\langle \pi, \psi, d^* \rangle$ is a normal triple, $\pi'' \text{set}(d^*) = b$ and $\forall k \in \text{set}(d^*) [\pi_1(k) = \pi_{\beta,\alpha}(\pi(k))]$.
- (4) if $\alpha < \beta < \delta$, then $\mathcal{U}_\beta \not\leq_T \mathcal{U}_\alpha$.
- (5) for every $\alpha < \delta$, $\pi_{\alpha,\alpha} = \text{id}$ and:
 - (a) $\forall \alpha \leq \beta < \delta \forall i < \mathfrak{c} [\pi''_{\beta,\alpha} \text{set}(c_i^\beta) \in \mathcal{U}_\alpha]$;
 - (b) $\forall \alpha \leq \beta \leq \gamma < \delta \exists i < \mathfrak{c} \forall^\infty k \in \text{set}(c_i^\gamma) [\pi_{\gamma,\alpha}(k) = \pi_{\beta,\alpha}(\pi_{\gamma,\beta}(k))]$;
 - (c) for $\alpha < \beta < \delta$ there are $i < \mathfrak{c}$, $b_{\beta,\alpha} \in \mathbb{P}$ and $\psi_{\beta,\alpha} \in \omega^\omega$ such that $\langle \pi_{\beta,\alpha}, \psi_{\beta,\alpha}, b_{\beta,\alpha} \rangle$ is a normal triple and $c_i^\beta \leq b_{\beta,\alpha}$;
- (6) if $\mu < \delta$ is a limit ordinal such that $\text{cof}(\mu) = \omega$, $X \subset \mu$ is a countable set such that $\sup(X) = \mu$, $\langle d_j : j < \omega \rangle$ is a decreasing sequence of conditions in \mathbb{P} , and $\langle \pi_\alpha : \alpha \in X \rangle$ is a sequence of maps in ω^ω such that:
 - (a) $\forall \alpha \in X \forall j < \omega [\pi''_\alpha \text{set}(d_j) \in \mathcal{U}_\alpha]$;
 - (b) $\forall \alpha, \beta \in X [\alpha \leq \beta \Rightarrow \exists j < \omega \forall^\infty k \in \text{set}(d_j) [\pi_\alpha(k) = \pi_{\beta,\alpha}(\pi_\beta(k))]]$;
 - (c) for all $\alpha \in X$ there are $j < \omega$, $b_\alpha \in \mathbb{P}$ and $\psi_\alpha \in \omega^\omega$ such that $\langle \pi_\alpha, \psi_\alpha, b_\alpha \rangle$ is a normal triple and $d_j \leq b_\alpha$;
then the set of all $i^* < \mathfrak{c}$ such that there are $d^* \in \mathbb{P}$ and $\pi, \psi \in \omega^\omega$ satisfying:
 - (d) $\forall j < \omega [d^* \leq d_j]$ and $\text{set}(c_{i^*}^\mu) = \pi'' \text{set}(d^*)$;
 - (e) $\forall \alpha \in X \forall^\infty k \in \text{set}(d^*) [\pi_\alpha(k) = \pi_{\mu,\alpha}(\pi(k))]$;
 - (f) $\langle \pi, \psi, d^* \rangle$ is a normal triple;
is cofinal in \mathfrak{c} ;

When CH is replaced with MA, the notion of a δ -generic sequence would be defined for every $\delta \leq \mathfrak{c}^+$. Clause (6) would need to be strengthened by allowing μ to be any limit ordinal such that $\text{cof}(\mu) < \mathfrak{c}$ and by allowing the decreasing sequence of conditions in \mathbb{P} to be of length $\text{cof}(\mu)$.

Remark 2.11. Suppose $\delta \leq \omega_2$ is some limit ordinal. We point out that if

$$S = \langle \langle c_i^\alpha : i < \mathfrak{c} \wedge \alpha < \delta \rangle, \langle \pi_{\beta,\alpha} : \alpha \leq \beta < \delta \rangle \rangle$$

²Raghavan and Verner [10] have proved that Clauses (3) and (6) below are redundant. In other words, Clauses (1), (2), (4), and (5) automatically imply (3) and (6).

is any sequence such that $S \upharpoonright \xi = \langle \langle c_i^\alpha : i < \mathfrak{c} \wedge \alpha < \xi \rangle, \langle \pi_{\beta, \alpha} : \alpha \leq \beta < \xi \rangle \rangle$ is ξ -generic for every ordinal $\xi < \delta$, then S is δ -generic. To see this we check conditions (1)–(6) of Definition 2.10. Conditions (1) and (2) are true because for a fixed $\alpha < \delta$ we can pick ξ so that $\alpha < \xi < \delta$. Then $S \upharpoonright \xi$ witnesses that \mathcal{U}_α and $\langle c_i^\alpha : i < \mathfrak{c} \rangle$ are as needed. For (3), (4) and (5c) take $\alpha < \beta < \delta$. There is ξ such that $\beta < \xi < \delta$ and $S \upharpoonright \xi$ witnesses (3), (4) and (5c). For (5a) and (5b) take $\alpha \leq \beta \leq \gamma < \delta$. Again there is ξ so that $\gamma < \xi < \delta$ and $S \upharpoonright \xi$ witnesses both (5a) and (5b). We still have to prove condition (6), so assume that all the objects are given as in (6). In this case we also pick ξ such that $\mu < \xi < \delta$. Then $S \upharpoonright \xi$ already has all the information about the assumed objects. So $S \upharpoonright \xi$ knows that the set of $i^* < \mathfrak{c}$ such that $c_{i^*}^\mu$ has the required properties is cofinal in \mathfrak{c} which implies that (6) is also satisfied in S .

3. MAIN LEMMAS

In this section we prove several crucial lemmas that will be used in Section 4 for proving things about the partial order \mathbb{Q}^δ to be defined there.

Definition 3.1. Let $\langle L, \prec \rangle$ be a finite linear order. For each $i < |L|$ let $L(i, \prec)$ denote the i th element of $\langle L, \prec \rangle$. More formally, if $\langle L, \prec \rangle$ is a finite linear order, then there is a unique order isomorphism $o : |L| \rightarrow L$ and $L(i, \prec) = o(i)$ ($i < |L|$).

Lemma 3.2 will be used to prove Lemmas 3.4 and 3.8. It essentially says that the sets D_m can be broken into intervals of the form $D_m[K_{m,n}, K_{m,n+1})$, for $m \leq n < \omega$, in such a way that whenever $m < m' \leq n$, the image of $D_{m'}[K_{m',n}, K_{m',n+1})$ under $\pi_{m',m}$ comes after everything in the interval $D_m[K_{m,n-1}, K_{m,n})$. The use of the elementary submodel M is only for convenience. We just need a way of saying that each D_n diagonalizes a “large enough” collection of sets from \mathcal{U}_n . $M \cap \mathcal{U}_n$ is a convenient way to specify this collection. The use of Lemma 3.2 simplifies the presentation of the proofs of Lemmas 3.4 and 3.8. There is no direct analogue of Lemma 3.2 when CH is replaced by MA. So the proofs of the analogues of Lemmas 3.4 and 3.8 under MA will be less elementary.

Lemma 3.2. Let $\langle \mathcal{U}_n : n < \omega \rangle$ be a sequence of distinct rapid P-points. Assume that $\bar{\pi} = \langle \pi_{m,n} : n \leq m < \omega \rangle$ is a sequence of maps in ω^ω such that $\pi_{n,n} = \text{id}$ ($n < \omega$) and:

- (1) $\forall n \leq m < \omega \forall a \in \mathcal{U}_m [\pi_{m,n}'' a \in \mathcal{U}_n]$;
- (2) $\forall n \leq m \leq k < \omega \exists a \in \mathcal{U}_k \forall l \in a [\pi_{k,n}(l) = \pi_{m,n}(\pi_{k,m}(l))]$;
- (3) $\forall n \leq m < \omega \exists a \in \mathcal{U}_m \forall x, y \in a [x \leq y \Rightarrow \pi_{m,n}(x) \leq \pi_{m,n}(y)]$.

Let $\langle E_n : n < \omega \rangle$ be a sequence such that $E_n \in \mathcal{U}_n$ ($n < \omega$). Suppose also that $f \in \omega^\omega$ is increasing and that M is a countable elementary submodel of $H_{(2^{\mathfrak{c}})^+}$ containing $\langle \mathcal{U}_n : n < \omega \rangle$, $\langle E_n : n < \omega \rangle$, $\bar{\pi}$ and the map f . If $\bar{D} = \langle D_n : n < \omega \rangle$ is a sequence such that $D_n \in \mathcal{U}_n$ ($n < \omega$) and $D_n \subset^* A$, for every $A \in \mathcal{U}_n \cap M$, then there are sequences $\langle C_n : n < \omega \rangle \in M$, $\langle F_n : n < \omega \rangle \in M$, $\bar{K} = \langle K_{m,n} : m \leq n < \omega \rangle \subset \omega$ and $g' \in \omega^\omega$ such that for every $n < \omega$ we have $\forall m \leq n [K_{m,n} > 0]$, $C_n, F_n \in \mathcal{U}_n$, $C_n \subset E_n$, $F_n = C_n \cap \pi_{n+1,n}'' C_{n+1}$, $\forall m \leq m' \leq n \forall v \in C_n [\pi_{n,m}(v) = \pi_{m',m}(\pi_{n,m'}(v))]$, $\forall m \leq n \forall v, v' \in C_n [v \leq v' \Rightarrow \pi_{n,m}(v) \leq \pi_{n,m}(v')]$, and letting $\rho = \langle \bar{D}, \bar{K}, \bar{\pi} \rangle$ (see Remark 2.1):

- (4) $\forall m \leq n \exists z \in F_n \cap F_n(g'(n)) [\pi_{n,m}(z) = D_m(K_{m,n} - 1)]$ and if $n > 0$ then $\forall m < n [K_{m,n} > K_{m,n-1}]$;
- (5) $\forall m \leq n [D_m[K_{m,n}] \subset \pi_{n,m}'' (F_n \setminus F_n(g'(n)))]$;
- (6) $\forall m \leq n \forall v \in C_n \setminus F_n(g'(n)) [\pi_{n,m}(v) > D_m(K_{m,n} - 1)]$;
- (7) $\forall m < n [D_n[K_{m,n}] \cap D_m = 0]$ and if $n > 0$, then for every $x \in \Delta_{n-1}^\rho$, there is a unique $m < n$ such that $x \in D_m[K_{m,n-1}, K_{m,n})$;
- (8) if $n > 0$, then define \prec_{n-1} to be the collection of all $\langle x, y \rangle$ such that $x, y \in \Delta_{n-1}^\rho$ and $\max \{z \in F_{n-1} : \pi_{n-1,m}(z) = x\} < \max \{z \in F_{n-1} : \pi_{n-1,m'}(z) = y\} -$

where $m, m' < n$ are unique with the property that $x \in D_m[K_{m,n-1}, K_{m,n}]$ and $y \in D_{m'}[K_{m',n-1}, K_{m',n}]$; then \prec_{n-1} is a linear order on Δ_{n-1}^ρ ;

(9) $2g'(n) \geq L_n^\rho$;

(10) if $n > 0$, then define the following notation: let $R_{n-1} = |\Delta_{n-1}^\rho|$ and for each $j < R_{n-1}$, let x_j^{n-1} be $\Delta_{n-1}^\rho(j, \prec_{n-1})$, let $m(n-1, j)$ be the unique $m < n$ such that $x_j^{n-1} \in D_m[K_{m,n-1}, K_{m,n}]$, and let

$$z_j^{n-1} = \max \{z \in F_{n-1} : \pi_{n-1, m(n-1, j)}(z) = x_j^{n-1}\};$$

then there exists $l \geq f(L_{n-1}^\rho + j)$ such that $z_j^{n-1} = E_{n-1}(l)$.

Proof. First pick a sequence $\bar{A} = \langle A_{n,m,k} : n \leq m \leq k < \omega \rangle \in M$ such that $A_{n,m,k} \in \mathcal{U}_k$ for each $n \leq m \leq k < \omega$ and $\pi_{k,n}(v) = \pi_{m,n}(\pi_{k,m}(v))$ for every $v \in A_{n,m,k}$. Similarly, pick a sequence $\bar{B} = \langle B_{n,m} : n \leq m < \omega \rangle \in M$ so that $B_{n,m} \in \mathcal{U}_m$, for every $n \leq m < \omega$ and $\forall x, y \in B_{n,m} [x \leq y \Rightarrow \pi_{m,n}(x) \leq \pi_{m,n}(y)]$. Let us now define sequence $\langle C'_k : k < \omega \rangle$ so that for every $k < \omega$ we have $C'_k = E_k \cap \left(\bigcap_{n \leq m \leq k} A_{n,m,k} \right) \cap \left(\bigcap_{m \leq k} B_{m,k} \right)$. Note that $C'_k \in \mathcal{U}_k$ and that $C'_k \subset E_k$ for every $k < \omega$. Moreover, $\langle C'_k : k < \omega \rangle \in M$. Next, choose $\bar{C} = \langle C_k : k < \omega \rangle \in M$ so that for $k < \omega$: $C_k \in \mathcal{U}_k$, $C_k \subset C'_k$ and $C_k(n) \geq E_k(f(2n))$ for every $n < \omega$. Let F_n be the set $C_n \cap \pi''_{n+1,n} C_{n+1}$ ($n < \omega$). Note that $\langle F_n : n < \omega \rangle$ belongs to M . Note also that for each $m \leq m' \leq n < \omega$, $C_n \subset A_{m,m',n}$. So $\forall v \in C_n [\pi_{n,m}(v) = \pi_{m',m}(\pi_{n,m'}(v))]$ as required in the statement of the lemma. Moreover if $m \leq n$, then $C_n \subset B_{m,n}$, implying that $\forall v, v' \in C_n [v \leq v' \Rightarrow \pi_{n,m}(v) \leq \pi_{n,m}(v')]$ as needed. Then by the definition of sets D_n ($n < \omega$) we have $D_n \subset^* \pi''_{m,n} F_m$ for $m \geq n$. Also, for $n < \omega$, let Y_n be minimal so that $\forall m < n [D_n[Y_n] \cap D_m = 0]$.

Now that we have chosen sets C_n and F_n ($n < \omega$) we construct, by induction on n , numbers $K_{m,n}$ and $g'(n)$ ($m \leq n < \omega$). First let l' be the least number such that $D_0[l'] \subset F_0$ and let $K_{0,0} = l' + 1$. Then define $g'(0)$ to be the least l so that $F_0(l) \geq D_0(K_{0,0})$. Note that properties (4)–(10) hold for $n = 0$. Now assume that for every $m \leq m' \leq n$ we have defined numbers $K_{m,m'}$ and $g'(m')$. We will define $K_{m,n+1}$ ($m \leq n+1$) and $g'(n+1)$. So for every $m \leq n$ let $X_{m,n+1}$ be the least number such that $X_{m,n+1} \geq K_{m,n}$ and $D_m[X_{m,n+1}] \subset \pi''_{n+1,m} F_{n+1}$, while $X_{n+1,n+1}$ is defined to be minimal such that $D_{n+1}[X_{n+1,n+1}] \subset F_{n+1} = \pi''_{n+1,n+1} F_{n+1}$ and $X_{n+1,n+1} \geq Y_{n+1}$. Put $x^* = L_n^\rho + \sum_{m \leq n} (X_{m,n+1} - K_{m,n})$. Now define $K_{0,n+1}$ to be the minimal l such that $l > X_{0,n+1} + x^*$ and that $D_0(l-1) \in \bigcap_{m \leq n+1} \pi''_{m,0} (D_m[X_{m,n+1}])$. Next define $g'(n+1)$ to be the minimal $l \in \omega$ with $\pi_{n+1,0}(F_{n+1}(l)) \geq D_0(K_{0,n+1})$. Observe that if $v \in F_{n+1} \cap F_{n+1}(g'(n+1))$, then $\pi_{n+1,0}(v) < D_0(K_{0,n+1})$. Also if $v \in F_{n+1}(g'(n+1))$, then $\pi_{n+1,0}(v) \geq \pi_{n+1,0}(F_{n+1}(g'(n+1))) \geq D_0(K_{0,n+1})$. Put $G = F_{n+1}(g'(n+1))$ for convenience. Now for $0 < m \leq n+1$ define $K_{m,n+1}$ as the minimal $l \in \omega$ so that $D_m[l] \subset \pi''_{n+1,m} G$. We remark that $K_{0,n+1}$ is also minimal such that $D_0[K_{0,n+1}] \subset \pi''_{n+1,0} G$. To see this, take any $w \in D_0[K_{0,n+1}]$. Since $K_{0,n+1} \geq X_{0,n+1}$, there exists $v \in F_{n+1}$ with $\pi_{n+1,0}(v) = w$. By the first observation above, $v \notin F_{n+1}(g'(n+1))$. Hence $v \in G$ showing that $D_0[K_{0,n+1}] \subset \pi''_{n+1,0} G$. By the second observation above, there is no $v \in G$ with $\pi_{n+1,0}(v) = D_0(K_{0,n+1} - 1)$. Hence there is no $l < K_{0,n+1}$ satisfying $D_0[l] \subset \pi''_{n+1,0} G$.

Now we prove that (4)–(10) are fulfilled for $n+1$. We begin with the second part of (4). Fix $m \leq n+1$. By the definition of $K_{0,n+1}$, there exists $u \in D_m[X_{m,n+1}]$ such that $\pi_{m,0}(u) = D_0(K_{0,n+1} - 1)$. We claim $u < D_m(K_{m,n+1})$. Suppose not. Then $u \in D_m[K_{m,n+1}]$, and so $u = \pi_{n+1,m}(v)$, for some $v \in G$. However $\pi_{n+1,0}(v) = \pi_{m,0}(\pi_{n+1,m}(v)) = \pi_{m,0}(u) = D_0(K_{0,n+1} - 1)$, contradicting an observation of the previous paragraph. Thus $u < D_m(K_{m,n+1})$, showing that $X_{m,n+1} < K_{m,n+1}$, for all $m \leq n+1$.

Before proving the rest of (4), (5), and (6) we make some useful observations. Put $y_m = D_m(K_{m,n+1} - 1)$, for each $m \leq n+1$. Let $m \leq n+1$ be fixed. First since $K_{m,n+1}$ is minimal so that $D_m[K_{m,n+1}] \subset \pi''_{n+1,m}G$, there is no $v \in G$ with $\pi_{n+1,m}(v) = y_m$. Next let $u \in D_m[K_{m,n+1}]$. Then there exists $v \in G$ with $u = \pi_{n+1,m}(v)$, and $\pi_{m,0}(u) = \pi_{m,0}(\pi_{n+1,m}(v)) = \pi_{n+1,0}(v) \geq D_0(K_{0,n+1})$. Thus $\pi_{m,0}(u) \geq D_0(K_{0,n+1})$, for every $u \in D_m[K_{m,n+1}]$. For the final observation, consider some $v \in F_{n+1} \cap F_{n+1}(g'(n+1))$. As pointed out before, $\pi_{n+1,0}(v) < D_0(K_{0,n+1})$. Now let $u = \pi_{n+1,m}(v)$. Then $\pi_{m,0}(u) = \pi_{n+1,0}(v) < D_0(K_{0,n+1})$. Applying the previous observation to u , we conclude that $\pi_{n+1,m}(v) \notin D_m[K_{m,n+1}]$, for every $v \in F_{n+1} \cap F_{n+1}(g'(n+1))$.

Now the rest of (4), (5), and (6) easily follow from the three observations in the previous paragraph. For the first part of (4), $y_m \in D_m[X_{m,n+1}]$, and so there is $v_m \in F_{n+1}$ with $y_m = \pi_{n+1,m}(v_m)$. By the first observation, $v_m \notin G$. Hence $v_m \in F_{n+1} \cap F_{n+1}(g'(n+1))$, as needed. For (5), let $u \in D_m[K_{m,n+1}]$. Then $u \in D_m[X_{m,n+1}]$ and so there is $v \in F_{n+1}$ with $\pi_{n+1,m}(v) = u$. By the third observation, $v \notin F_{n+1}(g'(n+1))$, as required. For (6), first note that since $v_m, F_{n+1}(g'(n+1)) \in B_{m,n+1}$, $\pi_{n+1,m}(F_{n+1}(g'(n+1))) \geq y_m$, and by the first observation, $\pi_{n+1,m}(F_{n+1}(g'(n+1))) > y_m$. Now let $v \in C_{n+1}$. Then $v \in B_{m,n+1}$ and if $v \geq F_{n+1}(g'(n+1))$, then $\pi_{n+1,m}(v) > y_m$, implying (6).

For (7) we have $Y_{n+1} \leq X_{n+1,n+1} < K_{n+1,n+1}$, and so $D_{n+1}[K_{n+1,n+1}] \cap D_m = 0$, for all $m < n+1$. The second part of (7) easily follows from the definition of Δ_n^ρ and from the induction hypotheses.

For (8), first consider any $x \in \Delta_n^\rho$. By (7) applied to $n+1$, let $m < n+1$ be unique so that $x \in D_m[K_{m,n}, K_{m,n+1}]$. By (5) applied to n , there is $z \in F_n$ with $\pi_{n,m}(z) = x$. Also $F_n \subset C_n$ and so $\pi_{n,m}$ is finite-to-one on F_n . Therefore $\max\{z \in F_n : \pi_{n,m}(z) = x\}$ is well-defined. Next it is clear that \prec_n is transitive and irreflexive. We check that it is total. Let $x, y \in \Delta_n^\rho$ and let $m, m' < n+1$ be unique so that $x \in D_m[K_{m,n}, K_{m,n+1}]$ and $y \in D_{m'}[K_{m',n}, K_{m',n+1}]$. We may assume $m \leq m'$. If x and y are incomparable under \prec_n , then there exists $z \in F_n$ so that $\pi_{n,m}(z) = x$ and $\pi_{n,m'}(z) = y$. As $F_n \subset C_n$, $\pi_{m',m}(y) = \pi_{m',m}(\pi_{n,m'}(z)) = \pi_{n,m}(z) = x$. If $m < m'$, then this contradicts the fact that $x \in H_{m,n}^\rho$. Therefore $m = m'$, and since $\pi_{m',m} = \text{id}$, $x = y$, implying comparability.

Now we check (9). For each $m < n+1$ define $H_{m,n,0}^\rho = H_{m,n}^\rho \cap D_m[K_{m,n}, X_{m,n+1}]$ and $H_{m,n,1}^\rho = H_{m,n}^\rho \cap D_m[X_{m,n+1}, K_{m,n+1}]$. Define $x_0 = \left| \bigcup_{m \leq n} H_{m,n,0}^\rho \right|$ and $x_1 = \left| \bigcup_{m \leq n} H_{m,n,1}^\rho \right|$. It is clear that $H_{m,n}^\rho = H_{m,n,0}^\rho \cup H_{m,n,1}^\rho$ and that $L_{n+1}^\rho \leq L_n^\rho + x_0 + x_1$. Also $L_n^\rho + x_0 \leq x^*$. So to prove (9) it is enough to show both $x^* \leq g'(n+1)$ and $x_1 \leq g'(n+1)$. For the first inequality, note that $|D_0[X_{0,n+1}, K_{0,n+1}]| \geq x^*$. For each $u \in D_0[X_{0,n+1}, K_{0,n+1}]$, there exists $v \in F_{n+1}$ with $\pi_{n+1,0}(v) = u$. By (6) applied to $n+1$, $v \in F_{n+1}(g'(n+1))$. It follows that $g'(n+1) \geq x^*$. For the second inequality, note first that for each $m \leq n$ and $u \in H_{m,n,1}^\rho$ we get a $v \in F_{n+1} \cap F_{n+1}(g'(n+1))$ with $\pi_{n+1,m}(v) = u$ by applying the same argument. Now suppose $u \neq u'$, $m, m' \leq n$, $u \in H_{m,n,1}^\rho$, $u' \in H_{m',n,1}^\rho$, $v, v' \in F_{n+1} \cap F_{n+1}(g'(n+1))$, $\pi_{n+1,m}(v) = u$, and $\pi_{n+1,m'}(v') = u'$. We would like to see that $v \neq v'$. Suppose not. We may assume $m \leq m'$. Since $v \in C_{n+1}$, $u = \pi_{n+1,m}(v) = \pi_{m',m}(\pi_{n+1,m'}(v)) = \pi_{m',m}(u')$. If $m < m'$, then this contradicts the fact that $u \in H_{m,n}^\rho$. Hence $m = m'$, whence $u = u'$. This is a contradiction which shows that $v \neq v'$. It follows that $g'(n+1) \geq x_1$ as needed.

Finally we come to (10). Fix $j < R_n$. By (5) applied to n , $z_j^n = F_n(t_j^n)$ for some $t_j^n \geq g'(n)$. Since \prec_n is a linear order $t_j^n \geq g'(n) + j$. By (9) applied to n , $2t_j^n \geq 2g'(n) + 2j \geq L_n^\rho + j$. Now $F_n \subset C_n \subset C'_n \subset E_n$ and $F_n(t_j^n) \geq C_n(t_j^n) \geq E_n(f(2t_j^n)) \geq E_n(f(L_n^\rho + j))$ because f is an increasing function. It follows that $z_j^n = E_n(l_j^n)$ for some $l_j^n \geq f(L_n^\rho + j)$, as needed. \square

Remark 3.3. Note that for each $n < \omega$, $H_{n,n}^\rho = D_n[K_{n,n}, K_{n,n+1}]$; so $\Delta_n^\rho \neq 0$, and so $L_{n+1}^\rho > L_n^\rho$. Note also that $z_j^n \geq F_n(g'(n))$, for each $n < \omega$ and $j < R_n$.

Lemma 3.4 will play an important role throughout the next section. It is essential to the proof that \mathbb{Q}^δ , which will be defined in Definition 4.1, is countably closed. It is also used in ensuring that \mathcal{U}_δ is a rapid ultrafilter and that \mathcal{U}_δ satisfies (3) and (6) of Definition 2.10.

Lemma 3.4. *Assume that $\delta < \omega_2$, $\text{cof}(\delta) = \omega$, $f \in \omega^\omega$ is increasing, $X \subset \delta$ is such that $\text{sup}(X) = \delta$ and:*

- (1) *the sequence $S = \langle \langle c_i^\alpha : i < \mathfrak{c} \wedge \alpha < \delta \rangle, \langle \pi_{\beta, \alpha} : \alpha \leq \beta < \delta \rangle \rangle$ is δ -generic;*
- (2) *there are $e \in \mathbb{P}$ and mappings $\langle \pi_{\delta, \alpha} : \alpha \in X \rangle$ such that:*
 - (a) $\forall \alpha \in X [\pi''_{\delta, \alpha} \text{set}(e) \in \mathcal{U}_\alpha]$;
 - (b) $\forall \alpha, \beta \in X [\alpha \leq \beta \Rightarrow \forall^\infty k \in \text{set}(e) [\pi_{\delta, \alpha}(k) = \pi_{\beta, \alpha}(\pi_{\delta, \beta}(k))]]$;
 - (c) *for all $\alpha \in X$ there are $b_{\delta, \alpha}$ and $\psi_{\delta, \alpha}$ such that $\langle \pi_{\delta, \alpha}, \psi_{\delta, \alpha}, b_{\delta, \alpha} \rangle$ is a normal triple and $e \leq b_{\delta, \alpha}$;*
- (3) *there is a decreasing sequence $\langle d_j : j < \omega \rangle$ of elements of \mathbb{P} and a sequence of mappings $\langle \pi_\alpha : \alpha \in X \rangle$ such that:*
 - (a) $\forall \alpha \in X \forall j < \omega [\pi''_\alpha \text{set}(d_j) \in \mathcal{U}_\alpha]$;
 - (b) $\forall \alpha, \beta \in X [\alpha \leq \beta \Rightarrow \exists j < \omega \forall^\infty k \in \text{set}(d_j) [\pi_\alpha(k) = \pi_{\beta, \alpha}(\pi_\beta(k))]]$;
 - (c) *for all $\alpha \in X$ there are $j < \omega$, ψ_α and b_α such that $\langle \pi_\alpha, \psi_\alpha, b_\alpha \rangle$ is a normal triple and $d_j \leq b_\alpha$.*

Then there are $d^, e^* \in \mathbb{P}$ and $\pi : \omega \rightarrow \omega$ such that:*

- (4) $\forall n < \omega \exists m \geq f(n) [e^*(n) \subset e(m)]$;
- (5) $\forall j < \omega [d^* \leq d_j]$ and $\text{set}(e^*) = \pi'' \text{set}(d^*)$;
- (6) $\forall \alpha \in X \forall^\infty k \in \text{set}(d^*) [\pi_\alpha(k) = \pi_{\delta, \alpha}(\pi(k))]$;
- (7) $\forall \alpha \in X [\pi''_{\delta, \alpha} \text{set}(e^*) \in \mathcal{U}_\alpha]$;
- (8) *there is $\psi \in \omega^\omega$ for which $\langle \pi, \psi, d^* \rangle$ is a normal triple.*

Proof. Let $\langle \delta_n : n < \omega \rangle \subset X$ be an increasing and cofinal sequence in δ . For $m \leq n < \omega$ choose $j_{m, n}$ and $L^d(m, n)$ so that $\forall k \in \text{set}(d_{j_{m, n}}) \llbracket L^d(m, n) \rrbracket [\pi_{\delta_m}(k) = \pi_{\delta_n, \delta_m}(\pi_{\delta_n}(k))]$. For every $n < \omega$ pick $j(n)$, b_{δ_n} and ψ_{δ_n} such that $\langle \pi_{\delta_n}, \psi_{\delta_n}, b_{\delta_n} \rangle$ is a normal triple and $d_{j(n)} \leq b_{\delta_n}$ holds by using (3c). Let $K^d(n)$ be minimal such that $d_{j(n)} \leq_{K^d(n)} b_{\delta_n}$. Define a strictly increasing sequence $\langle j_N : N < \omega \rangle$ by setting $j_N = \max(\{j(N)\} \cup \{j_k + 1 : k < N\} \cup \{j_{m, N} : m \leq N\})$. Let $Q^d(N)$ be minimal such that $d_{j_N} \leq_{Q^d(N)} d_{j(N)}$, $\forall k < N [d_{j_N} \leq_{Q^d(N)} d_{j_k}]$ and $\forall m \leq N [d_{j_N} \leq_{Q^d(N)} d_{j_{m, N}}]$. Define $M_N^d = \max(\{K^d(N), Q^d(N)\} \cup \{L^d(m, N) : m \leq N\} \cup \{M_k^d : k < N\})$.

For each $n < \omega$ let $K^e(n)$ be minimal such that $e \leq_{K^e(n)} b_{\delta_n}$. For $m \leq n < \omega$ let $L^e(m, n)$ be minimal such that $\forall k \in \text{set}(e) \llbracket L^e(m, n) \rrbracket [\pi_{\delta_n, \delta_m}(k) = \pi_{\delta_n, \delta_m}(\pi_{\delta_n}(k))]$. For $N < \omega$ let $M_N^e = \max(\{K^e(N)\} \cup \{L^e(m, N) : m \leq N\} \cup \{M_k^e : k < N\})$.

The proof of the following claim is simple so we leave it to the reader.

Claim 3.5. *Let $N < \omega$. The following hold:*

- (1) $\forall m \leq n \leq N \forall k \in \text{set}(e) \llbracket M_N^e \rrbracket [\pi_{\delta_n, \delta_m}(k) = \pi_{\delta_n, \delta_m}(\pi_{\delta_n}(k))]$;
- (2) $\forall m \leq n \leq N \forall k \in \text{set}(d_{j_N}) \llbracket M_N^d \rrbracket [\pi_{\delta_m}(k) = \pi_{\delta_n, \delta_m}(\pi_{\delta_n}(k))]$;
- (3) $\forall n \leq N \forall k, l \in \text{set}(e) \llbracket M_N^e \rrbracket [k \leq l \Rightarrow \pi_{\delta_n}(k) \leq \pi_{\delta_n}(l)]$;
- (4) $\forall n \leq N \forall k, l \in \text{set}(d_{j_N}) \llbracket M_N^d \rrbracket [k \leq l \Rightarrow \pi_{\delta_n}(k) \leq \pi_{\delta_n}(l)]$.

For every $n < \omega$ let $E_n = \pi''_{\delta_n} \text{set}(e) \llbracket M_n^e \rrbracket \cap \pi''_{\delta_n} \text{set}(d_{j_n}) \llbracket M_n^d \rrbracket \in \mathcal{U}_{\delta_n}$. Let M be a countable elementary submodel of $H_{(2^\epsilon)^+}$ containing S, δ, f , and sequences $\langle E_n : n < \omega \rangle$ and $\langle \delta_n : n < \omega \rangle$. For $n < \omega$ choose sets $D_n \in \mathcal{U}_{\delta_n}$ as follows: $D_n \subset^* A$ for every $A \in \mathcal{U}_{\delta_n} \cap M$. Note that these sets exist because \mathcal{U}_{δ_n} is a P-point and M is countable. Define $g \in \omega^\omega$ by $g(n) = \max\{f(n), t(n), s(n+1)\}$, for each $n \in \omega$. Note $g \in M$ and that g is increasing. Now Lemma 3.2 applies to the sequences $\langle \mathcal{U}_{\delta_n} : n < \omega \rangle$, $\bar{\pi} = \langle \pi_{\delta_n, \delta_m} : m \leq n < \omega \rangle$, $\langle E_n : n < \omega \rangle$, $\bar{D} = \langle D_n : n < \omega \rangle$ and the function g . Let $\langle C_n : n < \omega \rangle$, $\langle F_n : n < \omega \rangle$, $\bar{K} = \langle K_{m, n} : m \leq n < \omega \rangle$ and $\langle g'(n) : n < \omega \rangle$ be as in the conclusion of Lemma 3.2. We denote $\rho = \langle \bar{D}, \bar{K}, \bar{\pi} \rangle$, numbers $m(n, j)$, numbers R_n and numbers z_j^n as in the conclusion of Lemma 3.2.

At this point, for every $n < \omega$, we define set $I_n = \{m < \omega : L_n^\rho \leq m < L_{n+1}^\rho\}$. Clearly, $\{I_n : n < \omega\}$ is a partition of ω . We also have $I_n = \{L_n^\rho + j : j < R_n\}$. So every $k < \omega$ is of the form $L_n^\rho + j$ for some $n < \omega$ and $j < R_n$. For each $n < \omega$ and $j < R_n$, (10) of Lemma 3.2 implies that $z_j^n = E_n(l_j^n)$ for some $l_j^n \geq g(L_n^\rho + j)$. Now for a fixed $n < \omega$, $\langle \pi_{\delta, \delta_n}, \psi_{\delta, \delta_n}, b_{\delta, \delta_n} \rangle$ and $\langle \pi_{\delta_n}, \psi_{\delta_n}, b_{\delta_n} \rangle$ are normal triples, $e \leq_{M_n^e} b_{\delta, \delta_n}$, $d_{j_n} \leq_{M_n^d} b_{\delta_n}$, $E_n \subset \pi_{\delta, \delta_n}'' \text{set}(e) \llbracket M_n^e \rrbracket$, and $E_n \subset \pi_{\delta_n}'' \text{set}(d_{j_n}) \llbracket M_n^d \rrbracket$. So Lemma 2.9 applies and implies that for each $l < \omega$, $\zeta_l = \max\{m < \omega : \pi_{\delta, \delta_n}'' e(m) = \{E_n(l)\}\}$ and $\mathfrak{z}_l = \max\{m < \omega : \pi_{\delta_n}'' d_{j_n}(m) = \{E_n(l)\}\}$ are well-defined, that $\zeta_l < \zeta_{l+1}$, $\mathfrak{z}_l < \mathfrak{z}_{l+1}$, and that $\zeta_l \geq M_n^e$, $\mathfrak{z}_l \geq M_n^d$. Now for each $j < R_n$, define $\zeta_j^n = \zeta_{l_j^n}$ and $\mathfrak{z}_j^n = \mathfrak{z}_{l_j^n}$. It follows that $\zeta_j^n, \mathfrak{z}_j^n \geq g(L_n^\rho + j)$. Also if $j < j+1 < R_n$, then $z_j^n < z_{j+1}^n$, and so $\zeta_j^n < \zeta_{j+1}^n$ and $\mathfrak{z}_j^n < \mathfrak{z}_{j+1}^n$. Hence for $j < R_n$, $|e(\zeta_j^n)| \geq g(L_n^\rho + j) \geq L_n^\rho + j + 1$ and $|d_{j_n}(\mathfrak{z}_j^n)| \geq g(L_n^\rho + j) \geq t(L_n^\rho + j)$. Now fix n . For $k < \omega$ write $k = L_n^\rho + j$ and pick arbitrary $e^*(k) \in [e(\zeta_j^n)]^{k+1}$. Note that this choice of $e^*(k)$ ($k < \omega$) ensures that $e^*(k) \subset e(m)$ for $m \geq f(k)$ as required in the statement of the lemma. Similarly for $l < \omega$ find the unique m such that $s(m) \leq l < s(m+1)$ and write $m = L_n^\rho + j$. Pick $d^*(l) \in [d_{j_n}(\mathfrak{z}_j^n)]^{l+1}$ in such a way that for $s(m) \leq l < l+1 < s(m+1)$ we have $\max(d^*(l)) < \min(d^*(l+1))$. This is possible by already proved $|d_{j_n}(\mathfrak{z}_j^n)| \geq t(m)$.

Claim 3.6. *For every $k < \omega$ we have*

$$\max(e^*(k)) < \min(e^*(k+1)) \text{ and } \max(d^*(k)) < \min(d^*(k+1)).$$

Proof. For a fixed $n < \omega$, if $j < j+1 < R_n$, then $\zeta_j^n < \zeta_{j+1}^n$ and $\mathfrak{z}_j^n < \mathfrak{z}_{j+1}^n$ and so $\max(e(\zeta_j^n)) < \min(e(\zeta_{j+1}^n))$ and $\max(d_{j_n}(\mathfrak{z}_j^n)) < \min(d_{j_n}(\mathfrak{z}_{j+1}^n))$. Moreover, for a fixed $n < \omega$ and $j < R_n$, if $s(L_n^\rho + j) \leq l < l+1 < s(L_n^\rho + j+1)$, then $\max(d^*(l)) < \min(d^*(l+1))$ by definition. Therefore it suffices to show that for $n < \omega$, $j < R_n$, and $j' < R_{n+1}$, $\max(e(\zeta_j^n)) < \min(e(\zeta_{j'}^{n+1}))$ and $\max(d_{j_n}(\mathfrak{z}_j^n)) < \min(d_{j_{n+1}}(\mathfrak{z}_{j'}^{n+1}))$.

To see the first inequality, we argue by contradiction. Suppose $y \in e(\zeta_j^n)$, $y' \in e(\zeta_{j'}^{n+1})$, and $y' \leq y$. As noted above, $\zeta_{j'}^{n+1} \geq M_{n+1}^e$. Thus $y, y' \in \text{set}(e) \llbracket M_{n+1}^e \rrbracket$, and so by (1) and (3) of Claim 3.5,

$$\begin{aligned} \pi_{\delta_{n+1}, \delta_{m(n,j)}}(z_{j'}^{n+1}) &= \pi_{\delta_{n+1}, \delta_{m(n,j)}}(\pi_{\delta, \delta_{n+1}}(y')) = \pi_{\delta, \delta_{m(n,j)}}(y') \leq \pi_{\delta, \delta_{m(n,j)}}(y) = \\ \pi_{\delta_n, \delta_{m(n,j)}}(\pi_{\delta, \delta_n}(y)) &= \pi_{\delta_n, \delta_{m(n,j)}}(z_j^n) = x_j^n \leq D_{m(n,j)}(K_{m(n,j), n+1} - 1). \end{aligned}$$

However $z_{j'}^{n+1} \in C_{n+1} \setminus F_{n+1}(g'(n+1))$. But then by (6) of Lemma 3.2 applied to $n+1$ and $m = m(n, j)$, $\pi_{\delta_{n+1}, \delta_{m(n,j)}}(z_{j'}^{n+1}) > D_{m(n,j)}(K_{m(n,j), n+1} - 1)$. This is a contradiction which proves the first inequality.

The second inequality is also proved by contradiction. So suppose $y \in d_{j_n}(\mathfrak{z}_j^n)$, $y' \in d_{j_{n+1}}(\mathfrak{z}_{j'}^{n+1})$, and $y' \leq y$. As noted above, $\mathfrak{z}_j^n \geq M_n^d$ and $\mathfrak{z}_{j'}^{n+1} \geq M_{n+1}^d$. Moreover $d_{j_{n+1}} \leq_{Q^d(n+1)} d_{j_n}$, $M_{n+1}^d \geq Q^d(n+1)$, and $M_{n+1}^d \geq M_n^d$. So there exists $l \geq M_n^d$ with $d_{j_{n+1}}(\mathfrak{z}_{j'}^{n+1}) \subset d_{j_n}(l)$. Thus $y' \in \text{set}(d_{j_{n+1}}) \llbracket M_{n+1}^d \rrbracket$ and $y, y' \in \text{set}(d_{j_n}) \llbracket M_n^d \rrbracket$. Therefore by (2) and (4) of Claim 3.5,

$$\begin{aligned} \pi_{\delta_{n+1}, \delta_{m(n,j)}}(z_{j'}^{n+1}) &= \pi_{\delta_{n+1}, \delta_{m(n,j)}}(\pi_{\delta_{n+1}}(y')) = \pi_{\delta_{m(n,j)}}(y') \leq \pi_{\delta_{m(n,j)}}(y) = \\ \pi_{\delta_n, \delta_{m(n,j)}}(\pi_{\delta_n}(y)) &= \pi_{\delta_n, \delta_{m(n,j)}}(z_j^n) = x_j^n \leq D_{m(n,j)}(K_{m(n,j), n+1} - 1). \end{aligned}$$

However $\pi_{\delta_{n+1}, \delta_{m(n,j)}}(z_{j'}^{n+1}) > D_{m(n,j)}(K_{m(n,j), n+1} - 1)$ as pointed out in the previous paragraph. This is a contradiction which completes the proof. \square

So for now we have settled that $e^* \leq_0 e$ and that for every $n < \omega$ there is $m \geq f(n)$ such that $e^*(n) \subset e(m)$. Define $\pi : \omega \rightarrow \omega$ as follows: for every $k \in \omega \setminus \text{set}(d^*)$ let $\pi(k) = 0$, while for every $k \in \text{set}(d^*)$ let m be unique such that $k \in d^*(m)$ and define $\pi(k) = \text{set}(e^*)(m)$.

Claim 3.7. *The sequences $\langle e^*(n) : n < \omega \rangle$ and $\langle d^*(n) : n < \omega \rangle$ belong to \mathbb{P} and satisfy the following conditions:*

- (1) $\forall j < \omega [d^* \leq d_j]$ and $\text{set}(e^*) = \pi'' \text{set}(d^*)$;
- (2) $\forall \alpha \in X \forall^\infty k \in \text{set}(d^*) [\pi_\alpha(k) = \pi_{\delta, \alpha}(\pi(k))]$;
- (3) $\forall \alpha \in X [\pi''_{\delta, \alpha} \text{set}(e^*) \in \mathcal{U}_\alpha]$;
- (4) *There is ψ such that $\langle \pi, \psi, d^* \rangle$ is a normal triple.*

Proof. First note that e^* and d^* belong to \mathbb{P} . Next, we prove (1). It suffices to prove that $d^* \leq d_{j_n}$, for all $n < \omega$. Fix $n < \omega$. Take any $l \geq s(L_n^\rho)$. Let m be such that $s(m) \leq l < s(m+1)$ and n' and $j' < R_{n'}$ such that $m = L_{n'}^\rho + j'$. Note $m \geq L_n^\rho$ and $n' \geq n$. Then $d^*(l) \subset d_{j_{n'}}(\mathfrak{z}_{j'}^{n'})$ and $\mathfrak{z}_{j'}^{n'} \geq M_{n'}^d$. As noted earlier $\mathfrak{z}_{j'}^{n'} \geq g(L_{n'}^\rho + j') = g(m) \geq s(m+1) \geq l$. So since $d_{j_{n'}} \leq_{M_{n'}^d} d_{j_n}$, there is $l' \geq \mathfrak{z}_{j'}^{n'} \geq l$ so that $d^*(l) \subset d_{j_{n'}}(\mathfrak{z}_{j'}^{n'}) \subset d_{j_n}(l')$, showing $d^* \leq_{s(L_n^\rho)} d_{j_n}$. To see that $\text{set}(e^*) = \pi'' \text{set}(d^*)$ note that by the definition of π we have that for every $n < \omega$ holds $\pi'' d^*(n) = \{\text{set}(e^*)(n)\}$. So (1) is proved.

Now we prove (2). First we prove $\forall^\infty k \in \text{set}(d^*) [\pi_{\delta_n}(k) = \pi_{\delta, \delta_n}(\pi(k))]$, for every $n < \omega$. Fix $n < \omega$ and consider any $n' \geq n$ and $j' < R_{n'}$. It suffices to show that for any $s(L_{n'}^\rho + j') \leq l < s(L_{n'}^\rho + j' + 1)$ and $k \in d^*(l)$, $\pi_{\delta_n}(k) = \pi_{\delta, \delta_n}(\pi(k))$. By definition $d^*(l) \subset d_{j_{n'}}(\mathfrak{z}_{j'}^{n'})$ and $\pi(k) \in e^*(L_{n'}^\rho + j') \subset e(\zeta_{j'}^{n'})$. Therefore $\pi_{\delta_{n'}}(k) = \zeta_{j'}^{n'} = \pi_{\delta, \delta_{n'}}(\pi(k))$. Also $k \in \text{set}(d_{j_{n'}}) \llbracket M_{n'}^d \rrbracket$ and $\pi(k) \in \text{set}(e) \llbracket M_{n'}^e \rrbracket$ because $\zeta_{j'}^{n'} \geq M_{n'}^e$ and $\mathfrak{z}_{j'}^{n'} \geq M_{n'}^d$. Thus by (1) and (2) of Claim 3.5, $\pi_{\delta_n}(k) = \pi_{\delta_{n'}, \delta_n}(\pi_{\delta_{n'}}(k)) = \pi_{\delta_{n'}, \delta_n}(\pi_{\delta, \delta_{n'}}(\pi(k))) = \pi_{\delta, \delta_n}(\pi(k))$, as needed. For the more general claim fix $\alpha \in X$ and find $n < \omega$ so that $\alpha \leq \delta_n$. By (3b) of Lemma 3.4, there exist $i < \omega$ and L_0 so that $\forall k \in \text{set}(d_i) \llbracket L_0 \rrbracket [\pi_\alpha(k) = \pi_{\delta_n, \alpha}(\pi_{\delta_n}(k))]$. Let L_1 be minimal with $d^* \leq_{L_1} d_i$. By (2b) of Lemma 3.4 and by the fact that $\text{set}(e^*) \subset \text{set}(e)$, there is L_2 so that $\forall k \in \text{set}(e^*) \llbracket L_2 \rrbracket [\pi_{\delta, \alpha}(k) = \pi_{\delta_n, \alpha}(\pi_{\delta, \delta_n}(k))]$. Let L_3 be so that $\forall k \in \text{set}(d^*) \llbracket L_3 \rrbracket [\pi_{\delta_n}(k) = \pi_{\delta, \delta_n}(\pi(k))]$. Let $L = \max\{L_0, L_1, L_2, L_3\}$. If $k \in \text{set}(d^*) \llbracket L \rrbracket$, then $k \in \text{set}(d_i) \llbracket L_0 \rrbracket$ and $\pi(k) \in \text{set}(e^*) \llbracket L_2 \rrbracket$. So $\pi_\alpha(k) = \pi_{\delta_n, \alpha}(\pi_{\delta_n}(k)) = \pi_{\delta_n, \alpha}(\pi_{\delta, \delta_n}(\pi(k))) = \pi_{\delta, \alpha}(\pi(k))$. Thus $\forall k \in \text{set}(d^*) \llbracket L \rrbracket [\pi_\alpha(k) = \pi_{\delta, \alpha}(\pi(k))]$, proving (2).

Now we come to (3). We first show that for each $m < \omega$, $D_m \subset^* \pi''_{\delta, \delta_m} \text{set}(e^*)$. Fix $m < \omega$. As $\langle K_{m, n} : m \leq n < \omega \rangle$ is strictly increasing with n , it suffices to show that for each $n \geq m$, $D_m[K_{m, n}, K_{m, n+1}] \subset \pi''_{\delta, \delta_m} \text{set}(e^*)$. Let $n \geq m$ and $u \in D_m[K_{m, n}, K_{m, n+1}]$ be given. Put $m' =$

$$\max\{m'' \leq n : m \leq m'' \text{ and } \exists u'' \in D_{m''}[K_{m'', n}, K_{m'', n+1}] [\pi_{\delta_{m''}, \delta_m}(u'') = u]\},$$

and choose $u' \in D_{m'}[K_{m', n}, K_{m', n+1}]$ with $\pi_{\delta_{m'}, \delta_m}(u') = u$. We claim that $u' \in H_{m', n}^\rho$. Suppose not. Then there exist $m' < m'' \leq n$ and $u'' \in D_{m''}[K_{m'', n}, K_{m'', n+1}]$ with $\pi_{\delta_{m''}, \delta_{m'}}(u'') = u'$. Now $u'' \in D_{m''}[K_{m'', m''}]$ because $K_{m'', m''} \leq K_{m'', n}$. So by (5) of Lemma 3.2 applied to m'' , $u'' \in F_{m''} \subset C_{m''}$. By one of the properties of $C_{m''}$ listed in Lemma 3.2, $\pi_{\delta_{m''}, \delta_m}(u'') = \pi_{\delta_{m''}, \delta_m}(\pi_{\delta_{m''}, \delta_{m'}}(u'')) = \pi_{\delta_{m''}, \delta_m}(u') = u$. However this contradicts the choice of m' . Thus $u' \in H_{m', n}^\rho \subset \Delta_n^\rho$. So let $j < R_n$ be so that $u' = x_j^n$. Note that $m(n, j) = m'$. Also $z_j^n \in F_n \subset C_n$, and so $\pi_{\delta_n, \delta_m}(z_j^n) = \pi_{\delta_{m'}, \delta_m}(\pi_{\delta_n, \delta_{m'}}(z_j^n)) = \pi_{\delta_{m'}, \delta_m}(u') = u$. Now if $k \in e^*(L_n^\rho + j) \subset e(\zeta_j^n)$, then $k \in \text{set}(e) \llbracket M_n^e \rrbracket$ because $\zeta_j^n \geq M_n^e$. So by (1) of Claim 3.5, $\pi_{\delta, \delta_m}(k) = \pi_{\delta_n, \delta_m}(\pi_{\delta_n}(k)) = \pi_{\delta_n, \delta_m}(z_j^n) = u$, showing that $u \in \pi''_{\delta, \delta_m} \text{set}(e^*)$. This concludes the proof that $D_m \subset^* \pi''_{\delta, \delta_m} \text{set}(e^*)$. As $D_m \in \mathcal{U}_{\delta_m}$, this shows that $\pi''_{\delta, \delta_m} \text{set}(e^*) \in \mathcal{U}_{\delta_m}$, for all $m \in \omega$. Now for the more general statement, fix $\alpha \in X$. Find $m \in \omega$ with $\delta_m \geq \alpha$. By (2b) of Lemma 3.4, there is L so that $\forall k \in \text{set}(e^*) \llbracket L \rrbracket [\pi_{\delta, \alpha}(k) = \pi_{\delta_m, \alpha}(\pi_{\delta, \delta_m}(k))]$. Put $A = \text{set}(e^*) \llbracket L \rrbracket$. Since $\pi''_{\delta, \delta_m} A \in \mathcal{U}_{\delta_m}$, $\pi''_{\delta_m, \alpha} \pi''_{\delta, \delta_m} A \in \mathcal{U}_\alpha$. Hence $\pi''_{\delta_m, \alpha} \pi''_{\delta, \delta_m} A \subset \pi''_{\delta, \alpha} A \subset \pi''_{\delta, \alpha} \text{set}(e^*)$, implying $\pi''_{\delta, \alpha} \text{set}(e^*) \in \mathcal{U}_\alpha$, which proves (3).

For the proof of (4), consider function $\psi : \omega \rightarrow \omega$ defined in the following way: for $k < \omega$ let $\psi(k) = \text{set}(e^*)(k)$. It is clear that $\langle \pi, \psi, d^* \rangle$ is a normal triple. \square

The last claim proves the lemma. \square

When CH is replaced by MA, the statement of Lemma 3.4 needs to be generalized as follows. δ is allowed to be any ordinal with $\text{cof}(\delta) < \mathfrak{c}$, and the decreasing sequence $\langle d_j : j < \omega \rangle$ is replaced by the decreasing sequence $\langle d_j : j < \text{cof}(\delta) \rangle$. This version can be proved under MA by taking a suitably generic filter over a poset consisting of finite approximations to d^* , e^* , and π together with some finite side conditions. The exact definition of this poset can be formulated by examining the proofs of Lemmas 3.2 and 3.4.

Lemma 3.8 will be used in the proof that the poset \mathbb{Q}^δ is countably closed. The requirement in Lemma 3.2 that $F_n = C_n \cap \pi''_{n+1,n} C_{n+1}$ will be used crucially in this proof.

Lemma 3.8. *Let $\langle \mathcal{U}_n : n < \omega \rangle$ be a sequence of distinct rapid P-points. Assume that $\langle \pi_{n,m} : m \leq n < \omega \rangle \subset \omega^\omega$ is a sequence so that $\pi_{n,n} = \text{id}$ ($n < \omega$) and:*

- (1) $\forall m \leq n < \omega \forall a \in \mathcal{U}_n [\pi''_{n,m} a \in \mathcal{U}_m]$;
- (2) $\forall m \leq n \leq k < \omega \exists a \in \mathcal{U}_k \forall l \in a [\pi_{k,m}(l) = \pi_{n,m}(\pi_{k,n}(l))]$;
- (3) $\forall m \leq n < \omega \exists a \in \mathcal{U}_n \forall x, y \in a [x \leq y \Rightarrow \pi_{n,m}(x) \leq \pi_{n,m}(y)]$.

Then for every $e \in \mathbb{P}$ there is a sequence of maps in ω^ω , $\langle \pi_n : n < \omega \rangle$, satisfying:

- (4) $\forall n < \omega [\pi''_n \text{set}(e) \in \mathcal{U}_n]$;
- (5) $\forall m \leq n < \omega \forall^\infty k \in \text{set}(e) [\pi_m(k) = \pi_{n,m}(\pi_n(k))]$;
- (6) *for every $n < \omega$ there are $\psi_n \in \omega^\omega$ and $b_n \in \mathbb{P}$ such that $e \leq b_n$ and that $\langle \pi_n, \psi_n, b_n \rangle$ is a normal triple.*

Proof. Define $E_k = \omega$, for every $k < \omega$. Let M be a countable elementary submodel of $H_{(2^{\mathfrak{c}})^+}$ containing $\bar{\pi} = \langle \pi_{n,m} : m \leq n < \omega \rangle$, $\langle \mathcal{U}_n : n < \omega \rangle$. For $m < \omega$, let $D_m \in \mathcal{U}_m$ be such that $D_m \subset^* A$ for every $A \in \mathcal{U}_m \cap M$. Now Lemma 3.2 applies to M , function $f = \text{id}$, sequences $\bar{\pi}$, $\bar{D} = \langle D_m : m < \omega \rangle$, $\langle \mathcal{U}_n : n < \omega \rangle$ and $\langle E_n : n < \omega \rangle$. Let sequences $\langle F_n : n < \omega \rangle$, $\langle C_n : n < \omega \rangle$, $\langle g'(n) : n < \omega \rangle$ and $\bar{K} = \langle K_{m,n} : m \leq n < \omega \rangle$ be as in Lemma 3.2. Denote $\rho = \langle \bar{D}, \bar{K}, \bar{\pi} \rangle$, R_n , z_j^n , x_j^n and $m(n, j)$ ($j < R_n$) as in the conclusion of Lemma 3.2.

For each $n < \omega$, define $I_n = \{L_n^\rho + j : j < R_n\}$. Recall from the proof of Lemma 3.4 that $\langle I_n : n < \omega \rangle$ is an interval partition of ω . Fix $m < \omega$. For $n < m$ and $j < R_n$, define $\psi_m(L_n^\rho + j) = 0$, while for $m \leq n$ and $j < R_n$, define $\psi_m(L_n^\rho + j) = \pi_{n,m}(z_j^n)$. Thus $\psi_m \in \omega^\omega$ and we claim that it is increasing. It suffices to consider the following two cases. Case 1 is when $n < \omega$, $j \leq j' < R_n$ and we wish to compare $\psi_m(L_n^\rho + j)$ and $\psi_m(L_n^\rho + j')$. If $n < m$, then both these values are 0. If $m \leq n$, then $\psi_m(L_n^\rho + j) = \pi_{n,m}(z_j^n) \leq \pi_{n,m}(z_{j'}^n) = \psi_m(L_n^\rho + j')$ because $z_j^n \leq z_{j'}^n$ and because $z_j^n, z_{j'}^n \in C_n$. Now we come to case 2, which is when we wish to compare $\psi_m(L_n^\rho + j)$ and $\psi_m(L_{n+1}^\rho + j')$, for some $n < \omega$, $j < R_n$, and $j' < R_{n+1}$. First, if $n < m$, then $\psi_m(L_n^\rho + j) = 0 \leq \psi_m(L_{n+1}^\rho + j')$. So assume that $m \leq n$. Since $z_j^n \in F_n$, there exists $z \in C_{n+1}$ with $\pi_{n+1,n}(z) = z_j^n$. By a property of C_{n+1} from Lemma 3.2, $\pi_{n+1,m(n,j)}(z) = \pi_{n,m(n,j)}(\pi_{n+1,n}(z)) = \pi_{n,m(n,j)}(z_j^n) = x_j^n \leq D_{m(n,j)}(K_{m(n,j),n+1} - 1)$. It follows from (6) of Lemma 3.2 applied to $n+1$ that $z < F_{n+1}(g'(n+1)) \leq z_{j'}^{n+1}$. Since $z_{j'}^{n+1} \in C_{n+1}$, $\pi_{n,m}(z_j^n) = \pi_{n,m}(\pi_{n+1,n}(z)) = \pi_{n+1,m}(z) \leq \pi_{n+1,m}(z_{j'}^{n+1})$. So $\psi_m(L_n^\rho + j) = \pi_{n,m}(z_j^n) \leq \pi_{n+1,m}(z_{j'}^{n+1}) = \psi_m(L_{n+1}^\rho + j')$. Thus we have proved that ψ_m is increasing.

Now for each $m < \omega$, define $\pi_m \in \omega^\omega$ as follows. Let $k \in \omega$. If $k \notin \text{set}(e)$, then set $\pi_m(k) = 0$; else let $l \in \omega$ be unique such that $k \in e(l)$, and set $\pi_m(k) = \psi_m(l)$. We check that (4)–(6) are satisfied. We begin with (5). Fix $m \leq l < \omega$. Consider any $k \in \text{set}(e) \llbracket L_l^\rho \rrbracket$. Then $k \in e(L_n^\rho + j)$, for some $l \leq n < \omega$ and $j < R_n$. So $\pi_m(k) = \psi_m(L_n^\rho + j) = \pi_{n,m}(z_j^n)$ and $\pi_l(k) = \psi_l(L_n^\rho + j) = \pi_{n,l}(z_j^n)$. Since $z_j^n \in$

C_n , $\pi_{n,m}(z_j^n) = \pi_{l,m}(\pi_{n,l}(z_j^n))$. Therefore, $\pi_m(k) = \pi_{n,m}(z_j^n) = \pi_{l,m}(\pi_{n,l}(z_j^n)) = \pi_{l,m}(\pi_l(k))$, as needed for (5).

Next we prove (4). Fix $m < \omega$. We will show $D_m \subset^* \pi_m'' \text{set}(e)$. As the sequence $\langle K_{m,n} : m \leq n < \omega \rangle$ is strictly increasing with n , it suffices to show that for each $n \geq m$, $D_m[K_{m,n}, K_{m,n+1}] \subset \pi_m'' \text{set}(e)$. Let $n \geq m$ and $u \in D_m[K_{m,n}, K_{m,n+1}]$ be given. Apply the same argument as in the proof of (3) of Claim 3.7 to find m' and u' so that $m \leq m' \leq n$, $u' \in D_{m'}[K_{m',n}, K_{m',n+1}]$, $\pi_{m',m}(u') = u$, and $u' \in H_{m',n}^\rho \subset \Delta_n^\rho$. Let $j < R_n$ be such that $x_j^n = u'$. Note that $m(n, j) = m'$. Also $z_j^n \in F_n \subset C_n$. So by a property of C_n from Lemma 3.2, $\pi_{n,m}(z_j^n) = \pi_{m',m}(\pi_{n,m'}(z_j^n)) = \pi_{m',m}(u') = u$. Now if $k \in e(L_n^\rho + j)$, then since $m \leq n$, by definition, $\pi_m(k) = \pi_{n,m}(z_j^n) = u$. Thus $u \in \pi_m'' \text{set}(e)$. This proves $D_m \subset^* \pi_m'' \text{set}(e)$, which proves (4) because $D_m \in \mathcal{U}_m$.

We still have to prove (6). Fix $m < \omega$. We have already defined ψ_m and proved that it is increasing. Let $b_m = e$. By definition of π_m , $\pi_m'' b_m(l) = \{\psi_m(l)\}$, for each $l < \omega$, and $\pi_m(k) = 0$, for all $k \in \omega \setminus \text{set}(b_m)$. Also $\text{ran}(\psi_m)$ is infinite because $\pi_m'' \text{set}(e) \in \mathcal{U}_m$ and $\pi_m'' \text{set}(e) \subset \text{ran}(\psi_m)$. Therefore $\langle \pi_m, \psi_m, b_m \rangle$ is a normal triple and $e \leq b_m$, as needed. \square

In the context of MA, the statement of Lemma 3.8 will be modified as follows. The sequence $\langle \mathcal{U}_n : n < \omega \rangle$ will be replaced with the sequence $\langle \mathcal{U}_\alpha : \alpha < \lambda \rangle$, where λ is a cardinal $< \mathfrak{c}$. Moreover each \mathcal{U}_α will be assumed to be a rapid P_c -point. And, of course, there will be a map $\pi_{\beta,\alpha}$, for each $\alpha \leq \beta < \lambda$. The sequence $\langle \pi_n : n < \omega \rangle$ in the conclusion of Lemma 3.8 will be replaced by the sequence $\langle \pi_\alpha : \alpha < \lambda \rangle$. This version can be proved under MA by taking a suitably generic filter over a poset consisting of finite approximations to the sequence $\langle \pi_\alpha : \alpha < \lambda \rangle$ together with some finite side conditions. The exact definition of the forcing notion that is required is somewhat cumbersome, and so we do not include it here. Its precise definition as well as the proof that it has all of the necessary properties will appear in a future paper.

The next lemma will also be used in the proof that the poset \mathbb{Q}^δ is countably closed. It is like a simple special case of Lemma 3.4 in spirit, but does not directly follow from the statement of Lemma 3.4.

Lemma 3.9. *Let \mathcal{U} be a rapid P-point, π a mapping in ω^ω and $\langle d_m : m < \omega \rangle$ a decreasing sequence of conditions in \mathbb{P} such that $\pi'' \text{set}(d_n) \in \mathcal{U}$ for every $n < \omega$. Suppose that there are $b \in \mathbb{P}$ and $\psi \in \omega^\omega$ so that $\langle \pi, \psi, b \rangle$ is a normal triple and $d_0 \leq b$. Then there is $d \in \mathbb{P}$ such that $\pi'' \text{set}(d) \in \mathcal{U}$ and $d \leq d_n$ for every $n < \omega$.*

Proof. First we define sequence of numbers n_k ($k < \omega$) as follows: n_0 is minimal such that $d_0 \leq_{n_0} b$, while $n_{k+1} = \max\{l, n_k\}$ for l minimal such that $d_{k+1} \leq_l d_k$. By Remark 2.6(2) we have $d_k \leq_{n_k} b$, $d_k \leq_{n_k} d_l$ for $l \leq k$ and consequently $\text{set}(d_{k+1}) \llbracket n_{k+1} \rrbracket \subset \text{set}(d_k) \llbracket n_k \rrbracket$ for $k < \omega$. Let $C_k = \pi'' \text{set}(d_k) \llbracket n_k \rrbracket$ for $k < \omega$ and notice that $C_{k+1} \subset C_k$ and $C_k \in \mathcal{U}$ for $k < \omega$. So since \mathcal{U} is a rapid ultrafilter, by Lemma 2.3, for every $k < \omega$ there is $D_k \in \mathcal{U}$ such that for every $n < \omega$ there is $m \geq 2(n+1)$ such that $D_k(n) = C_k(m)$. Because \mathcal{U} is a P-point there is $D \in \mathcal{U}$ such that $D \subset^* D_k$ for every $k < \omega$ and $D \subset D_0$. For every $k, l < \omega$ define set $F_l^k = \{m < \omega : \pi'' d_k(m) = \{D(l)\}\}$. By Lemma 2.9, if $D(l) \in C_k$, then $\pi_l^k = \max(F_l^k)$ and $\sigma_l^k = \min(F_l^k \setminus n_k)$ are well defined. For a fixed k and $l_1 < l_2$ such that $D(l_1), D(l_2) \in C_k$, again by Lemma 2.9, we have $\pi_{l_1}^k < \sigma_{l_2}^k \leq \pi_{l_2}^k$. Also, if $l_1 < l_2 < \omega$, $k_1 < k_2 < \omega$, $D(l_1) \in D_{k_1}$ and $D(l_2) \in D_{k_2}$, then it is easy to see that $\max(d_{k_1}(\pi_{l_1}^{k_1})) < \min(d_{k_2}(\pi_{l_2}^{k_2}))$. Now, by induction on k , we construct numbers $g(k)$ and sets $d(m)$ for $m < g(k)$ so that for $k < \omega$:

- (1) $\pi_{g(k)}^k \geq g(k)$;
- (2) $D[g(k)] \subset D_k$;
- (3) if $k > 0$ then $\forall l \in [g(k-1), g(k)]$ $[d(l) \in [d_{k-1}(\pi_l^{k-1})]^{l+1}]$.
- (4) if $k > 0$ then $g(k) > g(k-1)$ and $\forall l < g(k) - 1$ $[\max(d(l)) < \min(d(l+1))]$.

Let $g(0) = 0$ and note that (1)–(4) are satisfied. So fix $k \in \omega$ and assume that for every $m \leq k$ numbers $g(m)$ are defined, and that for every $l < g(k)$ sets $d(l)$ are defined. Let X_k be the minimal number such that $X_k > g(k)$ and $D[X_k] \subset D_{k+1}$ and define $g(k+1) = 2X_k$. First note that since $X_k > g(k)$ we have that $g(k+1) = 2X_k > g(k)$. Since by inductive hypothesis $D[g(k)] \subset D_k$ and $\mathfrak{d}_{g(k)}^k \geq g(k)$, Lemma 2.9 implies that for $g(k) \leq l < l' < g(k+1)$ we have $\mathfrak{d}_l^k < \mathfrak{d}_{l'}^k$. So we can pick $d(l) \in [d_k(\mathfrak{d}_l^k)]^{l+1}$ for $g(k) \leq l < g(k+1)$. Now we prove that (1)–(4) hold. To prove (1) note that by the choice of X_k we know that $D[X_k, g(k+1)] \subset D_{k+1}$. So $D(g(k+1)) = D_{k+1}(m)$ for some $m \geq g(k+1) - X_k = X_k$. This implies that $D(g(k+1)) = C_{k+1}(m')$ for some $m' \geq 2m \geq 2X_k \geq g(k+1)$. Now by Lemma 2.9 applied to C_{k+1} , d_{k+1} and $\langle \pi, \psi, b \rangle$ we have $\mathfrak{d}_{g(k+1)}^{k+1} \geq g(k+1)$. Condition (2) follows from the fact that $g(k+1) = 2X_k \geq X_k$ and $D[X_k] \subset D_{k+1}$. Condition (3) holds by construction. To see that (4) is true we distinguish three cases: either $l > g(k) - 1$ or $l < g(k) - 1$ or $l = g(k) - 1$. If $l < g(k) - 1$ then it follows from the inductive hypothesis and the fact that $g(0) = 0$. If $l = g(k) - 1$ then because $g(k-1) < g(k)$ we have $g(k) - 1 \geq g(k-1)$ so $k > 0$. By (2) applied to $k-1$ and k we know $D(g(k)-1) \in D_{k-1}$ and $D(g(k)) \in D_k$ so the statement follows from the observation in the first paragraph that $\max(d_{k-1}(\mathfrak{d}_{g(k)-1}^{k-1})) < \min(d_k(\mathfrak{d}_{g(k)}^k))$. If $l > g(k) - 1$ then it follows from the facts that $\mathfrak{d}_l^k < \mathfrak{d}_{l+1}^k$. \square

4. ADDING AN ULTRAFILTER ON TOP

In this section, for a given $\delta < \omega_2$, we introduce the poset for adding a rapid P-point \mathcal{U}_δ together with a sequence of maps $\langle \pi_{\delta, \alpha} : \alpha \leq \delta \rangle$ on top of an already constructed δ -generic sequence of P-points $\langle \mathcal{U}_\alpha : \alpha < \delta \rangle$ and Rudin-Keisler maps $\langle \pi_{\beta, \alpha} : \alpha \leq \beta < \delta \rangle$. So fix a $\delta < \omega_2$ and a δ -generic sequence $S = \langle \langle c_i^\alpha : \alpha < \delta \wedge i < \mathfrak{c} \rangle, \langle \pi_{\beta, \alpha} : \alpha \leq \beta < \delta \rangle \rangle$ for the rest of this section.

We briefly explain the idea behind the definition of \mathbb{Q}^δ given below. We would like a generic filter for \mathbb{Q}^δ to produce two sequences $\bar{C} = \langle c_i^\delta : i < \mathfrak{c} \rangle$ and $\bar{\pi} = \langle \pi_{\delta, \alpha} : \alpha \leq \delta \rangle$ which, when added to S , will result in a $\delta + 1$ -generic sequence. Conditions in \mathbb{Q}^δ are essentially countable approximations to such objects. The first coordinate of the condition q will be an element of \bar{C} , and the fourth coordinate fixes $\bar{\pi}$ on a countable subset of δ . Clauses (4a), (4b), and (4c) below say that the maps that have already been determined by q work in accordance with clauses (5a), (5b), and (5c) of Definition 2.10. Clause (3) below says that X_q , which is the countable set on which $\bar{\pi}$ has been fixed, always has a maximal element unless X_q is cofinal in δ . This assumption will simplify some arguments.

Definition 4.1. Let \mathbb{Q}^δ be the set of all $q = \langle c_q, \gamma_q, X_q, \langle \pi_{q, \alpha} : \alpha \in X_q \rangle \rangle$ such that:

- (1) $c_q \in \mathbb{P}$;
- (2) $\gamma_q \leq \delta$;
- (3) $X_q \in [\delta]^{\leq \aleph_0}$ is such that $\gamma_q = \sup(X_q)$ and $\gamma_q \in X_q$ iff $\gamma_q < \delta$;
- (4) $\pi_{q, \alpha}$ ($\alpha \in X_q$) are mappings in ω^ω such that:
 - (a) $\pi_{q, \alpha}'' \text{set}(c_q) \in \mathcal{U}_\alpha$;
 - (b) $\forall \alpha, \beta \in X_q [\alpha \leq \beta \Rightarrow \forall^\infty k \in \text{set}(c_q) [\pi_{q, \alpha}(k) = \pi_{q, \beta}(\pi_{q, \beta}(k))]]$;
 - (c) there is $\psi_{q, \alpha} \in \omega^\omega$ and $b_{q, \alpha} \geq c_q$ such that $\langle \pi_{q, \alpha}, \psi_{q, \alpha}, b_{q, \alpha} \rangle$ is a normal triple;

Let the ordering on \mathbb{Q}^δ be given by: $q_1 \leq q_0$ if and only if

$$c_{q_1} \leq c_{q_0} \text{ and } X_{q_1} \supset X_{q_0} \text{ and for every } \alpha \in X_{q_0}, \pi_{q_1, \alpha} = \pi_{q_0, \alpha}.$$

In the situation where CH is replaced by MA, \mathbb{Q}^δ would consist of approximations of size $< \mathfrak{c}$ instead of countable ones. Thus X_q would be a set of size less than \mathfrak{c} .

Remark 4.2. It is easy to check that $\langle \mathbb{Q}^\delta, \leq \rangle$ defined in this way is a partial order. Note also that $\mathbb{Q}^\delta \neq 0$. Namely, if $\delta = 0$, then we can take $q = \langle c, 0, 0, 0 \rangle$ for any $c \in \mathbb{P}$. If

$\delta \neq 0$, then let $q = \langle c_q, \gamma_q, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle$ be such that: c_q is arbitrary in \mathbb{P} ; $\gamma_q = 0$; $X_q = \{0\}$; $\pi_{q,0} \in \omega^\omega$ is given by: for $k \in \text{set}(c_q)$ let $\pi_{q,0}(k) = n$ for $k \in c_q(n)$, while $\pi_{q,0}(k) = 0$ otherwise. First note that conditions (1)–(3) of Definition 4.1 are satisfied. Because $\pi''_{q,0} \text{set}(c_q) = \omega$ we know that (4a) holds. It is also easy to see that $\langle \pi_{q,0}, \text{id}, c_q \rangle$ is a normal triple by definition of $\pi_{q,0}$ so condition (4c) is true. To see that condition (4b) is also true note that $\pi_{0,0} = \text{id}$ by Definition 2.10(5). So $q \in \mathbb{Q}^\delta$.

Remark 4.3. Let $q = \langle c_q, \gamma_q, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle \in \mathbb{Q}^\delta$. Let $c_{q'} \in \mathbb{P}$ be such that $c_{q'} \leq c_q$. Then $q' = \langle c_{q'}, \gamma_q, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle$ satisfies conditions (1), (2), (3), (4b) and (4c) of Definition 4.1. Moreover, if q' also satisfies Definition 4.1(4a), then $q' \in \mathbb{Q}^\delta$ and $q' \leq q$.

Instead of forcing with the poset \mathbb{Q}^δ , we would like to build a sufficiently generic filter over it in the ground model itself. \mathbb{Q}^δ needs to be countably closed for this to be feasible. We prove this fact next. The next lemma is the crux of the whole construction. We briefly sketch the idea of its proof. So suppose that $\langle q_n : n \in \omega \rangle$ is a decreasing sequence of conditions in \mathbb{Q}^δ . We want to find a lower bound. There are four natural cases to consider. We start with the simpler ones. The most trivial case is when $\delta = 0$. Then we just have a decreasing sequence in \mathbb{P} and bounding them is easy. Next, it could be the case that for all $n \in \omega$, $\gamma_{q_n} = \gamma$, for some fixed $\gamma < \delta$. Then we essentially have a fixed ultrafilter \mathcal{U}_γ , a descending sequence in \mathbb{P} , and a fixed map taking each element of this sequence into \mathcal{U}_γ . We wish to find a bound for this sequence in \mathbb{P} whose image is still in \mathcal{U}_γ . Lemma 3.9 is set up precisely to handle this situation, so we apply it. The third case is when the γ_{q_n} form an increasing sequence converging to δ . Then we have a decreasing sequence in \mathbb{P} , some countable cofinal $Y \subset \delta$, and a sequence of maps taking members of the decreasing sequence in \mathbb{P} to various ultrafilters indexed by Y . We would like to find a lower bound for this decreasing sequence in \mathbb{P} whose images under each of the given maps are in the corresponding ultrafilters. This is almost like the situation in Lemma 3.4, except that e and its associated maps are missing. So we first apply Lemma 3.8 to find these things, and then apply Lemma 3.4 to them. The final and trickiest case is when the γ_{q_n} form an increasing sequence converging to some $\mu < \delta$. Then the ultrafilter \mathcal{U}_μ must have been constructed to anticipate this situation. This is where clause (6) of Definition 2.10 enters. We have a decreasing sequence in \mathbb{P} , a countable cofinal $Y \subset \mu$, and a sequence of maps as before. We would like to find a lower bound for this decreasing sequence in \mathbb{P} as well as a new map associated with \mathcal{U}_μ in such a way that the images of this lower bound under all of the maps, both old and new, are in the corresponding ultrafilters. Clause (6) of Definition 2.10 says precisely that this is possible.

Lemma 4.4. *For any decreasing sequence of conditions $\langle q_n : n < \omega \rangle$ in \mathbb{Q}^δ there is $q \in \mathbb{Q}^\delta$ so that $\forall n < \omega [q \leq q_n]$. Moreover, if $\forall n < \omega [X_{q_{n+1}} = X_{q_n}]$, then $X_q = X_{q_0}$.*

Proof. Assume that we are given a decreasing sequence of conditions $\langle q_n : n < \omega \rangle$ in \mathbb{Q}^δ , i.e. $q_{n+1} \leq q_n$ for $n < \omega$. Define $Y = \bigcup_{n < \omega} X_{q_n}$ and $\gamma = \sup(Y)$. Note that $Y \in [\delta]^{< \aleph_0}$. Also, if $\forall n < \omega [X_{q_{n+1}} = X_{q_n}]$, then $Y = X_{q_0}$. So the moreover part of the lemma holds as long as we find q such that $X_q = Y$. We will consider two cases: either $\gamma \in Y$ or $\gamma \notin Y$.

Case I: $\gamma \in Y$. Then there is $n_0 < \omega$ such that $\gamma \in X_{q_{n_0}}$. So $\gamma = \gamma_{q_{n_0}}$ and note that $\gamma < \delta$ because $X_{q_{n_0}} \subset \delta$. Notice that $\gamma_{q_{n+1}} \geq \gamma_{q_n}$ for every $n < \omega$, so $\gamma_{q_n} = \gamma_{q_{n_0}}$ for every $n \geq n_0$. Also, by Definition 4.1 we know that $\gamma \in X_{q_n}$ for $n \geq n_0$. We apply Lemma 3.9 in such a way that: d_n in Lemma 3.9 is $c_{q_{n+n_0}}$ ($n < \omega$); \mathcal{U} is \mathcal{U}_γ ; π is $\pi_{q_{n_0}, \gamma}$; ψ is $\psi_{q_{n_0}, \gamma}$ and b is $b_{q_{n_0}, \gamma}$. It is easy to see that the hypotheses of Lemma 3.9 are satisfied. So there is $d \in \mathbb{P}$ such that $\pi'' \text{set}(d) \in \mathcal{U}_\gamma$ and $d \leq d_n$ for every $n < \omega$. Now we will prove that the condition $q = \langle d, \gamma, Y, \langle \pi_{q,\alpha} : \alpha \in Y \rangle \rangle$ is as required, where $\pi_{q,\alpha}$ is $\pi_{q_{n+n_0}, \alpha}$ for any $n < \omega$ such that $\alpha \in X_{q_{n+n_0}}$. To show that $q \in \mathbb{Q}^\delta$ note that conditions

(1)–(3) of Definition 4.1 are clearly satisfied. To prove Definition 4.1(4b), fix $\alpha, \beta \in Y$ such that $\alpha \leq \beta$. There is $n < \omega$ such that $\alpha, \beta \in X_{q_{n_0+n}}$. Since $\text{set}(d) \subset^* \text{set}(c_{q_{n_0+n}})$ and $\pi_{q,\alpha} = \pi_{q_{n_0+n},\alpha}$ and $\pi_{q,\beta} = \pi_{q_{n_0+n},\beta}$, by Definition 4.1(4b) for q_{n_0+n} we have $\forall^\infty k \in \text{set}(d) [\pi_{q,\alpha}(k) = \pi_{\beta,\alpha}(\pi_{q,\beta}(k))]$ as required. To see that Definition 4.1(4a) is true take arbitrary $\beta \in Y$. First notice that $\pi''_{q_{n_0},\gamma} \text{set}(d) \in \mathcal{U}_\gamma$. So (4a) is true in case $\beta = \gamma$. If $\beta < \gamma$ consider the set $Z = \pi''_{\gamma,\beta}(\pi''_{q_{n_0},\gamma} \text{set}(d))$. It belongs to \mathcal{U}_β by already proved (4a) for γ and Definition 2.10(5a). However, by already proved (4b) we have $Z \subset^* \pi''_{q,\beta} \text{set}(d)$ which implies that $\pi''_{q,\beta} \text{set}(d) \in \mathcal{U}_\beta$. We still have to prove (4c). Take arbitrary $\alpha \in Y$ and let $n < \omega$ be such that $\alpha \in X_{q_{n_0+n}}$. We know that $\langle \pi_{q_{n_0+n},\alpha}, \psi_{q_{n_0+n},\alpha}, b_{q_{n_0+n},\alpha} \rangle$ is a normal triple, that $d \leq c_{q_{n_0+n}} \leq b_{q_{n_0+n},\alpha}$ and that $\pi_{q,\alpha} = \pi_{q_{n_0+n},\alpha}$. So $d \leq b_{q_{n_0+n},\alpha}$ and $\langle \pi_{q,\alpha}, \psi_{q_{n_0+n},\alpha}, b_{q_{n_0+n},\alpha} \rangle$ is as required.

Case II: $\gamma \notin Y$. Therefore $Y \subset \gamma$. In this case either $\gamma = 0$ or γ is a limit ordinal such that $\text{cof}(\gamma) = \omega$. So there are three subcases: either $\gamma = 0$ or $\gamma < \delta$ and $\text{cof}(\gamma) = \omega$ or $\gamma = \delta$ and $\text{cof}(\gamma) = \omega$.

Subcase IIa: $\gamma = 0$. Since $Y \subset \gamma$ we have $Y = 0$, so $X_{q_0} = 0$ and $\gamma_{q_0} = 0$ and $\gamma_{q_0} \notin X_{q_0}$. So $\delta = \gamma_{q_0} = 0$. In this case all the conditions q_n ($n < \omega$) are of the form $q_n = \langle c_{q_n}, 0, 0, 0 \rangle$. So it is enough to construct condition $c_q \leq c_{q_n}$ ($n < \omega$) because in that case $q = \langle c_q, 0, 0, 0 \rangle$ will satisfy $q \leq q_n$ for every $n < \omega$, and also the moreover part of the lemma. For $n < \omega$ let k_n be such that $c_{q_{n+1}} \leq_{k_n} c_{q_n}$. Define $m_0 = 0$ and $m_{n+1} = \max\{k_n, \max(c_{q_n}(m_n)) + 2\}$ for $n < \omega$. Let $c_q(n) = c_{q_n}(m_n)$ for $n < \omega$. It is obvious that $c_q \in \mathbb{P}$ and $c_q \leq c_{q_n}$ for every $n < \omega$.

Subcase IIb: $\text{cof}(\gamma) = \omega$ and $\gamma < \delta$. We apply Definition 2.10(6) as follows: μ is γ , X is Y and d_n is c_{q_n} ($n < \omega$). For $\alpha \in Y$ let $n < \omega$ be minimal such that $\alpha \in X_{q_n}$. Then we consider π_α to be $\pi_{q_n,\alpha}$, ψ_α to be $\psi_{q_n,\alpha}$ and b_α to be $b_{q_n,\alpha}$ - note that if $m \leq n$ then $\text{set}(c_{q_n}) \subset^* \text{set}(c_{q_m})$, so $\pi''_{q_n,\alpha} \text{set}(c_{q_m}) \in \mathcal{U}_\alpha$, while if $m > n$, then $\pi_{q_n,\alpha} = \pi_{q_m,\alpha}$ and Definition 4.1(4a) implies $\pi''_{q_n,\alpha} \text{set}(c_{q_m}) \in \mathcal{U}_\alpha$, and so Definition 2.10(6a) holds; Definition 2.10(6c) is true because $\langle \pi_\alpha, \psi_\alpha, b_\alpha \rangle$ is a normal triple and $d_n = c_{q_n} \leq b_{q_n,\alpha} = b_\alpha$; to show that Definition 2.10(6b) is satisfied, pick $\alpha, \beta \in Y$ such that $\alpha \leq \beta$, let $n < \omega$ be minimal such that $\alpha \in X_{q_n}$, let $m < \omega$ minimal such that $\beta \in X_{q_m}$ and assume $n \leq m$ (case $m \leq n$ is symmetric). Then $\alpha, \beta \in X_{q_m}$ so according to Definition 4.1(4b) for q_m we have $\forall^\infty k \in \text{set}(d_m) [\pi_{q_m,\alpha}(k) = \pi_{\beta,\alpha}(\pi_{q_m,\beta}(k))]$.

Hypothesis of Definition 2.10(6) is satisfied as explained above. So there are $i^* < \mathfrak{c}$, $d^* \in \mathbb{P}$ and $\pi, \psi \in \omega^\omega$ which satisfy the conclusion of Definition 2.10(6). Now define condition $q = \langle d^*, \gamma, Y \cup \{\gamma\}, \langle \pi_{q,\alpha} : \alpha \in Y \cup \{\gamma\} \rangle \rangle$, where for $\alpha \in Y$, $\pi_{q,\alpha}$ is $\pi_{q_n,\alpha}$ for the minimal $n < \omega$ such that $\alpha \in X_{q_n}$, while $\pi_{q,\gamma}$ is π . When we prove $q \in \mathbb{Q}^\delta$ it will follow easily that $q \leq q_n$ for $n < \omega$. So we check conditions (1)–(4) of Definition 4.1. The only non-trivial condition is (4). First we show (4b). Take any $\alpha, \beta \in Y$ such that $\alpha \leq \beta$. There are two cases, either $\beta = \gamma$ or $\beta \neq \gamma$. If $\beta = \gamma$, then by Definition 2.10(6e) we have $\forall^\infty k \in \text{set}(d^*) [\pi_{q,\alpha}(k) = \pi_{\gamma,\alpha}(\pi_{q,\gamma}(k))]$ as required. If $\beta < \gamma$, then pick $n < \omega$ such that $\alpha, \beta \in X_{q_n}$. Then since $\text{set}(d^*) \subset \text{set}(c_{q_n})$, by Definition 4.1(4b) applied to q_n we have $\forall^\infty k \in \text{set}(d^*) [\pi_{q,\alpha}(k) = \pi_{\beta,\alpha}(\pi_{q,\beta}(k))]$ as required. Next we prove (4a). Let $\alpha \in Y$. If $\alpha = \gamma$, then by Definition 2.10(6d) we have $\pi''_{q,\gamma} \text{set}(d^*) = \text{set}(c_{i^*}^\gamma) \in \mathcal{U}_\gamma$. If $\alpha < \gamma$, then by already proved (4b) we have $\pi''_{\gamma,\alpha}(\pi''_{q,\gamma} \text{set}(d^*)) \subset^* \pi''_{q,\alpha} \text{set}(d^*)$. This together with Definition 2.10(5a) gives $\pi''_{q,\alpha} \text{set}(d^*) \in \mathcal{U}_\alpha$ as required. We still have to prove (4c). Take arbitrary $\alpha \in Y$. If $\alpha = \gamma$, then $\langle \pi_{q,\gamma}, \psi, d^* \rangle$ is itself a normal triple. If $\alpha < \gamma$ let $n < \omega$ be minimal such that $\alpha \in X_{q_n}$. Then $\langle \pi_{q,\alpha}, \psi_{q_n,\alpha}, b_{q_n,\alpha} \rangle$ is a normal triple and $d^* \leq c_{q_n} \leq b_{q_n,\alpha}$ as required.

The situation from the moreover part of the lemma does not occur in this subcase. To see this, suppose otherwise. Then $Y = X_{q_0}$ and $\gamma = \gamma_{q_0}$. Since $\gamma < \delta$, by Definition 4.1(3) $\gamma = \gamma_{q_0} \in X_{q_0} \subset Y$, a contradiction to Case II.

Subcase IIc: $\text{cof}(\gamma) = \omega$ and $\gamma = \delta$. Choose $\langle \gamma_n : n < \omega \rangle$ such that $\sup \{\gamma_n : n < \omega\} = \delta$, and $\gamma_n < \gamma_{n+1}$ and $\gamma_n \in Y$, for every $n < \omega$. Now we apply Lemma 3.8 as follows: \mathcal{U}_n is \mathcal{U}_{γ_n} ($n < \omega$) - note that the \mathcal{U}_{γ_n} 's are distinct rapid P-points; $\pi_{m,n}$ is π_{γ_m, γ_n} ($n \leq m < \omega$) - note that by Definition 2.10(5) conditions (1)–(3) of Lemma 3.8 are satisfied.

As we have explained above, hypothesis of Lemma 3.8 is satisfied, so there are $e \in \mathbb{P}$ and maps π_{δ, γ_n} ($n < \omega$) such that

- (1) $\forall n < \omega$ [$\pi''_{\delta, \gamma_n} \text{set}(e) \in \mathcal{U}_{\gamma_n}$];
- (2) $\forall n \leq m < \omega \forall^\infty k \in \text{set}(e)$ [$(\pi_{\delta, \gamma_n}(k) = \pi_{\gamma_m, \gamma_n}(\pi_{\delta, \gamma_m}(k)))$];
- (3) for every $n < \omega$ there are ψ_{δ, γ_n} and b_{δ, γ_n} such that $\langle \pi_{\delta, \gamma_n}, \psi_{\delta, \gamma_n}, b_{\delta, \gamma_n} \rangle$ is a normal triple and $e \leq b_{\delta, \gamma_n}$.

We will apply Lemma 3.4 as follows: d_n is c_{q_n} for $n < \omega$, e is e , δ is δ and $f = \text{id}$ - note that $\text{cof}(\delta) = \omega$; X is $\{\gamma_n : n < \omega\}$ - note that $\delta = \sup(X)$ as required in Lemma 3.4; the $\pi_{\delta, \alpha}$ are $\pi_{\delta, \alpha}$, for $\alpha \in X$ - note that Lemma 3.4(2) is true by (1)–(3); for $n < \omega$, π_{γ_n} is π_{q_m, γ_n} , b_{γ_n} is b_{q_m, γ_n} , ψ_{γ_n} is ψ_{q_m, γ_n} for the minimal $m < \omega$ such that $\gamma_n \in X_{q_m}$. We have to show that Lemma 3.4(3a)–(3c) are satisfied. First we prove (3a). Fix $n < \omega$ and let $m < \omega$ be minimal such that $\gamma_n \in X_{q_m}$. We will show that $\forall j < \omega$ [$\pi''_{\gamma_n} \text{set}(c_{q_j}) = \pi''_{q_m, \gamma_n} \text{set}(c_{q_j}) \in \mathcal{U}_{\gamma_n}$]. There are two cases: either $j \leq m$ or $j > m$. If $j \leq m$, then $\text{set}(c_{q_m}) \subset^* \text{set}(c_{q_j})$ and by Definition 4.1(4a) applied to q_m , $\pi''_{q_m, \gamma_n} \text{set}(c_{q_j}) \in \mathcal{U}_{\gamma_n}$. If $j > m$, then $\gamma_n \in X_{q_j}$ and $\pi_{q_j, \gamma_n} = \pi_{q_m, \gamma_n}$; so we have that $\pi''_{q_m, \gamma_n} \text{set}(c_{q_j}) \in \mathcal{U}_{\gamma_n}$. Next, we prove (3b). Fix $n \leq m < \omega$. Let $k < \omega$ be minimal such that $\gamma_n \in X_{q_k}$ and $l < \omega$ minimal such that $\gamma_m \in X_{q_l}$. Define $j = \max\{k, l\}$. Then $\gamma_n, \gamma_m \in X_{q_j}$, and $\pi_{q_k, \gamma_n} = \pi_{q_j, \gamma_n}$ and $\pi_{q_l, \gamma_m} = \pi_{q_j, \gamma_m}$. By Definition 4.1(4b) applied to q_j we have that $\forall^\infty k \in \text{set}(c_{q_j})$ [$\pi_{q_j, \gamma_n}(k) = \pi_{\gamma_m, \gamma_n}(\pi_{q_j, \gamma_m}(k))$]. Hence j witnesses (3b). Finally for (3c), fix $n < \omega$ and let $m < \omega$ be minimal such that $\gamma_n \in X_{q_m}$. Since q_m satisfies Definition 4.1(4c) we know that $c_{q_m} \leq b_{q_m, \gamma_n}$ and $\langle \pi_{q_m, \gamma_n}, \psi_{q_m, \gamma_n}, b_{q_m, \gamma_n} \rangle$ is a normal triple. So (3c) is witnessed by $j = m$.

As explained above, the assumptions of Lemma 3.4 are satisfied, so there are $e^*, d^* \in \mathbb{P}$ and $\pi, \psi \in \omega^\omega$ which satisfy conditions (4)–(8) in the conclusion of Lemma 3.4. Consider $q = \langle d^*, \delta, Y, \langle \pi_{q, \alpha} : \alpha \in Y \rangle \rangle$, where for $\alpha \in Y$, $\pi_{q, \alpha} = \pi_{q_m, \alpha}$ for the minimal $m < \omega$ such that $\alpha \in X_{q_m}$. Note that for each $n < \omega$, $\pi_{\gamma_n} = \pi_{q, \gamma_n}$. If we prove that $q \in \mathbb{Q}^\delta$ it will follow easily that $q \leq q_n$ for $n < \omega$ and that q satisfies the moreover part of the lemma. So we check the properties (1)–(4) of Definition 4.1. Conditions (1)–(3) are clearly satisfied. We prove (4a)–(4c). First we show that (4b) is true. Let $\alpha, \beta \in Y$ be such that $\alpha \leq \beta$. Let $m < \omega$ and $k < \omega$ be minimal with $\alpha \in X_{q_m}$ and $\beta \in X_{q_k}$ respectively. Put $l = \max\{m, k\}$, and note that $\pi_{q, \alpha} = \pi_{q_l, \alpha}$ and $\pi_{q, \beta} = \pi_{q_l, \beta}$. So by definition 4.1(4b) applied to q_l and by the fact that $\text{set}(d^*) \subset^* \text{set}(c_{q_l})$, $\forall^\infty k^* \in \text{set}(d^*)$ [$\pi_{q, \alpha}(k^*) = \pi_{\beta, \alpha}(\pi_{q, \beta}(k^*))$], as required. Now we prove (4a). Fix $\alpha \in Y$, and let $n < \omega$ be such that $\gamma_n \geq \alpha$. Note that $\pi''_{\delta, \gamma_n} \text{set}(e^*) \in \mathcal{U}_{\gamma_n}$ and that $\pi''_{\delta, \gamma_n} \text{set}(e^*) \subset^* \pi''_{\gamma_n} \text{set}(d^*)$. Thus $\pi''_{q, \gamma_n} \text{set}(d^*) \in \mathcal{U}_{\gamma_n}$, and so $\pi''_{\gamma_n, \alpha} \pi''_{q, \gamma_n} \text{set}(d^*) \in \mathcal{U}_\alpha$. By (4b), $\pi''_{\gamma_n, \alpha} \pi''_{q, \gamma_n} \text{set}(d^*) \subset^* \pi''_{q, \alpha} \text{set}(d^*)$, whence $\pi''_{q, \alpha} \text{set}(d^*) \in \mathcal{U}_\alpha$ as needed. Finally for (4c), fix $\alpha \in Y$ and let $m < \omega$ be minimal such that $\alpha \in X_{q_m}$. Then setting $b_{q, \alpha} = b_{q_m, \alpha}$ and $\psi_{q, \alpha} = \psi_{q_m, \alpha}$ fulfills (4c). \square

\mathbb{Q}^δ is required to be $< \mathfrak{c}$ closed when carrying out the constructing under MA. This can be proved in the same way as Lemma 4.4 by using the appropriate generalizations of the lemmas from Section 3 and the regularity of \mathfrak{c} , which follows from MA.

We next turn towards showing that various sets are dense in \mathbb{Q}^δ . These are the dense sets we will want to meet when building our “sufficiently generic” filter for \mathbb{Q}^δ . Meeting these dense sets will ensure that the sequences $\langle c_i^\delta : i < \mathfrak{c} \rangle$ and $\langle \pi_{\delta, \alpha} : \alpha \leq \delta \rangle$, which we intend to read off from the generic filter, will satisfy the conditions of Definition 2.10 when they are added to S . The first density condition states that for each $q \in \mathbb{Q}^\delta$, there is a

$q' \leq q$ such that $c_{q'}$ is a “fast” subsequence of c_q . This is needed to ensure that \mathcal{U}_δ is rapid, and it will also play a role in ensuring that it is an ultrafilter.

Lemma 4.5. *For $q \in \mathbb{Q}^\delta$ and strictly increasing $f \in \omega^\omega$ there is $q' \leq q$ such that $X_q = X_{q'}$ and that for every $n < \omega$ there is $m \geq f(n)$ so that $c_{q'}(n) = c_q(m)$. Moreover, there is $q'' \leq q'$ such that for every $n < \omega$ we have $c_{q''}(n) \in [c_{q'}(n)]^{n+1}$.*

Proof. We first show how to get q' . We will distinguish two cases: when $\gamma_q = \delta$ and when $\gamma_q < \delta$.

Case I: $\gamma_q = \delta$. We know that $X_q \subset \delta$, $\sup(X_q) = \gamma_q = \delta$ and $|X_q| \leq \omega$, so either $\delta = 0$ or δ is a limit ordinal with $\text{cof}(\delta) = \omega$.

Subcase Ia: $\gamma_q = \delta$ and $\delta = 0$. In this case q is of the form $\langle c_q, 0, 0, 0 \rangle$. For every $n < \omega$, let $c_{q'}(n) = c_q(f(n))$. Then $c_{q'} \in \mathbb{P}$ because f is strictly increasing. Also it is clear that $c_{q'} \leq c_q$. Consequently $q' = \langle c_{q'}, 0, 0, 0 \rangle \leq q$ is as required.

Subcase Ib: $\gamma_q = \delta$ and $\text{cof}(\delta) = \omega$. We apply Lemma 3.4 in such a way that: e is c_q and δ is δ ; d_n is c_q for $n < \omega$; f is f ; X is X_q ; maps π_α are maps $\pi_{q,\alpha}$ ($\alpha \in X_q$); maps $\pi_{\delta,\alpha}$ are maps $\pi_{q,\alpha}$ ($\alpha \in X_q$). The conditions of Lemma 3.4 are clearly satisfied. Hence, there is $e^* \leq_0 c_q$ such that for every $n < \omega$ there is $m \geq f(n)$ so that $e^*(n) \subset c_q(m)$. We will construct numbers k_n by induction on n so that for every $n < \omega$ there is $m \geq n$ so that $e^*(m) \subset c_q(k_n)$ and that $\text{set}(e^*) \subset \bigcup_{n < \omega} c_q(k_n)$. Let k_0 be such that $e^*(0) \subset c_q(k_0)$. Now assume that numbers k_m have been chosen for every $m \leq n$, and define k_{n+1} as follows: let l be maximal such that $e^*(l) \subset c_q(k_n)$ and define k_{n+1} as the unique number such that $e^*(l+1) \subset c_q(k_{n+1})$. Now for every $n < \omega$ define $c_{q'}(n) = c_q(k_n)$. We will prove that the condition $q' = \langle c_{q'}, \delta, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle$ is as required. Since for every n , $e^*(n) \subset c_q(m)$ for $m \geq f(n)$ we have that $k_n \geq f(n)$, so $\forall n < \omega \exists l \geq f(n) c_{q'}(n) = c_q(l)$, as required in the statement of the lemma. By Remark 4.3, in order to prove $q' \in \mathbb{Q}^\delta$ and $q' \leq q$ it is enough to prove that q' satisfies Definition 4.1(4a). So pick $\alpha \in X_q$. Since $\text{set}(e^*) \subset \bigcup_{n < \omega} c_q(k_n) = \text{set}(c_{q'})$ we know that $\pi''_{q,\alpha} \text{set}(e^*) \subset^* \pi''_{q,\alpha} \text{set}(c_{q'})$, but since $\pi''_{q,\alpha} \text{set}(e^*) \in \mathcal{U}_\alpha$ we have $\pi''_{q,\alpha} \text{set}(c_{q'}) \in \mathcal{U}_\alpha$ as required.

Case II: $\gamma_q < \delta$. Note that by Definition 4.1(3) $\gamma_q \in X_q$. Let n_0 be such that $c_q \leq_{n_0} b_{q,\gamma_q}$. Then $a = \pi''_{q,\gamma_q} \text{set}(c_q) \llbracket n_0 \rrbracket \in \mathcal{U}_{\gamma_q}$. Now by Lemma 2.9, for each $n < \omega$, $m_n = \max\{m < \omega : \pi''_{q,\gamma_q} c_q(m) = \{a(n)\}\}$ is well-defined and $m_n < m_{n+1}$. As \mathcal{U}_{γ_q} is rapid, there is $Y \in \mathcal{U}_{\gamma_q}$ such that $Y \subset a$ and for each $n \in \omega$, there is $l_n \geq f(n)$ such that $Y(n) = a(l_n)$. Now it is clear that for each $n \in \omega$, $m_{l_n} \geq l_n \geq f(n)$. Define $c_{q'}(n) = c_q(m_{l_n})$. It is clear that $c_{q'} \in \mathbb{P}$ and that $\pi''_{q,\gamma_q} \text{set}(c_{q'}) = Y$. So by Remark 4.3, we will finish the proof by showing that $q' = \langle c_{q'}, \gamma_q, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle$ satisfies Definition 4.1(4a). So let $\alpha \in X_q$. We know that $\pi''_{q,\gamma_q} \text{set}(c_{q'}) = Y \in \mathcal{U}_{\gamma_q}$ so $\pi''_{\gamma_q,\alpha} Y \in \mathcal{U}_\alpha$. Now we have that $\pi''_{q,\alpha} \text{set}(c_{q'}) =^* \pi''_{\gamma_q,\alpha} (\pi''_{q,\gamma_q} \text{set}(c_{q'})) = \pi''_{\gamma_q,\alpha} Y \in \mathcal{U}_\alpha$ as required.

To get q'' , define $c_{q''}$ as follows: for every $n < \omega$ pick an arbitrary $c_{q''}(n) \in [c_{q'}(n)]^{n+1}$. This is possible because $|c_{q'}(n)| \geq n+1$. Let $q'' = \langle c_{q''}, \gamma_q, X_q, \langle \pi_{q,\alpha} : \alpha \in X_q \rangle \rangle$. To see that $q'' \in \mathbb{Q}^\delta$ note that conditions (1)–(3), (4b), and (4c) of Definition 4.1 are clearly satisfied. Condition (4a) holds because for every $\alpha \in X_q$, $\pi''_{q,\alpha} \text{set}(c_{q''}) =^* \pi''_{q,\alpha} \text{set}(c_{q'})$. \square

The next lemma ensures that for any given $Y \in \mathcal{P}(\omega)$, every condition in \mathbb{Q}^δ has an extension that “decides” Y . This will make \mathcal{U}_δ into an ultrafilter.

Lemma 4.6. *For every $q \in \mathbb{Q}^\delta$ and for every $Y \in \mathcal{P}(\omega)$ there is $q' \leq q$ such that $X_{q'} = X_q$ and that $\text{set}(c_{q'}) \subset Y$ or $\text{set}(c_{q'}) \subset \omega \setminus Y$.*

Proof. In the same way as in the proof of Lemma 4.5 we distinguish the following cases: either $\gamma_q = \delta = 0$ or $\gamma_q = \delta$ and $\text{cof}(\delta) = \omega$ or $\gamma_q < \delta$.

Case I: $\gamma_q = \delta$. As already mentioned this case has two subcases.

Subcase Ia: $\gamma_q = \delta = 0$. In this case q is of the form $q = \langle c_q, 0, 0, 0 \rangle$. Recall the notation introduced in Section 2: for $Y \subset \omega$, define $Y^0 = Y$ and $Y^1 = \omega \setminus Y$. For $i \in 2$ consider the sets $A_i = \{n < \omega : |c_q(n) \cap Y^i| \geq (n+1)/2\}$. Note $A_0 \cup A_1 = \omega$ so either A_0 or A_1 is infinite. Assume without loss of generality that A_0 is infinite. Then $|c_q(A_0(2n+1)) \cap Y| \geq n+1$ for every $n < \omega$. Define $c_{q'} \in \mathbb{P}$ as follows: for $n < \omega$ let $c_{q'}(n) = [c_q(A_0(2n+1)) \cap Y]^{n+1}$. It is easy to see that $c_{q'} \in \mathbb{P}$ and $c_{q'} \leq c_q$. So for $q' = \langle c_{q'}, 0, 0, 0 \rangle$ we have $q' \leq q$ and $\text{set}(c_{q'}) \subset Y$. If we assumed A_1 is infinite, then we would obtain $\text{set}(c_{q'}) \subset \omega \setminus Y$.

Subcase Ib: $\gamma_q = \delta$ and $\text{cof}(\delta) = \omega$. First according to Lemma 4.5 there is $q' \leq q$ such that $X_{q'} = X_q$ and that for every $n < \omega$ there is $m \geq 2^{n+1}$ such that $c_{q'}(n) = c_q(m)$. Note that this implies that for every $n < \omega$ we have $|c_{q'}(n)| \geq 2^{n+1}$. Let us consider two sets $A_i = \{n < \omega : |c_{q'}(n) \cap Y^i| \geq 2^n\}$ ($i = 0, 1$). Fix $\alpha \in X_{q'}$. Because $q' \in \mathbb{Q}^\delta$ we have $\pi''_{q', \alpha} \text{set}(c_{q'}) \in \mathcal{U}_\alpha$. So since $A_0 \cup A_1 = \omega$ we have that

$$\pi''_{q', \alpha} \left(\bigcup_{n \in A_0} c_{q'}(n) \right) \cup \pi''_{q', \alpha} \left(\bigcup_{n \in A_1} c_{q'}(n) \right) \in \mathcal{U}_\alpha.$$

Since \mathcal{U}_α is an ultrafilter, there is $i_\alpha \in 2$ such that $\pi''_{q', \alpha} \left(\bigcup_{n \in A_{i_\alpha}} c_{q'}(n) \right) \in \mathcal{U}_\alpha$. Now that we have defined i_α for every $\alpha \in X_{q'}$, pick ordinals $\beta_n \in X_{q'}$ so that the sequence $\langle \beta_n : n < \omega \rangle$ is strictly increasing and cofinal in δ . There is $K \in [\omega]^{<\aleph_0}$ and $i \in \{0, 1\}$ so that $i_{\beta_n} = i$ for every $n \in K$. Now pick any $\beta \in X_{q'}$. Because K is infinite and $\langle \beta_n : n < \omega \rangle$ is cofinal in δ , there is $n \in K$ so that $\beta_n > \beta$. We know that $\pi''_{q', \beta_n} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \in \mathcal{U}_{\beta_n}$. But according to Definition 2.10(5a) and Definition 4.1(4b) we have

$$\pi''_{q', \beta} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \supset^* \pi''_{\beta_n, \beta} \left(\pi''_{q', \beta_n} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \right) \in \mathcal{U}_\beta$$

which shows that for every $\beta \in X_{q'}$ we have that $\pi''_{q', \beta} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \in \mathcal{U}_\beta$. Now define $d \in \mathbb{P}$ as follows: for every $n < \omega$ pick arbitrary $d(n) \in [c_{q'}(A_i(n)) \cap Y^i]^{n+1}$. The sequence $d = \langle d(n) : n < \omega \rangle$ belongs to \mathbb{P} because A_i was chosen in such a way that for $n < \omega$ we have $|c_{q'}(A_i(n)) \cap Y^i| \geq 2^n \geq n+1$. Finally, we will show that $q'' = \langle d, \gamma_{q'}, X_q, \langle \pi_{q', \alpha} : \alpha \in X_{q'} \rangle \rangle$ is as required (note $X_{q''} = X_{q'} = X_q$). It is enough to prove that $q'' \in \mathbb{Q}^\delta$, because then $q'' \leq q$ and $\text{set}(c_{q''}) = \text{set}(d) \subset Y^i$ easily follows. By Remark 4.3 it is enough to show that Definition 4.1(4a) is satisfied. We show that $\pi''_{q'', \beta} \text{set}(d) \supset^* \pi''_{q'', \beta} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \in \mathcal{U}_\beta$ holds for $\beta \in X_{q'}$. Consider the set $C = \pi''_{q'', \beta} \left(\bigcup_{n \in A_i} c_{q'}(n) \right) \setminus \pi''_{q'', \beta} \text{set}(d)$. Let $m < \omega$ be such that $c_{q'} \leq_m b_{q'', \beta}$. Note that for any $n \geq m$ we have $\pi''_{q'', \beta} d(n) = \pi''_{q'', \beta} c_{q'}(A_i(n))$. This implies that $C \subset \pi''_{q'', \beta} \left(\bigcup_{n < A_i(m)} c_{q'}(n) \right)$ which shows that $|C| < \omega$ as required.

Case II: $\gamma_q < \delta$. Let $q' \leq q$ be such that $X_{q'} = X_q$ and that for each $n \in \omega$, $c_{q'}(n) = c_q(m)$ for some $m \geq 2n+1$. Note that $\gamma_q = \gamma_{q'} \in X_{q'}$ and that for each $n \in \omega$, $|c_{q'}(n)| \geq 2n+2$. For $i \in 2$, let $A_i = \{n \in \omega : |Y^i \cap c_{q'}(n)| \geq n+1\}$. Note that $\omega = A_0 \cup A_1$. Therefore $\left(\pi''_{q', \gamma_q} \bigcup_{n \in A_0} c_{q'}(n) \right) \cup \left(\pi''_{q', \gamma_q} \bigcup_{n \in A_1} c_{q'}(n) \right) = \pi''_{q', \gamma_q} \text{set}(c_{q'}) \in \mathcal{U}_{\gamma_q}$. Fix $i \in 2$ such that $\pi''_{q', \gamma_q} \bigcup_{n \in A_i} c_{q'}(n) \in \mathcal{U}_{\gamma_q}$. Then A_i is infinite and $|c_{q'}(A_i(k)) \cap Y^i| \geq A_i(k) + 1 \geq k+1$, for each $k \in \omega$. Choose $c_{q''}(k) \in [c_{q'}(A_i(k)) \cap Y^i]^{k+1}$. Then $c_{q''} = \langle c_{q''}(k) : k \in \omega \rangle \in \mathbb{P}$ and $c_{q''} \leq c_{q'}$. Moreover, $\pi''_{q', \gamma_q} \bigcup_{n \in A_i} c_{q'}(n) \subset^* \pi''_{q', \gamma_q} \text{set}(c_{q''})$. Thus $\pi''_{q', \gamma_q} \text{set}(c_{q''}) \in \mathcal{U}_{\gamma_q}$. Furthermore, $\pi''_{\gamma_q, \alpha} \pi''_{q', \gamma_q} \text{set}(c_{q''}) \subset^* \pi''_{q', \alpha} \text{set}(c_{q''})$, for each $\alpha \in X_{q'}$. So we also have that $\pi''_{q', \alpha} \text{set}(c_{q''}) \in \mathcal{U}_\alpha$, for each $\alpha \in X_{q'}$. Therefore by Remark 4.3 $q'' = \langle c_{q''}, \gamma_q, X_{q'}, \langle \pi_{q', \alpha} : \alpha \in X_{q'} \rangle \rangle$ is as required. \square

We would like it to be the case that for each $\beta < \delta$, there is a q in our ‘‘sufficiently generic’’ filter over \mathbb{Q}^δ with $\beta \in X_q$ because we would like to read the map $\pi_{\delta, \beta}$ from

the filter. So we next prove that for each $\beta < \delta$, every $q \in \mathbb{Q}^\delta$ has an extension q' with $\beta \in X_{q'}$. But let us first interject two technical lemmas that are easy to prove.

Lemma 4.7. *For $q \in \mathbb{Q}^\delta$, $\alpha \in X_q$ and $a \in \mathcal{U}_\alpha$ there is $q' \leq q$ such that $X_{q'} = X_q$ and $\pi''_{q',\alpha} \text{set}(c_{q'}) \subset a$.*

Proof. Consider the set $b = \pi''_{q,\alpha} \text{set}(c_q)$. By Definition 4.1(4a) $b \in \mathcal{U}_\alpha$, which implies that $a \cap b \in \mathcal{U}_\alpha$. Denote $V = \pi_{q,\alpha}^{-1}(a \cap b)$. By Lemma 4.6 there is $q' \leq q$ such that $\text{set}(c_{q'}) \subset V$ or $\text{set}(c_{q'}) \subset \omega \setminus V$. Assume that $\text{set}(c_{q'}) \subset \omega \setminus V$. By Definition 4.1(4a) we have $\pi''_{q',\alpha} \text{set}(c_{q'}) \in \mathcal{U}_\alpha$ and $\pi_{q',\alpha} = \pi_{q,\alpha}$. So $(\pi''_{q',\alpha} \text{set}(c_{q'})) \cap (a \cap b) = \emptyset$ which is impossible. Hence $\text{set}(c_{q'}) \subset V$ which implies that $\pi''_{q',\alpha} \text{set}(c_{q'}) \subset a \cap b$ as required. \square

Lemma 4.8. *Let $q \in \mathbb{Q}^\delta$, $\beta \in \delta$, and $Y = X_q \cup \{\beta\}$. There is $q' \leq q$ such that $X_{q'} = X_q$ and that for every ζ, ξ, μ satisfying $\mu \in X_{q'}$, $\zeta, \xi \in Y$, and $\zeta \leq \xi \leq \mu$, there is $N < \omega$ such that for every $k, l \in \text{set}(c_{q'})$ if $N \leq k \leq l$, then $\pi_{\mu,\zeta}(\pi_{q',\mu}(k)) = \pi_{\xi,\zeta}(\pi_{\mu,\xi}(\pi_{q',\mu}(k)))$ and $\pi_{\mu,\xi}(\pi_{q',\mu}(k)) \leq \pi_{\mu,\xi}(\pi_{q',\mu}(l))$.*

Proof. Let $V = \{(\zeta, \xi, \mu) : \mu \in X_q \wedge \zeta, \xi \in Y \wedge \zeta \leq \xi \leq \mu\}$. V is countable, so let $\{(\zeta_n, \xi_n, \mu_n) : n < \omega\}$ enumerate it, possibly with repetitions. Build by induction on n a decreasing sequence $\langle q_n : n \in \omega \rangle \subset \mathbb{Q}^\delta$ such that $\forall n \in \omega [X_{q_{n+1}} = X_{q_n}]$. Let $q_0 = q$. Fix $n \in \omega$, and suppose that $q_n \leq q$ is given. By the definition of a δ -generic sequence, there exists $a_n \in \mathcal{U}_{\mu_n}$ such that $\forall k^* \in a_n [\pi_{\mu_n, \zeta_n}(k^*) = \pi_{\xi_n, \zeta_n}(\pi_{\mu_n, \xi_n}(k^*))]$ and $\forall k^*, l^* \in a_n [k^* \leq l^* \implies \pi_{\mu_n, \xi_n}(k^*) \leq \pi_{\mu_n, \xi_n}(l^*)]$. Apply Lemma 4.7 to $q_n \in \mathbb{Q}^\delta$, $\mu_n \in X_{q_n}$, and $a_n \in \mathcal{U}_{\mu_n}$, to find $q_{n+1} \leq q_n$ such that $\pi''_{q_{n+1}, \mu_n} \text{set}(c_{q_{n+1}}) \subset a_n$ and $X_{q_{n+1}} = X_{q_n}$. This concludes the construction of $\langle q_n : n \in \omega \rangle$. Find $q' \in \mathbb{Q}^\delta$ such that $\forall n \in \omega [q' \leq q_n]$ and $X_{q'} = X_{q_0} = X_q$. We check that q' is as needed. Fix $n < \omega$. As $\mu_n \in X_{q'}$ and $q' \leq q_{n+1}$, there is N such that for all $k, l \in \text{set}(c_{q'})$, if $N \leq k \leq l$, then $k, l \in \text{set}(c_{q_{n+1}})$ and $\pi_{q', \mu_n}(k) \leq \pi_{q', \mu_n}(l)$. Fixing any such k and l , let $k^* = \pi_{q', \mu_n}(k)$ and $l^* = \pi_{q', \mu_n}(l)$. Then $k^*, l^* \in a_n$ and $k^* \leq l^*$. Therefore, $\pi_{\mu_n, \zeta_n}(k^*) = \pi_{\xi_n, \zeta_n}(\pi_{\mu_n, \xi_n}(k^*))$ and $\pi_{\mu_n, \xi_n}(k^*) \leq \pi_{\mu_n, \xi_n}(l^*)$, as needed. \square

Lemma 4.9. *For $q \in \mathbb{Q}^\delta$ and $\beta < \delta$ there is $q' \in \mathbb{Q}^\delta$ such that $q' \leq q$ and $\beta \in X_{q'}$.*

Proof. Assume $\beta \notin X_q$. According to Lemma 4.8 applied to q and β there is $q^* \leq q$ such that $X_{q^*} = X_q$ and that for every ζ, ξ, μ satisfying $\mu \in X_{q^*}$, $\zeta, \xi \in X_{q^*} \cup \{\beta\}$ and $\zeta \leq \xi \leq \mu$ there is $N < \omega$ such that for every $k, l \in \text{set}(c_{q^*})$ if $N \leq k \leq l$, then $\pi_{\mu,\zeta}(\pi_{q^*,\mu}(k)) = \pi_{\xi,\zeta}(\pi_{\mu,\xi}(\pi_{q^*,\mu}(k)))$ and $\pi_{\mu,\xi}(\pi_{q^*,\mu}(k)) \leq \pi_{\mu,\xi}(\pi_{q^*,\mu}(l))$. In the same way as in the proof of Lemma 4.5 we have the following cases: either $\gamma_{q^*} = \delta = 0$ or $\gamma_{q^*} = \delta$ and $\text{cof}(\delta) = \omega$ or $\gamma_{q^*} < \delta$.

Case I: $\gamma_{q^*} = \delta$. As we mentioned above there are two subcases.

Subcase Ia: $\gamma_{q^*} = \delta = 0$. Note that in this case the statement is vacuous because there is no $\beta < \delta$.

Subcase Ib: $\gamma_{q^*} = \delta$ and $\text{cof}(\delta) = \omega$. Since $\text{sup}(X_{q^*}) = \gamma_{q^*} = \delta$ and $\beta < \delta$, let $\gamma^* \in X_{q^*}$ be minimal such that $\beta \leq \gamma^*$. Let $m < \omega$ be minimal such that $c_{q^*} \leq_m b_{q^*, \gamma^*}$ and for all $k, l \in \text{set}(c_{q^*}) \llbracket m \rrbracket$, if $k \leq l$, then $\pi_{\gamma^*, \beta}(\pi_{q^*, \gamma^*}(k)) \leq \pi_{\gamma^*, \beta}(\pi_{q^*, \gamma^*}(l))$. Define

$$q' = \langle c_{q^*}, \delta, X_{q^*} \cup \{\beta\}, \langle \pi_{q^*, \alpha} : \alpha \in X_{q^*} \cup \{\beta\} \rangle \rangle,$$

where $\pi_{q^*, \beta}$ is as follows: for $k \in \text{set}(c_{q^*}) \llbracket m \rrbracket$ let $\pi_{q^*, \beta}(k) = \pi_{\gamma^*, \beta}(\pi_{q^*, \gamma^*}(k))$, while $\pi_{q^*, \beta}(k) = 0$ otherwise. It suffices to prove that $q' \in \mathbb{Q}^\delta$ because it is then easy to see that $q' \leq q^*$ and $\beta \in X_{q'}$ hold. Properties (1)–(3) are clearly satisfied. So we check (4). First it is clear that (4a) holds by the definition of $\pi_{q^*, \beta}$ and by the fact that $c_{q'} = c_{q^*}$. Next, we check (4b). Pick arbitrary $\alpha, \gamma \in X_{q'}$ such that $\alpha \leq \gamma$. We will distinguish four cases: either $(\alpha \neq \beta \text{ and } \gamma \neq \beta)$, or $(\alpha = \beta = \gamma)$, or $(\gamma = \beta \text{ and } \alpha \neq \beta)$, or $(\alpha = \beta \text{ and } \gamma \neq \beta)$. First, if $\alpha \neq \beta$ and $\gamma \neq \beta$, then (4b) holds because $q^* \in \mathbb{Q}^\delta$ and

$\alpha, \gamma \in X_{q^*}$. Next, if $\alpha = \beta = \gamma$, then (4b) trivially holds. Now assume that $\gamma = \beta$ and $\alpha \neq \beta$. Then $\alpha \in X_{q^*}$. There exists k_1 such that for each $k \in \text{set}(c_{q^*}) \llbracket k_1 \rrbracket$ the following hold: $\pi_{q^*, \gamma}(k) = \pi_{\gamma^*, \gamma}(\pi_{q^*, \gamma^*}(k))$, $\pi_{\gamma^*, \alpha}(\pi_{q^*, \gamma^*}(k)) = \pi_{\gamma, \alpha}(\pi_{\gamma^*, \gamma}(\pi_{q^*, \gamma^*}(k)))$, and $\pi_{q^*, \alpha}(k) = \pi_{\gamma^*, \alpha}(\pi_{q^*, \gamma^*}(k))$. Thus for every $k \in \text{set}(c_{q^*}) \llbracket k_1 \rrbracket$ we have $\pi_{q^*, \alpha}(k) = \pi_{\gamma^*, \alpha}(\pi_{q^*, \gamma^*}(k)) = \pi_{\gamma, \alpha}(\pi_{\gamma^*, \gamma}(\pi_{q^*, \gamma^*}(k))) = \pi_{\gamma, \alpha}(\pi_{q^*, \gamma}(k))$ as required. Finally assume that $\alpha = \beta$ and $\gamma \neq \beta$. Then $\gamma \in X_{q^*}$ and $\beta \leq \gamma$. By minimality of γ^* , $\gamma^* \leq \gamma$. As before, there exists $k_0 \in \omega$ such that for each $k \in \text{set}(c_{q^*}) \llbracket k_0 \rrbracket$ the following hold: $\pi_{q^*, \alpha}(k) = \pi_{\gamma^*, \alpha}(\pi_{q^*, \gamma^*}(k))$, $\pi_{\gamma, \alpha}(\pi_{q^*, \gamma}(k)) = \pi_{\gamma^*, \alpha}(\pi_{\gamma^*, \gamma}(\pi_{q^*, \gamma^*}(k)))$, and $\pi_{q^*, \gamma^*}(k) = \pi_{\gamma, \gamma^*}(\pi_{q^*, \gamma}(k))$. Thus for every $k \in \text{set}(c_{q^*}) \llbracket k_0 \rrbracket$ we have $\pi_{q^*, \alpha}(k) = \pi_{\gamma^*, \alpha}(\pi_{q^*, \gamma^*}(k)) = \pi_{\gamma^*, \alpha}(\pi_{\gamma, \gamma^*}(\pi_{q^*, \gamma}(k))) = \pi_{\gamma, \alpha}(\pi_{q^*, \gamma}(k))$ as required. So (4b) holds. Finally, we check (4c). If $\alpha \in X_{q^*}$ then (4c) is true because $q^* \in \mathbb{Q}^\delta$. Let us now define $b_{q^*, \beta}$ and $\psi_{q^*, \beta}$. Put $b_{q^*, \beta} = c_{q^*}$. For each $m^* \geq m$, π_{q^*, γ^*} is constant on $c_{q^*}(m^*)$ because $c_{q^*} \leq_m b_{q^*, \gamma^*}$. So for each $m^* \geq m$, $\pi_{q^*, \beta}$ is constant on $c_{q^*}(m^*)$. Also $\pi_{\gamma^*, \beta} \circ \pi_{q^*, \gamma^*}$ is increasing on $\text{set}(c_{q^*}) \llbracket m \rrbracket$. So for each $m^* \geq m$ define $\psi_{q^*, \beta}(m^*) = \pi_{\gamma^*, \beta}(\pi_{q^*, \gamma^*}(k)) = \pi_{q^*, \beta}(k)$, for an arbitrary $k \in c_{q^*}(m^*)$. When $m^* < m$, $\pi_{q^*, \beta}$ is constantly equal to 0 on $c_{q^*}(m^*)$. So set $\psi_{q^*, \beta}(m^*) = 0$, for $m^* < m$. It is clear that $\langle \pi_{q^*, \beta}, \psi_{q^*, \beta}, b_{q^*, \beta} \rangle$ is a normal triple with $c_{q^*} \leq b_{q^*, \beta}$.

Case II: $\gamma_{q^*} < \delta$. Note that $\gamma_{q^*} \in X_{q^*}$. There are two subcases: when $\beta < \gamma_{q^*}$ and when $\gamma_{q^*} < \beta$. If $\beta < \gamma_{q^*}$, then define $\pi_{q^*, \beta}$ as follows. Let m_1 be such that the following two things hold: $c_{q^*} \leq_{m_1} b_{q^*, \gamma_{q^*}}$; and for any $k, l \in \text{set}(c_{q^*}) \llbracket m_1 \rrbracket$, if $k \leq l$, then $\pi_{\gamma_{q^*}, \beta}(\pi_{q^*, \gamma_{q^*}}(k)) \leq \pi_{\gamma_{q^*}, \beta}(\pi_{q^*, \gamma_{q^*}}(l))$. For $k \in \text{set}(c_{q^*}) \llbracket m_1 \rrbracket$, define $\pi_{q^*, \beta}(k) = \pi_{\gamma_{q^*}, \beta}(\pi_{q^*, \gamma_{q^*}}(k))$, while for $k \notin \text{set}(c_{q^*}) \llbracket m_1 \rrbracket$, define $\pi_{q^*, \beta}(k) = 0$. We will prove that $q' = \langle c_{q^*}, \gamma_{q^*}, X_{q^*} \cup \{\beta\}, \langle \pi_{q^*, \alpha} : \alpha \in X_{q^*} \cup \{\beta\} \rangle \rangle$ is as required. It is enough to show that $q' \in \mathbb{Q}^\delta$, because then we will have that $q' \leq q^*$ and $\beta \in X_{q'}$. Clearly, conditions (1)–(3) are satisfied. To see that (4) is true, note that (4a) is clear from the definition of $\pi_{q^*, \beta}$ and from that fact that $c_{q'} = c_{q^*}$. Next, we check (4b) for q' . Fix $\alpha, \gamma \in X_{q'}$ such that $\alpha \leq \gamma$. There are again four cases: either ($\beta \neq \alpha$ and $\beta \neq \gamma$), or ($\alpha = \beta = \gamma$), or ($\gamma = \beta$ and $\alpha \neq \beta$), or ($\gamma \neq \beta$ and $\alpha = \beta$). If $\beta \neq \alpha$ and $\beta \neq \gamma$, then the statement follows directly from Definition 4.1(4b) applied to q^* . The case when $\alpha = \beta = \gamma$ is trivial. Next, consider the case when $\gamma = \beta$ and $\alpha \neq \beta$. Then $\alpha \in X_{q^*}$. There exists k_2 such that for each $k \in \text{set}(c_{q^*}) \llbracket k_2 \rrbracket$ the following hold: $\pi_{q^*, \gamma}(k) = \pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k))$, $\pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{\gamma, \alpha}(\pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k)))$, and $\pi_{q^*, \alpha}(k) = \pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k))$. Thus for any $k \in \text{set}(c_{q^*}) \llbracket k_2 \rrbracket$, $\pi_{\gamma, \alpha}(\pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k))) = \pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{q^*, \alpha}(k)$, as needed. Finally suppose that $\gamma \neq \beta$ and $\alpha = \beta$. Then $\gamma \in X_{q^*}$. As before, there exists k_3 such that for each $k \in \text{set}(c_{q^*}) \llbracket k_3 \rrbracket$ the following hold: $\pi_{q^*, \alpha}(k) = \pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k))$, $\pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{\gamma, \alpha}(\pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k)))$, and $\pi_{q^*, \gamma}(k) = \pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k))$. Thus for $k \in \text{set}(c_{q^*}) \llbracket k_3 \rrbracket$, $\pi_{q^*, \alpha}(k) = \pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{\gamma, \alpha}(\pi_{\gamma_{q^*}, \gamma}(\pi_{q^*, \gamma_{q^*}}(k))) = \pi_{\gamma, \alpha}(\pi_{q^*, \gamma}(k))$, as required. So (4b) is checked, and we now check (4c) for q' . If $\alpha \in X_{q^*}$, then (4c) is satisfied for q' because it was satisfied for q^* . It remains to define $b_{q^*, \beta}$ and $\psi_{q^*, \beta}$. Put $b_{q^*, \beta} = c_{q^*}$. Note that for each $m^* \geq m_1$, $\pi_{q^*, \gamma_{q^*}}$ is constant on $c_{q^*}(m^*)$ because $c_{q^*} \leq_{m_1} b_{q^*, \gamma_{q^*}}$. So for each $m^* \geq m_1$, $\pi_{q^*, \beta}$ is constant on $c_{q^*}(m^*)$. Also $\pi_{\gamma_{q^*}, \beta} \circ \pi_{q^*, \gamma_{q^*}}$ is increasing on $\text{set}(c_{q^*}) \llbracket m_1 \rrbracket$. So for $m^* \geq m_1$, define $\psi_{q^*, \beta}(m^*) = \pi_{\gamma_{q^*}, \beta}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{q^*, \beta}(k)$, for an arbitrary $k \in c_{q^*}(m^*)$. When $m^* < m_1$, $\pi_{q^*, \beta}$ is constantly equal to 0 on $c_{q^*}(m^*)$. So define $\psi_{q^*, \beta}(m^*) = 0$, for $m^* < m_1$. It is clear that $\langle \pi_{q^*, \beta}, \psi_{q^*, \beta}, b_{q^*, \beta} \rangle$ is a normal triple and that $c_{q^*} \leq b_{q^*, \beta}$. Hence q' is as required.

Now consider the case when $\beta > \gamma_{q^*}$. For each $\alpha \in X_{q^*}$, since $\alpha \leq \gamma_{q^*} < \beta$, by Definition 2.10(5b) pick $a_\alpha \in \mathcal{U}_\beta$ so that $\forall k \in a_\alpha [\pi_{\beta, \alpha}(k) = \pi_{\gamma_{q^*}, \alpha}(\pi_{\beta, \gamma_{q^*}}(k))]$. Since X_{q^*} is countable and \mathcal{U}_β is a P-point there is $a \in \mathcal{U}_\beta$ such that $a \subset^* a_\alpha$ for every $\alpha \in X_{q^*}$. Then we apply Definition 2.10(3) with β, α being γ_{q^*} , c_{q^*} being d , π_1

being $\pi_{q^*, \gamma_{q^*}}$, b_1 being $b_{q^*, \gamma_{q^*}}$ and ψ_1 being $\psi_{q^*, \gamma_{q^*}}$ and a . Note that hypothesis of Definition 2.10(3) are satisfied. By Definition 2.10(3) there are $b \in \mathcal{U}_\beta$, $\pi, \psi \in \omega^\omega$ and $d^* \leq_0 c_{q^*}$ so that $b \subset^* a$, $\langle \pi, \psi, d^* \rangle$ is a normal triple, $\pi'' \text{set}(d^*) = b$ and $\forall k \in \text{set}(d^*) [\pi_{q^*, \gamma_{q^*}}(k) = \pi_{\beta, \gamma_{q^*}}(\pi(k))]$. Denote $\pi_{q^*, \beta} = \pi$ and $\psi_{q^*, \beta} = \psi$. Now define $q' = \langle d^*, \beta, X_{q^*} \cup \{\beta\}, \langle \pi_{q^*, \alpha} : \alpha \in X_{q^*} \cup \{\beta\} \rangle \rangle$. It is easy to see that if we prove that $q' \in \mathbb{Q}^\delta$, then $q' \leq q^*$ and $\beta \in X_{q'}$ follow. So we check conditions of Definition 4.1. Note that conditions (1)–(3) are clearly true. We still have to check Definition 4.1(4). First note that (4c) is satisfied for $\alpha \in X_{q^*}$ because $d^* \leq c_{q^*}$, while it is true for β because $\langle \pi, \psi, d^* \rangle$ is a normal triple. To see that (4b) is true let $\alpha, \gamma \in X_{q^*} \cup \{\beta\}$ be such that $\alpha \leq \gamma$. There are three cases: either $\alpha \neq \beta$ and $\gamma \neq \beta$ or $\alpha = \beta$ or $\gamma = \beta$. First note that if $\alpha = \beta$, then it must also be $\gamma = \beta$ and the statement holds. If $\alpha \neq \beta$ and $\gamma \neq \beta$ then by Definition 4.1 and because $\text{set}(d^*) \subset^* \text{set}(c_{q^*})$ we have $\forall^\infty k \in \text{set}(d^*) [\pi_{q^*, \alpha}(k) = \pi_{\gamma, \alpha}(\pi_{q^*, \gamma}(k))]$. If $\gamma = \beta$ then because $b \subset^* a \subset^* a_\alpha$ and $\pi'' \text{set}(d^*) = b$ we have that there is $k_0 < \omega$ such that for every $k \in \text{set}(d^*) \llbracket k_0 \rrbracket$ we have $\pi_{q^*, \alpha}(k) = \pi_{\gamma_{q^*}, \alpha}(\pi_{q^*, \gamma_{q^*}}(k)) = \pi_{\gamma_{q^*}, \alpha}(\pi_{\beta, \gamma_{q^*}}(\pi(k))) = \pi_{\beta, \alpha}(\pi(k)) = \pi_{\beta, \alpha}(\pi_{q^*, \beta}(k))$ as required. To see that (4a) is true note that $\pi''_{q^*, \beta} \text{set}(d^*) = b \in \mathcal{U}_\beta$ and consequently $\pi''_{\beta, \alpha} (\pi''_{q^*, \beta} \text{set}(d^*)) \in \mathcal{U}_\beta$ for any $\alpha \in X_{q^*}$. Together with already proved (4b), this implies $\pi''_{\beta, \alpha} (\pi''_{q^*, \beta} \text{set}(d^*)) \subset^* \pi''_{q^*, \alpha} \text{set}(d^*) \in \mathcal{U}_\beta$ for $\alpha \in X_{q^*}$. \square

The next lemma ensures that we can “kill” unwanted Tukey maps. That is, if $\beta < \delta$ and $\phi : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$ is a monotone map that is a potential witness for the unwanted Tukey reduction $\mathcal{U}_\delta \leq_T \mathcal{U}_\beta$, then we would like every condition in \mathbb{Q}^δ to have an extension forcing that ϕ is not such a witness.

Lemma 4.10. *For any $q \in \mathbb{Q}^\delta$, any $\beta < \delta$ and any monotone $\phi : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$, if for every $A \in \mathcal{U}_\beta$, $\phi(A) \neq \emptyset$ then there is $q' \leq q$ such that $\beta \in X_{q'}$ and that for every $A \in \mathcal{U}_\beta$ we have $\phi(A) \not\subset \text{set}(c_{q'})$.*

Proof. By Lemma 4.9 there is $q' \leq q$ such that $\beta \in X_{q'}$ and by Lemma 4.5 there is $q'' \leq q' \leq q$ such that for every $n < \omega$ there is $m \geq 2n + 1$ such that $c_{q''}(n) = c_{q'}(m)$. For every $n < \omega$ choose sets $d_1(n)$ and $d_2(n)$ which are elements of $[c_{q''}(n)]^{n+1}$ and are such that $d_1(n) \cap d_2(n) = \emptyset$. This can be done because $|c_{q''}(n)| \geq 2n + 2$. Note that both $q_1 = \langle d_1, \gamma_{q''}, X_{q''}, \langle \pi_{q'', \alpha} : \alpha \in X_{q''} \rangle \rangle$ and $q_2 = \langle d_2, \gamma_{q''}, X_{q''}, \langle \pi_{q'', \alpha} : \alpha \in X_{q''} \rangle \rangle$ belong to \mathbb{Q}^δ and that $q_1, q_2 \leq q'' \leq q$. Now we consider two cases: either for every $A \in \mathcal{U}_\beta$, $\phi(A) \not\subset \text{set}(d_1)$, or there is some $A \in \mathcal{U}_\beta$ such that $\phi(A) \subset \text{set}(d_1)$. If for every $A \in \mathcal{U}_\beta$, $\phi(A) \not\subset \text{set}(d_1)$, then q_1 is as required. Otherwise, q_2 is as required because $\text{set}(d_1) \cap \text{set}(d_2) = \emptyset$ and ϕ is monotone. \square

Note that q' forces what we want because it forces $\text{set}(c_{q'}) \in \mathcal{U}_\delta$. Hence it forces that the image of \mathcal{U}_β under ϕ is not cofinal in \mathcal{U}_δ . It is also worth noting that the descriptive complexity of ϕ plays no role in the proof of Lemma 4.10. So Theorem 1.4 is only needed for bounding the number of relevant maps.

The next lemma is needed for ensuring clause (6) of Definition 2.10, and hence it is only relevant when $\text{cof}(\delta) = \omega$. It follows by a direct application of Lemma 3.4.

Lemma 4.11. *Suppose that $\text{cof}(\delta) = \omega$, $q \in \mathbb{Q}^\delta$ is such that $\gamma_q = \delta$, $\langle d_j : j < \omega \rangle$ is a decreasing sequence in \mathbb{P} , $X \subset X_q$ is such that $\text{sup}(X) = \delta$ and that $\langle \pi_\alpha : \alpha \in X \rangle$ is a sequence of maps in ω^ω satisfying:*

- (1) $\forall \alpha \in X \forall j < \omega [\pi''_\alpha \text{set}(d_j) \in \mathcal{U}_\alpha]$;
- (2) $\forall \alpha, \beta \in X [\alpha \leq \beta \Rightarrow \exists j < \omega \forall^\infty k \in \text{set}(d_j) [\pi_\alpha(k) = \pi_{\beta, \alpha}(\pi_\beta(k))]]$;
- (3) for all $\alpha \in X$ there are $j < \omega$ and $\psi_\alpha \in \omega^\omega$ and $b_\alpha \in \mathbb{P}$ such that $\langle \pi_\alpha, \psi_\alpha, b_\alpha \rangle$ is a normal triple and $d_j \leq b_\alpha$.

Then there are $q' \leq q$, $d^ \in \mathbb{P}$ and $\pi : \omega \rightarrow \omega$ such that:*

- (4) $\forall j < \omega [d^* \leq d_j]$ and $\text{set}(c_{q'}) = \pi'' \text{set}(d^*)$;
- (5) $\forall \alpha \in X \forall^\infty k \in \text{set}(d^*) [\pi_\alpha(k) = \pi_{q',\alpha}(\pi(k))]$;
- (6) there is ψ for which $\langle \pi, \psi, d^* \rangle$ is a normal triple.

Proof. We will use Lemma 3.4 where: $\delta, X, \langle d_j : j < \omega \rangle$ and $\pi_\alpha (\alpha \in X)$ are as in the statement of this lemma, $f = \text{id}$; e is c_q ; for $\alpha \in X$ map $\pi_{\delta,\alpha}$ is $\pi_{q,\alpha}$ ($\alpha \in X$). So there are e^*, d^* and π satisfying properties (4)–(8) of the conclusion of Lemma 3.4. We will show that $q' = \langle e^*, \delta, X_{q'}, \langle \pi_{q,\alpha} : \alpha \in X_{q'} \rangle \rangle, d^*$ and π are as required. The conditions (4)–(6) will be witnessed by conditions (5)–(8) in the conclusion of the Lemma 3.4. By Remark 4.3, in order to finish the proof we only have to show that Definition 4.1(4a) holds for q' . First assume that $\alpha \in X$. Then by Lemma 3.4(7), $\pi''_{q,\alpha} \text{set}(e^*) \in \mathcal{U}_\alpha$. Now assume that $\alpha \in X_q \setminus X$. Let $\alpha' \in X$ be such that $\alpha' > \alpha$. Then $\pi''_{\alpha',\alpha} (\pi''_{q,\alpha'} \text{set}(e^*)) \in \mathcal{U}_\alpha$. Also we have $\pi''_{\alpha',\alpha} (\pi''_{q,\alpha'} \text{set}(e^*)) \subset^* \pi''_{q,\alpha} \text{set}(e^*)$. These observations together give us $\pi''_{q,\alpha} \text{set}(e^*) \in \mathcal{U}_\alpha$ as required. \square

Next we show how to make sure that \mathcal{U}_δ is rapid. This lemma follows from a direct application of Lemma 4.5.

Lemma 4.12. *Suppose that $\delta < \omega_2$, that $q \in \mathbb{Q}^\delta$ and that $f \in \omega^\omega$ is a strictly increasing function. There is $q' \leq q$ such that for every $n < \omega$ we have $\text{set}(c_{q'})(n) \geq f(n)$.*

Proof. According to Lemma 4.5 there is $q' \leq q$ so that for every $n < \omega$ there is $m \geq f(s(n+1))$ such that $c_{q'}(n) \in [c_q(m)]^{n+1}$. We will prove that q' is as required. So fix $n < \omega$, and let $k < \omega$ be such that $\text{set}(c_{q'})(n) \in c_{q'}(k)$. Equivalently $s(k) \leq n < s(k+1)$ which implies $\text{set}(c_{q'})(n) \geq \text{set}(c_{q'})(s(k))$. Since for some $m \geq f(s(k+1))$ we have $c_{q'}(k) \subset c_q(m)$ and $\text{set}(c_{q'})(n) \in c_{q'}(k)$, then by Remark 2.6(1), $\text{set}(c_{q'})(n) \geq m \geq f(s(k+1)) > f(n)$, the last inequality being true because f is an increasing function. So we showed that $\text{set}(c_{q'})(n) \geq f(n)$ as required. \square

We now come to the final density lemma. This lemma ensures that clause (3) of Definition 2.10 can be satisfied during the construction of \mathcal{U}_δ . One of the cases in its proof makes use of Lemma 3.4.

Lemma 4.13. *Let $q \in \mathbb{Q}^\delta$, $\pi_1, \psi_1 \in \omega^\omega$, $b_1, d \in \mathbb{P}$, and $\alpha < \delta$ be such that $\langle \pi_1, \psi_1, b_1 \rangle$ is a normal triple, $d \leq b_1$, and $\pi''_1 \text{set}(d) \in \mathcal{U}_\alpha$. Then there are $q^* \leq q$, $d^* \leq d$, $\pi, \psi \in \omega^\omega$ such that $\langle \pi, \psi, d^* \rangle$ is a normal triple, $\alpha \in X_{q^*}$, $\pi'' \text{set}(d^*) = \text{set}(c_{q^*})$ and $\forall k \in \text{set}(d^*) [\pi_1(k) = \pi_{q^*,\alpha}(\pi(k))]$.*

Proof. By Lemma 4.9 there is $q_0 \leq q$ such that $\alpha \in X_{q_0}$. In the same way as in the proof of Lemma 4.5 we have the following cases: either $\gamma_{q_0} = \delta = 0$ or $\gamma_{q_0} = \delta$ and $\text{cof}(\delta) = \omega$ or $\gamma_{q_0} < \delta$.

Case I: $\gamma_{q_0} = \delta$. As we mentioned above there are two subcases.

Subcase Ia: $\gamma_{q_0} = \delta = 0$. Then the statement is vacuous because there is no $\alpha < \delta$.

Subcase Ib: $\gamma_{q_0} = \delta$ and $\text{cof}(\delta) = \omega$. In particular δ is limit ordinal. There is $q_1 \leq q_0$ such that $X_{q_1} = X_{q_0}$ and that q_1 satisfies conclusion of Lemma 4.8. Note $\sup(X_{q_1}) = \delta$. So pick an increasing sequence $\langle \alpha_n : n < \omega \rangle$ such that $\alpha_0 = \alpha$, $\sup \{\alpha_n : n < \omega\} = \delta$ and $\alpha_n \in X_{q_1}$ for $n < \omega$. Build by induction sequences $\langle d_n : n < \omega \rangle$ and $\langle \pi_{\alpha_n} : n < \omega \rangle$ satisfying the following for each $n < \omega$:

- (1) $d_0 = d$, $\pi_{\alpha_0} = \pi_1$, and $\forall m \leq n [d_n \leq d_m]$;
- (2) $\pi''_{\alpha_n} \text{set}(d_n) \in \mathcal{U}_{\alpha_n}$ and $\forall m \leq n \forall^\infty k \in \text{set}(d_n) [\pi_{\alpha_m}(k) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_n}(k))]$;
- (3) if $n > 0$, then there is $\psi_{\alpha_n} \in \omega^\omega$ such that $\langle \pi_{\alpha_n}, \psi_{\alpha_n}, d_n \rangle$ is a normal triple.

Put $d_0 = d$ and $\pi_{\alpha_0} = \pi_1$. Fix $n \in \omega$ and assume that d_n and π_{α_n} are given satisfying (1)–(3). To get d_{n+1} we apply Definition 2.10(3) with $\alpha = \alpha_n$, $\beta = \alpha_{n+1}$, $\pi_1 = \pi_{\alpha_n}$, $d = d_n$, and if $n = 0$, then $\psi_1 = \psi_1$ and $b_1 = b_1$, while if $n > 0$, then

$\psi_1 = \psi_{\alpha_n}$ and $b_1 = d_n$. Note that in all cases the hypothesis of Definition 2.10(3) is satisfied. Let a in Definition 2.10(3) be $\pi''_{d_1, \alpha_{n+1}} \text{set}(c_{q_1})$. Then there are $b \in \mathcal{U}_{\alpha_{n+1}}$, $\psi_{\alpha_{n+1}}$, $\pi_{\alpha_{n+1}} \in \omega^\omega$ and $d_{n+1} \leq d_n$ such that $b \subset^* \pi''_{q_1, \alpha_{n+1}} \text{set}(c_{q_1})$, $\pi''_{\alpha_{n+1}} \text{set}(d_{n+1}) = b$, $\forall k \in \text{set}(d_{n+1}) [\pi_{\alpha_n}(k) = \pi_{\alpha_{n+1}, \alpha_n}(\pi_{\alpha_{n+1}}(k))]$ and that $\langle \pi_{\alpha_{n+1}}, \psi_{\alpha_{n+1}}, d_{n+1} \rangle$ is a normal triple. We will prove that d_{n+1} and $\pi_{\alpha_{n+1}}$ satisfy (1)–(3). (1) is clear. Second, we have $\pi''_{\alpha_{n+1}} \text{set}(d_{n+1}) = b \in \mathcal{U}_{\alpha_{n+1}}$. Next, we check that for every $m \leq n+1$, $\forall^\infty k \in \text{set}(d_{n+1}) [\pi_{\alpha_m}(k) = \pi_{\alpha_{n+1}, \alpha_m}(\pi_{\alpha_{n+1}}(k))]$. We distinguish two cases: either $m = n+1$ or $m \leq n$. If $m = n+1$, then since $\pi_{\alpha_{n+1}, \alpha_{n+1}} = \text{id}$, for every $k \in \text{set}(d_{n+1})$ we have $\pi_{\alpha_{n+1}}(k) = \pi_{\alpha_{n+1}, \alpha_{n+1}}(\pi_{\alpha_{n+1}}(k))$. If $m \leq n$, then it is easy to find a $k_0 \in \omega$ so that for every $k \in \text{set}(d_{n+1}) \llbracket k_0 \rrbracket$ the following hold: $\pi_{\alpha_n}(k) = \pi_{\alpha_{n+1}, \alpha_n}(\pi_{\alpha_{n+1}}(k))$, $\pi_{\alpha_m}(k) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_n}(k))$, and $\pi_{\alpha_{n+1}, \alpha_m}(\pi_{\alpha_{n+1}}(k)) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_{n+1}, \alpha_n}(\pi_{\alpha_{n+1}}(k)))$. Hence for each $k \in \text{set}(d_{n+1}) \llbracket k_0 \rrbracket$, $\pi_{\alpha_m}(k) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_n}(k)) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_{n+1}, \alpha_n}(\pi_{\alpha_{n+1}}(k))) = \pi_{\alpha_{n+1}, \alpha_m}(\pi_{\alpha_{n+1}}(k))$, as required. Fourth, $\langle \pi_{\alpha_{n+1}}, \psi_{\alpha_{n+1}}, d_{n+1} \rangle$ is a normal triple. So the sequences $\langle d_n : n < \omega \rangle$ and $\langle \pi_{\alpha_n} : n < \omega \rangle$ are as required.

Next we apply Lemma 3.4 in such a way that δ is δ , f is id , $X = \{\alpha_n : n < \omega\}$, e is c_{q_1} , π_{δ, α_n} is π_{q_1, α_n} for $n < \omega$, d_n is d_n for $n < \omega$, π_{α_n} is π_{α_n} for $n < \omega$, b_{δ, α_n} and ψ_{δ, α_n} are b_{q_1, α_n} and ψ_{q_1, α_n} for $n < \omega$. Note that $\text{cof}(\delta) = \omega$, $\text{sup}(X) = \delta$, $X \subset \delta$ and that Lemma 3.4(2) is satisfied because q_1 satisfies Definition 4.1(4). So we still have to prove that condition (3) of Lemma 3.4 is satisfied. First we prove (3b). Fix $m \leq n < \omega$. By the construction of π_{α_n} we know that $\forall^\infty k \in \text{set}(d_n) [\pi_{\alpha_m}(k) = \pi_{\alpha_n, \alpha_m}(\pi_{\alpha_n}(k))]$ as required. To see (3c) take $n < \omega$ and note that $\langle \pi_{\alpha_n}, \psi_{\alpha_n}, d_n \rangle$ is a normal triple. Next we prove (3a). We have to prove that for every $m, n < \omega$, $\pi''_{\alpha_n} \text{set}(d_m) \in \mathcal{U}_{\alpha_n}$. We consider two cases: either $m \leq n$ or $m > n$. If $m \leq n$ note $\text{set}(d_n) \subset^* \text{set}(d_m)$ so since by construction $\pi''_{\alpha_n} \text{set}(d_n) \in \mathcal{U}_{\alpha_n}$ we have $\pi''_{\alpha_n} \text{set}(d_m) \in \mathcal{U}_{\alpha_n}$. If $m > n$ then by construction $\pi''_{\alpha_m} \text{set}(d_m) \in \mathcal{U}_{\alpha_m}$. By already proved (3b) $\pi''_{\alpha_m, \alpha_n}(\pi''_{\alpha_m} \text{set}(d_m)) =^* \pi''_{\alpha_n} \text{set}(d_m)$. By Definition 2.10(5a), $\pi''_{\alpha_m, \alpha_n}(\pi''_{\alpha_m} \text{set}(d_m)) \in \mathcal{U}_{\alpha_n}$ so we have $\pi''_{\alpha_n} \text{set}(d_m) \in \mathcal{U}_{\alpha_n}$ as required.

So all conditions of Lemma 3.4 are satisfied. Hence, there are e' , d' , π' and ψ' satisfying conditions (4)–(8) of the conclusion of Lemma 3.4. In particular, by (6) we know that there is k_1 such that $\forall k \in \text{set}(d') \llbracket k_1 \rrbracket [\pi_{\alpha_0}(k) = \pi_{q_1, \alpha_0}(\pi'(k))]$. Let $n < \omega$ be such that $(\pi')'' d'(k_1) \subset e'(n)$, and k_2 such that $\text{set}(e') \llbracket n+1 \rrbracket = (\pi')'' \text{set}(d') \llbracket k_2 \rrbracket$. Since $\langle \pi', \psi', d' \rangle$ is a normal triple, k_2 is well defined and $k_2 \geq k_1$. Now let d^* be defined by $d^*(k) = d'(k_2 + k)$ ($k < \omega$), let e^* be defined by $e^*(k) = e'(n + k + 1)$ ($k < \omega$). Note that $e^* \leq_0 e' \leq_0 c_{q_1}$, $\text{set}(e^*) =^* \text{set}(e')$ and $d^* \leq d'$. Define also $\pi \in \omega^\omega$ as follows: for $k \in \text{set}(d^*)$ let $\pi(k) = \pi'(k)$ and $\pi(k) = 0$ otherwise. Let $\psi \in \omega^\omega$ be defined by $\psi(k) = \psi'(k_2 + k)$ for $k < \omega$. Note that $\langle \pi, \psi, d^* \rangle$ is a normal triple and that $\pi'' \text{set}(d^*) = \text{set}(e^*)$. Define $q^* = \langle e^*, \delta, X_{q_1}, \langle \pi_{q_1, \alpha} : \alpha \in X_{q_1} \rangle \rangle$. Since $e^* \leq c_{q_1}$, by Remark 4.3, in order to show $q^* \leq q_1$ we only have to prove that q^* satisfies Definition 4.1(4a). First we show that it holds for all α_n ($n < \omega$). Take $n < \omega$. By (7) of the conclusion of Lemma 3.4 and because $\text{set}(e^*) =^* \text{set}(e')$ we have $\pi''_{q_1, \alpha_n} \text{set}(e^*) \in \mathcal{U}_{\alpha_n}$. Now we prove (4a). Let $\alpha \in X_{q_1}$. Pick $\alpha_n \geq \alpha$. By Remark 4.3, q^* satisfies (4b) so $\pi''_{\alpha_n, \alpha}(\pi''_{q_1, \alpha_n} \text{set}(e^*)) =^* \pi_{q_1, \alpha} \text{set}(e^*)$. By Definition 2.10(5a) we know $\pi''_{\alpha_n, \alpha}(\pi''_{q_1, \alpha_n} \text{set}(e^*)) \in \mathcal{U}_\alpha$. Hence $\pi''_{q_1, \alpha} \text{set}(e^*) \in \mathcal{U}_\alpha$ as required. We will show that q^* , d^* , π and ψ satisfy conclusion of this lemma. First, $\langle \pi, \psi, d^* \rangle$ is a normal triple. Second, $\alpha \in X_{q_0} = X_{q_1} = X_{q^*}$. Third, $\pi'' \text{set}(d^*) = \text{set}(e^*)$. Fourth, for every $k \in \text{set}(d^*)$ we know that $k \in \text{set}(d') \llbracket k_2 \rrbracket$, so $\pi_1(k) = \pi_\alpha(k) = \pi_{q_1, \alpha}(\pi(k))$. Note that $\pi_1 = \pi_\alpha$ by definition of π_{α_0} . So q^* is as required.

Case II: $\gamma_{q_0} < \delta$. Let n_0 be such that $c_{q_0} \leq_{n_0} b_{q_0, \gamma_{q_0}}$ and that for every $k \in \text{set}(c_{q_0}) \llbracket n_0 \rrbracket$ we have $\pi_{q_0, \alpha}(k) = \pi_{\gamma_{q_0}, \alpha}(\pi_{q_0, \gamma_{q_0}}(k))$

Claim 4.14. *There are $d' \leq d$, $b \subset \pi''_{q_0, \gamma_{q_0}} \text{set}(c_{q_0}) \llbracket n_0 \rrbracket$ and $\pi_2, \psi_2 \in \omega^\omega$ such that $\pi''_2 \text{set}(d') = b$, $\forall k \in \text{set}(d') [\pi_1(k) = \pi_{\gamma_{q_0}, \alpha}(\pi_2(k))]$ and that $\langle \pi_2, \psi_2, d' \rangle$ is a normal triple.*

Proof. We will consider two cases: either $\alpha = \gamma_{q_0}$ or $\alpha < \gamma_{q_0}$. If $\alpha < \gamma_{q_0}$ then we apply Definition 2.10(3) with $\alpha = \alpha$, $\beta = \gamma_{q_0}$, $\pi_1 = \pi_1$, $\psi_1 = \psi_1$, $b_1 = b_1$, $d = d$ and $a = \pi''_{q_0, \gamma_{q_0}} \text{set}(c_{q_0}) \llbracket n_0 \rrbracket$. Note that hypothesis of Definition 2.10(3) are satisfied, and that b in the conclusion of Definition 2.10(3) can be chosen so that $b \subset a$, and not only $b \subset^* a$. Hence there are $b \in \mathcal{U}_{\gamma_{q_0}}$, $\pi, \psi \in \omega^\omega$ and $d' \leq_0 d$ so that $\langle \pi, \psi, d' \rangle$ is a normal triple, $\pi'' \text{set}(d') = b \subset a$ and $\forall k \in \text{set}(d') [\pi_1(k) = \pi_{\gamma_{q_0}, \alpha}(\pi(k))]$ as required.

If $\alpha = \gamma_{q_0}$, first let l be such that $d \leq_l b_1$. Put $b = \pi'_1 \text{set}(d) \llbracket l \rrbracket \cap \pi''_{q_0, \gamma_{q_0}} \text{set}(c_{q_0}) \llbracket n_0 \rrbracket$ and note $b \in \mathcal{U}_{\gamma_{q_0}}$ and $b \subset \pi''_{q_0, \gamma_{q_0}} \text{set}(c_{q_0}) \llbracket n_0 \rrbracket$. Put $F_n = \{m < \omega : \pi'_1 d(m) = \{b(n)\}\}$. By Lemma 2.9, $L_n = \max(F_n)$ is well defined and $L_n < L_{n+1}$ is true for $n < \omega$. Define d' as follows: for $n < \omega$ let $d'(n) = d(L_n)$. Since $L_n < L_{n+1}$ ($n < \omega$) we know that $d' \in \mathbb{P}$. Define π_2 as follows: for $k \notin \text{set}(d')$ let $\pi_2(k) = \pi_1(k)$ while $\pi_2(k) = 0$ otherwise. Since $L_n \geq l$ for every $n < \omega$, we can define L'_n such that $d(L_n) \subset b_1(L'_n)$. Then $\pi'_1 b_1(L'_n) = \{\psi_1(L'_n)\}$. Define $\psi_2(n) = \psi_1(L'_n)$. Then it is easy to see that $\langle \pi_2, \psi_2, d' \rangle$ is a normal triple. We know that $\pi''_2 \text{set}(d') = b$ because for every $n < \omega$, $\pi''_2 d'(n) = \pi'_1 d(L_n) = \pi'_1 b_1(L'_n) = \{b(n)\}$. To see that $\forall k \in \text{set}(d') [\pi_1(k) = \pi_{\gamma_{q_0}, \alpha}(\pi_2(k))]$ note that $\gamma_{q_0} = \alpha$ and $\pi_{\alpha, \alpha} = \text{id}$ so for every $k \in \text{set}(d')$ we have that $\pi_1(k) = \pi_{\alpha, \alpha}(\pi_2(k))$ is true. \square

Now that we have b with the required properties, since $\mathcal{U}_{\gamma_{q_0}}$ is rapid, there is $c \in \mathcal{U}_{\gamma_{q_0}}$ so that for every $n < \omega$ there is $m \geq t(n+1)$ so that $c(n) = b(m)$. We will build e^* , d^* , ψ , and π so that the following hold: $e^* \leq_0 c_{q_0}$, $d^* \leq_0 d'$, $\pi'' \text{set}(d^*) = e^*$, $\pi''_{q_0, \gamma_{q_0}} \text{set}(e^*) = c$, $\forall k \in \text{set}(d^*) [\pi_2(k) = \pi_{q_0, \gamma_{q_0}}(\pi(k))]$, and $\langle \pi, \psi, d^* \rangle$ is a normal triple. For each $n < \omega$ define $M_n = \max\{m < \omega : \pi''_{q_0, \gamma_{q_0}} c_{q_0}(m) = \{c(n)\}\}$ and $K_n = \max\{m < \omega : \pi''_2 d'(m) = \{c(n)\}\}$. By Lemma 2.9, $M_n \geq n_0$, $M_{n+1} > M_n$ and $K_{n+1} > K_n$ ($n < \omega$). We show that $K_n \geq t(n+1)$ for every $n < \omega$. Define $l_n = \max\{m < \omega : \pi''_2 d'(m) = b(n)\}$ ($n < \omega$), and note that by Lemma 2.9 numbers l_n are well defined and that $l_{n+1} > l_n$ ($n < \omega$). Hence $l_{n+1} \geq n+1$ ($n < \omega$). Fix $n < \omega$. Then there is $v_n \geq t(n+1)$ such that $c(n) = b(v_n)$. So $K_n = l_{v_n} \geq l_{t(n+1)} \geq t(n+1)$. Define e^* as follows: for $n < \omega$ let $e^*(n) \in [c_{q_0}(M_n)]^{n+1}$. Define d^* as follows. For each $n' < \omega$ choose a sequence $\langle d^*(n) : s(n') \leq n < s(n'+1) \rangle$ in such a way that for all $s(n') \leq n < s(n'+1)$, $d^*(n) \in [d'(K_{n'})]^{n+1}$ and for all $s(n') \leq n < n+1 < s(n'+1)$, $\max(d^*(n)) < \min(d^*(n+1))$. Next, define $\pi \in \omega^\omega$ as follows: for $k \notin \text{set}(d^*)$ let $\pi(k) = 0$, while for $k \in \text{set}(d^*)$ let $\pi(k) = \text{set}(e^*)(n)$ where n is such that $k \in d^*(n)$. Let $\psi \in \omega^\omega$ be defined as $\psi(n) = \text{set}(e^*)(n)$ for every $n < \omega$. Note that $\langle \pi, \psi, d^* \rangle$ is a normal triple because $\pi'' d^*(n) = \{\psi(n)\} = \{\text{set}(e^*)(n)\}$ for $n < \omega$. To show that e^* , d^* , π and ψ are as required, we still have to show that for every $k \in \text{set}(d^*) [\pi_2(k) = \pi_{q_0, \gamma_{q_0}}(\pi(k))]$. Fix $k \in \text{set}(d^*)$. Let n be such that $k \in d^*(n)$ and let m be such that $s(m) \leq n < s(m+1)$. Then $k \in d'(K_m)$ so $\pi_2(k) = c(m)$. Also $\pi(k) = \text{set}(e^*)(n) \in e^*(m) \subset c_{q_0}(M_m)$ so $\pi_{q_0, \gamma_{q_0}}(\pi(k)) = c(m) = \pi_2(k)$ as required. Define $q^* = \langle e^*, \gamma_{q_0}, X_{q_0}, \langle \pi_{q_0, \alpha} : \alpha \in X_{q_0} \rangle \rangle$. Since $e^* \leq_0 c_{q_0}$, by Remark 4.3, in order to prove $q^* \in \mathbb{Q}^\delta$ and $q^* \leq q_0$ it is enough to show that q^* satisfies property (4a) of Definition 4.1. So let $\beta \in X_{q_0}$. There are two cases: either $\beta = \gamma_{q_0}$ or $\beta < \gamma_{q_0}$. If $\beta = \gamma_{q_0}$ then $\pi''_{q_0, \gamma_{q_0}} \text{set}(e^*) = c \in \mathcal{U}_\beta$. If $\beta < \gamma_{q_0}$ then by Remark 4.3 property (4b) of Definition 4.1 holds for q^* so $\pi''_{\gamma_{q_0}, \beta}(\pi''_{q_0, \gamma_{q_0}} \text{set}(e^*)) = \pi''_{q_0, \beta} \text{set}(e^*)$. Also, by Definition 2.10(5a), $\pi''_{\gamma_{q_0}, \beta}(\pi''_{q_0, \gamma_{q_0}} \text{set}(e^*)) \in \mathcal{U}_\beta$ so $\pi''_{q_0, \beta} \text{set}(e^*) \in \mathcal{U}_\beta$. Hence $q^* \in \mathbb{Q}^\delta$ and $q^* \leq q_0$. Finally, we prove that q^* satisfies conclusion of this lemma. By the choice of q_0 we have $\alpha \in X_{q_0}$. By the choice of d^* and e^* we have $\pi'' \text{set}(d^*) =$

$\text{set}(e^*)$. We already explained why $\langle \pi, \psi, d^* \rangle$ is a normal triple. So we still have to prove that $\forall k \in \text{set}(d^*) [\pi_1(k) = \pi_{q^*, \alpha}(\pi(k))]$. By Claim 4.14 and since $d^* \leq_0 d'$ we have $\forall k \in \text{set}(d^*) [\pi_1(k) = \pi_{\gamma_{q_0}, \alpha}(\pi_2(k))]$. We also proved $\forall k \in \text{set}(d^*) [\pi_2(k) = \pi_{q_0, \gamma_{q_0}}(\pi(k))]$. Since $M_n \geq n_0$ for every $n < \omega$ and $\pi'' \text{set}(d^*) = \text{set}(e^*)$, we also have that $\forall k \in \text{set}(d^*) [\pi_{q_0, \alpha}(\pi(k)) = \pi_{\gamma_{q_0}, \alpha}(\pi_{q_0, \gamma_{q_0}}(\pi(k)))]$. From these three equations we get $\pi_1(k) = \pi_{\gamma_{q_0}, \alpha}(\pi_2(k)) = \pi_{\gamma_{q_0}, \alpha}(\pi_{q_0, \gamma_{q_0}}(\pi(k))) = \pi_{q_0, \alpha}(\pi(k))$ as required. Hence q^* satisfies conclusion of this lemma. \square

The proofs of Lemmas 4.5–4.13 go through with no essential modifications under MA. Of course the proofs would depend on the fact that \mathbb{Q}^δ would be $< \mathfrak{c}$ closed and the generalized form of Lemma 3.4 would hold in this context. The inductive construction occurring in Subcase Ib of the proof of Lemma 4.13 would need to be of length ξ , for some $\xi < \mathfrak{c}$. The limit stages of this inductive construction can be passed by appealing to the generalized forms of Lemma 3.4 and Clause (6) of Definition 2.10.

5. A LONG CHAIN

We now have all the tools necessary for constructing the desired chain of P-points. As our construction requires CH, we assume $2^{\aleph_0} = \aleph_1$ in this section. The chain of length ω_2 will be obtained from an ω_2 -generic sequence.

Theorem 5.1 (CH). *There is an ω_2 -generic sequence.*

Proof. We build by induction sequence $\langle S_{\delta'} : \delta' \leq \omega_2 \rangle$ such that for each $\delta' \leq \omega_2$:

- (1) $S_{\delta'}$ is δ' -generic;
- (2) $\forall \gamma < \delta' [S_\gamma = S_{\delta'} \upharpoonright \gamma]$.

For $\delta' = 0$, let $S_0 = \langle 0, 0 \rangle$. Next assume that δ' is a limit ordinal and that for every $\gamma < \delta'$, we are given S_γ as required. Define

$$S_{\delta'} = \left\langle \bigcup_{\gamma < \delta'} \langle c_i^\alpha : \alpha < \gamma \wedge i < \mathfrak{c} \rangle, \bigcup_{\gamma < \delta'} \langle \pi_{\beta, \alpha} : \alpha \leq \beta < \gamma \rangle \right\rangle.$$

Remark 2.11 ensures that $S_{\delta'}$ satisfies (1) and (2)³. Finally assume that $\delta' = \delta + 1$ and that S_δ satisfies (1) and (2). Note $\delta < \omega_2$. In the next paragraph we build $S_{\delta+1}$.

First partition $\omega_1 = T_0 \cup T_1 \cup T_2 \cup T_3 \cup T_4$ into five disjoint sets so that $|T_i| = \omega_1$ ($i \in 5$). Next we enumerate certain sets. Let $\mathcal{P}(\omega) = \{Y_i : i \in T_0\}$. Let $V = \{f_i : i \in T_1\}$, where V is the set of all increasing functions in ω^ω . Let $T = \omega^\omega \times \omega^\omega \times \mathbb{P} \times \mathbb{P} \times \delta = \{t_i : i \in T_2\}$ be enumeration of T in such a way that every element occurs ω_1 many times on the list. Let $\Phi \times \delta = \{\langle \phi_i, \alpha_i \rangle : i \in T_3\}$, where Φ is the set of all continuous monotone maps $\phi : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$, and note that this enumeration is possible because $|\Phi| = \mathfrak{c}$. Let $\Gamma = \{s_i : i \in T_4\}$ be enumeration of Γ such that every element of Γ appears ω_1 many times, where Γ is the set of all $\langle X, \bar{d}, \bar{\pi}, \bar{b}, \bar{\psi} \rangle$ such that $X \in [\delta]^{\leq \aleph_0}$, $\bar{d} \in \mathbb{P}^\omega$, $\bar{\pi} \in (\omega^\omega)^X$, $\bar{b} \in \mathbb{P}^X$, $\bar{\psi} \in (\omega^\omega)^X$. Now we build a decreasing sequence $\langle q_i : i < \omega_1 \rangle$ in \mathbb{Q}^δ . Since $\mathbb{Q}^\delta \neq 0$ pick arbitrary $q_0 \in \mathbb{Q}^\delta$. Assume that for $i < \omega_1$ we already built $\langle q_j : j < i \rangle$. If i is limit then by Lemma 4.4 we choose q_i such that $q_i \leq q_j$ ($j < i$). If $i = i_0 + 1$ then we distinguish five cases. Suppose that $i_0 \in T_0$. Then $Y_{i_0} \in \mathcal{P}(\omega)$. By Lemma 4.6 we pick $q_i \leq q_{i_0}$ such that $\text{set}(c_{q_i}) \subset Y_{i_0}$ or $\text{set}(c_{q_i}) \subset \omega \setminus Y_{i_0}$. Suppose that $i_0 \in T_1$. Then $f_{i_0} \in \omega^\omega$ is a strictly increasing so by Lemma 4.12 there is $q_i \leq q_{i_0}$ such that $\forall n < \omega [\text{set}(c_{q_i})(n) \geq f_{i_0}(n)]$. Suppose that $i_0 \in T_2$. If $\langle \pi_{i_0}, \psi_{i_0}, b_{i_0} \rangle$ is a normal triple, $d_{i_0} \leq b_{i_0}$ and $\pi''_{i_0} \text{set}(d_{i_0}) \in \mathcal{U}_{\alpha_{i_0}}$ then by Lemma 4.13 pick $q_i \leq q_{i_0}$, $d_i^* \leq d_{i_0}$, $\pi_i^*, \psi_i^* \in \omega^\omega$ so that $\langle \pi_i^*, \psi_i^*, d_i^* \rangle$ is a normal triple, $\alpha_{i_0} \in X_{q_i}$, $(\pi_i^*)'' \text{set}(d_i^*) =^* \text{set}(c_{q_i})$ and $\forall k \in \text{set}(d_i^*) [\pi_{i_0}(k) = \pi_{q_i, \alpha_{i_0}}(\pi_i^*(k))]$. Otherwise let $q_i = q_{i_0}$. Suppose that $i_0 \in T_3$.

³We consider the sequence $\langle c_i^\alpha : \alpha < \gamma \wedge i < \mathfrak{c} \rangle$ as the function from $\gamma \times \mathfrak{c}$ into \mathbb{P} while the sequence $\langle \pi_{\beta, \alpha} : \alpha \leq \beta < \gamma \rangle$ is considered as the function from $\{\langle \alpha, \beta \rangle : \alpha \leq \beta < \gamma\}$ into ω^ω .

Then $\phi_{i_0} : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$ is monotone and continuous. If $\phi_{i_0}(A) \neq 0$ for every $A \in \mathcal{U}_{\alpha_{i_0}}$ then by Lemma 4.10 pick $q_i \leq q_{i_0}$ such that $\alpha_{i_0} \in X_{q_i}$ and $\phi(A) \not\subseteq \text{set}(c_{q_i})$ for every $A \in \mathcal{U}_{\alpha_{i_0}}$. Otherwise let $q_i = q_{i_0}$. Suppose that $i_0 \in T_4$. Then there are $X \in [\delta]^{\leq \aleph_0}$, $\bar{d} \in \mathbb{P}^\omega$, $\bar{b} \in \mathbb{P}^X$, and $\bar{\pi}, \bar{\psi} \in (\omega^\omega)^X$ such that $\mathfrak{s}_{i_0} = \langle X, \bar{d}, \bar{\pi}, \bar{b}, \bar{\psi} \rangle$. If $\text{cof}(\delta) = \omega$, $\gamma_{q_{i_0}} = \delta$, \bar{d} is a decreasing sequence in \mathbb{P} , $X \subset X_{q_{i_0}}$, $\text{sup}(X) = \delta$ and $\bar{d}, \bar{\pi}, \bar{\psi}$ and \bar{b} satisfy Lemma 4.11(1)–(3), then by Lemma 4.11 pick $q_i \leq q_{i_0}$, d_i^* and π_i^* satisfying Lemma 4.11(4)–(6). Otherwise let $q_i = q_{i_0}$. Now define $S_{\delta'}$ as follows: $\pi_{\delta, \delta} = \text{id}$, for $\alpha \leq \beta < \delta$ and $i < \omega_1$, c_i^α and $\pi_{\beta, \alpha}$ are as in S_δ , while for $\alpha < \delta$ and $i < \omega_1$, c_i^δ is c_{q_i} and $\pi_{\delta, \alpha}$ is $\pi_{q_j, \alpha}$ where j is minimal such that $\alpha \in X_{q_j}$.

Claim 5.2. *For every $\alpha < \delta$ there is $i < \omega_1$ such that $\alpha \in X_{q_i}$.*

Proof. Take $\alpha < \delta$. Consider the function $\phi : \mathcal{P}(\omega) \rightarrow \mathcal{P}(\omega)$ given by $\phi(A) = \omega$ for $A \subset \omega$. Clearly, $\phi(A) \neq 0$ for $A \in \mathcal{U}_\alpha$. So there is $i_0 \in T_3$ so that $\langle \phi, \alpha \rangle = \langle \phi_{i_0}, \alpha_{i_0} \rangle$. Then for $i = i_0 + 1$, by choice of q_i we have $\alpha_{i_0} \in X_{q_i}$. \square

Note that by Claim 5.2 $\pi_{\delta, \alpha}$ is defined for every $\alpha < \delta$. Namely, if $i < \omega_1$ is such that $\alpha \in X_{q_i}$, then $\pi_{\delta, \alpha} = \pi_{q_i, \alpha}$. We still have to prove that $S_{\delta'}$ is $\delta + 1$ -generic sequence. Only conditions (3) and (6) of Definition 2.10 need checking.

To see that (3) holds, take any $\alpha < \beta \leq \delta$. If $\beta < \delta$ the statement follows because S_δ is δ -generic and $S_\delta = S_{\delta+1} \upharpoonright \delta$. If $\beta = \delta$ let $\pi_1, \psi_1, d_1, b_1, \alpha$ be as in the statement of (3) and let $a \in \mathcal{U}_\delta$. Since $a \in \mathcal{U}_\delta$ there is $j < \mathfrak{c}$ such that $c_j^\delta = c_{q_j} \subset^* a$. Then $\langle \pi_1, \psi_1, d_1, b_1, \alpha \rangle = \mathfrak{t}_i$ for some $i \in T_2$ such that $i \geq j$. Note that this is true because every element of \mathbb{T} appears ω_1 many times in its enumeration. So by the choice of q_{i+1} we know that $c_{q_{i+1}} \subset^* c_{q_j} \subset^* a$ and that $d_{i+1}^*, \pi_{i+1}^*, \psi_{i+1}^*$ and b_{i+1}^* are such that $\langle \pi_{i+1}^*, \psi_{i+1}^*, b_{i+1}^* \rangle$ is a normal triple, $\alpha \in X_{q_{i+1}}$, $(\pi_{i+1}^*)'' \text{set}(d_{i+1}^*) = {}^* \text{set}(c_{q_{i+1}})$ and $\forall k \in \text{set}(d_{i+1}^*) [\pi_1(k) = \pi_{q_{i+1}, \alpha}(\pi_{i+1}^*(k))]$. Denote $b = (\pi_{i+1}^*)'' \text{set}(d_{i+1}^*)$ and note that $b \subset^* a$. Now $b, d_{i+1}^*, \pi_{i+1}^*, \psi_{i+1}^*$ and b_{i+1}^* witness that Definition 2.10(3) is true in this case also.

Next we show that $S_{\delta'}$ satisfies condition (6) of Definition 2.10. Let μ, X, \bar{d} , and $\bar{\pi}$ be as in Definition 2.10(6). Since we require i^* satisfying Definition 2.10(6) to be cofinal in \mathfrak{c} let $i_0 < \mathfrak{c}$ be fixed. If $\mu < \delta$, the condition is satisfied because $S_{\delta'} = S_\delta \upharpoonright \delta$ and S_δ is δ -generic. If $\mu = \delta$ then, by Claim 5.2 and because every element of Γ appears \mathfrak{c} many times in its enumeration, there is $i \in T_4$ such that $i \geq i_0$ and $\langle X, \bar{d}, \bar{\pi}, \bar{b}, \bar{\psi} \rangle = \mathfrak{s}_i$ and $X \subset X_{q_i}$. By the choice of q_{i+1} we know that d_{i+1}^*, π_{i+1}^* and $c_{i+1}^\delta = c_{q_{i+1}}$ satisfy conditions (4)–(6) of Lemma 4.11 which implies that they also satisfy condition (6) of Definition 2.10. Since $\pi_{\delta, \alpha} = \pi_{q_{i+1}, \alpha}$ for $\alpha \in X$ this shows that $i + 1 \geq i_0$ witnesses that Definition 2.10(6) is satisfied. \square

The proof of Theorem 5.1 shows something slightly more general. We could have started the construction with a fixed δ -generic sequence for some $\delta < \omega_2$, and then extended it to an ω_2 -generic sequence in the same way. So we have the following corollary to the proof.

Corollary 5.3 (CH). *Suppose that $\delta < \omega_2$ and that S_δ is a δ -generic sequence. Then there is an ω_2 -generic sequence S such that $S \upharpoonright \delta = S_\delta$.*

Thus, we have the following theorem.

Theorem 5.4 (CH). *There is a chain of P-points of length \mathfrak{c}^+ under both Rudin-Keisler and Tukey ordering.*

Proof. For each $\alpha < \omega_2$, let $\mathcal{U}_\alpha = \{a \in \mathcal{P}(\omega) : \exists i < \mathfrak{c} [\text{set}(c_i^\alpha) \subset^* a]\}$. As we have noted in Section 2, $\langle \mathcal{U}_\alpha : \alpha < \omega_2 \rangle$ is a sequence of P-points that is strictly increasing with respect to both \leq_{RK} and \leq_T . Thus the ordinal ω_2 embeds into the P-points under both orderings. \square

Raghavan and Verner introduce the concept of a weakly δ -generic sequence in [10], and they prove that their apparently weaker notion is equivalent to our notion of a δ -generic sequence in that paper. Thus Corollary 5.3, when combined with the results in [10], yields that for every $\delta < \omega_2$, any weakly δ -generic sequence can be extended to an ω_2 -generic sequence.

When CH is replaced with MA and the lemmas from Section 4 have been appropriately generalized, the proof of the natural generalization of Theorem 5.1 presents little difficulty. In the crucial successor step of the construction, ω_1 can be replaced everywhere with \mathfrak{c} ; all of the sets that need to be enumerated have size \mathfrak{c} because $\mathfrak{c}^{<\mathfrak{c}} = \mathfrak{c}$ under MA. The generalizations of the lemmas from Section 4 imply that each condition in \mathbb{Q}^δ has an extension that meets some given requirement, and the fact that \mathbb{Q}^δ is $< \mathfrak{c}$ closed allows us to find lower bounds at the limit steps. Therefore a \mathfrak{c}^+ -generic sequence exists under MA.

REFERENCES

- [1] A. Blass, *The Rudin-Keisler ordering of P-points*, Trans. Amer. Math. Soc. **179** (1973), 145–166.
- [2] N. Dobrinen and S. Todorćević, *Tukey types of ultrafilters*, Illinois J. Math. **55** (2013), no. 3, 907–951.
- [3] ———, *A new class of Ramsey-classification theorems and their application in the Tukey theory of ultrafilters, Part 1*, Trans. Amer. Math. Soc. **366** (2014), no. 3, 1659–1684.
- [4] ———, *A new class of Ramsey-classification theorems and their applications in the Tukey theory of ultrafilters, Part 2*, Trans. Amer. Math. Soc. **367** (2015), no. 7, 4627–4659.
- [5] C. Laflamme, *Forcing with filters and complete combinatorics*, Ann. Pure Appl. Logic **42** (1989), no. 2, 125–163.
- [6] C. Laflamme and J. Zhu, *The Rudin-Blass ordering of ultrafilters*, J. Symbolic Logic **63** (1998), no. 2, 584–592.
- [7] D. Milovich, *Tukey classes of ultrafilters on ω* , Topology Proc. **32** (2008), no. Spring, 351–362, Spring Topology and Dynamics Conference.
- [8] D. Raghavan and S. Shelah, *On embedding certain partial orders into the P-points under Rudin-Keisler and Tukey reducibility*, Trans. Amer. Math. Soc. **369** (2017), no. 6, 4433–4455.
- [9] D. Raghavan and S. Todorćević, *Cofinal types of ultrafilters*, Ann. Pure Appl. Logic **163** (2012), no. 3, 185–199.
- [10] D. Raghavan and J. L. Verner, *Chains of P-points*, Submitted.
- [11] N. I. Rosen, *P-points with countably many constellations*, Trans. Amer. Math. Soc. **290** (1985), no. 2, 585–596.
- [12] A. Rosłanowski and S. Shelah, *Norms on possibilities. I. Forcing with trees and creatures*, Mem. Amer. Math. Soc. **141** (1999), no. 671, xii+167.
- [13] S. Shelah, *Proper and improper forcing*, second ed., Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1998.
- [14] ———, *Diamonds*, Proc. Amer. Math. Soc. **138** (2010), no. 6, 2151–2161.
- [15] S. Shelah, C. Laflamme, and B. Hart, *Models with second order properties. V. A general principle*, Ann. Pure Appl. Logic **64** (1993), no. 2, 169–194.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076, AND
 MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11001 BELGRADE, SERBIA.

E-mail address: borisa@mi.sanu.ac.rs

URL: <http://www.mi.sanu.ac.rs/~borisa>

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076.

E-mail address: raghavan@math.nus.edu.sg

URL: <http://www.math.nus.edu.sg/~raghavan>