

SINGULAR SHOCK WAVES IN INTERACTIONS

MARKO NEDELJKOV

ABSTRACT. In a number of papers it was shown that there are one-dimensional systems such that they contain solutions with, so called, overcompressive singular shock waves besides the usual elementary waves (shock and rarefaction ones as well as contact discontinuities).

One can see their definition for a general 2×2 system with fluxes linear in one of dependent variables in [8]. This paper is devoted to examining their interactions with themselves and elementary waves. After a discussion of systems given in a general form, a complete analysis will be given for the ion-acoustic system given in [6].

1. INTRODUCTION

Consider the system

$$\begin{aligned} (f_2(u))_t + (f_3(u)v + f_4(u))_x &= 0 \\ (g_1(u)v + g_2(u))_t + (g_3(u)v + g_4(u))_x &= 0. \end{aligned} \tag{1}$$

where f_i, g_j , $i = 2, \dots, 4$, $j = 1, \dots, 4$ are polynomials with the maximal degree m , $(u, v) = (u(x, t), v(x, t))$ are unknown functions with a physical range Ω , $(x, t) \in \mathbb{R}_+^2 := \mathbb{R} \times \mathbb{R}_+$. We shall fix the following notation for the rest of the paper:

$$f_i(y) = \sum_{k=0}^m a_{i,k} y^k, \quad g_j(y) = \sum_{k=0}^m b_{j,k} y^k, \quad i = 2, 3, 4, \quad j = 1, 2, 3, 4.$$

There are cases when there is no classical solution to Riemann problem for the above system. Sometimes, there is a solution in the form of delta or singular shock wave. In [8] one can see when a system in evolution form (i.e. when $f_2 = u$, $g_1 = 1$ and $g_2 = 0$) permits a solution in the shape of singular shock wave. With the same type of reasoning and a more effort, one can give the answer to the same question in the case system (1).

The aim of this paper is to investigate an interaction of a singular shock wave with another wave. After a general statement about new initial data taken at interaction point (of course, true for delta shock waves, too) in Section 3, we shall present a detailed investigation in the case of the system (so called ion-acoustic system)

$$\begin{aligned} u_t + (u^2 - v)_x &= 0 \\ v_t + (u^3/3 - u)_x &= 0 \end{aligned} \tag{2}$$

given in [6].

1991 *Mathematics Subject Classification.* 35L65, 35L67.

Key words and phrases. conservation law systems, singular shock wave, interaction of singularities.

THE WORK IS SUPPORTED BY SERBIAN MINISTRY OF SCIENCE AND ECOLOGY

Definitions and concepts used here are similar to the ones in [8], based on the notion of approximate solution. They are described in the Section 2. Shortly, a solution to the above system is given by a net of smooth functions with equality substituted by a distributional limit.

Few interesting facts observed during the investigations of system (2) are arising a question about possibilities in a general case. Observed facts are:

- (1) The singular shock wave solution to a Riemann problem for (2) always has an increasing strength of the rate $\mathcal{O}(t)$, $t \rightarrow \infty$. (The strength of the shock is a function which multiplies the delta function contained in a solution, $s(t)$ in (10)). After the interaction, the resulting singular shock wave is supported by a curve, not necessary straight line as before, and its strength can be an increasing, but also a constant or a decreasing function with the respect to the time variable.
- (2) When the resulting singular shock wave has a decreasing strength (this can occurs during an interaction of a admissible singular shock wave with a rarefaction wave), after some time it can decompose into two shock waves. This is a quite new phenomenon.

The structure of this paper can be described in the following way.

In the second section we will introduce necessary notation and give basic notions based on the paper [8].

In the third section, one can find a way how to continue a solution to the general case of system (1) after an interaction point (Theorem 1). The basic assumption is that a left-hand side of the first, and the right-hand side of the second wave can be connected by a new singular shock wave. The conditions for such a possibility are formulated trough a notion of *second delta singular locus*, see Definition 5. Explicit calculations for a geometric description of the locus are possible to some extend in the case of system (1), but we shall omit them completely.

The results given in these sections are used in the next ones devoted to special case (2) and that is the main part of the paper.

The first part of the fourth section is devoted to description of a situation which can occur after a singular shock and a shock wave interact. In the same way one can do the same for two singular shock waves, as one can at the end of this section.

The final, 5th section, contains the most interesting and important results about singular shock and rarefaction wave interaction. In that case the decoupling of a singular shock into a pair of shock waves, already mentioned before, can occur. The analysis is done when a singular shock wave is on the left-hand side of a rarefaction wave. But one can easily see that these results can be obtained using the same procedure when a singular shock is on the other side of a rarefaction wave.

We hope that the present paper give some opportunity for extending the procedure in this paper for arbitrary initial data when a system posses singular or delta shock wave as a solution.

2. NOTATION

For a net of smooth functions, $U_\varepsilon : \mathbb{R}_+^2 \rightarrow \Omega$, $\varepsilon \in (0, 1)$ is *distributionally zero*, $U_\varepsilon \approx 0$, if $\iint U_\varepsilon \phi dxdt \rightarrow 0$, as $\varepsilon \rightarrow 0$ for every $\phi \in \mathcal{C}_0^\infty(\mathbb{R}_+^2)$. A maximal positive real α such that $\iint U_\varepsilon \phi dxdt = \mathcal{O}(\varepsilon^\alpha)$, as $\varepsilon \rightarrow 0$ is called the *rate of convergence*. Due to the linearity, $U_\varepsilon \approx V_\varepsilon$ if $U_\varepsilon - V_\varepsilon \approx 0$.

We say that a smooth function net $(u_\varepsilon, v_\varepsilon)$ is an *approximate solution* to (1) if

$$\begin{aligned} \iint (f_2(u_\varepsilon)\phi_t + (f_3(u_\varepsilon)v_\varepsilon + f_4(u_\varepsilon))\phi_x) dxdt &\rightarrow 0 \\ \iint ((g_1(u_\varepsilon)v_\varepsilon + g_2(u_\varepsilon))\phi_t + (g_3(u_\varepsilon)v_\varepsilon + g_4(u_\varepsilon))\phi_x) dxdt &\rightarrow 0, \end{aligned}$$

as $\varepsilon \rightarrow 0$ for every $\phi \in C_0^\infty(\mathbb{R}_+^2)$. The above means

$$\begin{aligned} (f_2(u_\varepsilon))_t + (f_3(u_\varepsilon)v_\varepsilon + f_4(u_\varepsilon))_x &\approx 0 \\ (g_1(u_\varepsilon)v_\varepsilon + g_2(u_\varepsilon))_t + (g_3(u_\varepsilon)v_\varepsilon + g_4(u_\varepsilon))_x &\approx 0. \end{aligned}$$

A net G_ε is said to be of a *bounded type* if

$$\sup_{(x,t) \in \mathbb{R} \times (0,T)} |G_\varepsilon(x,t)| = \mathcal{O}(1) \text{ as } \varepsilon \rightarrow 0,$$

for every $T > 0$.

Before the definition of basic building blocks for a solution to Riemann problem for (1) we give some useful functions.

Definition 1. (a) A net G_ε is said to be a *step function* with value (y_0, y_1) if it is of bounded type and

$$G_\varepsilon(y) = \begin{cases} y_0, & y < -\varepsilon \\ y_1, & y > \varepsilon \end{cases}$$

Denote $[G] := y_1 - y_0$.

- (b) A net D_ε is said to be a *split delta function* (S δ -function, for short) with value (α_0, α_1) if $D_\varepsilon = \alpha_0 D_\varepsilon^- + \alpha_1 D_\varepsilon^+$, where $\alpha_0 + \alpha_1 = 1$ and

$$D_\varepsilon G_\varepsilon \approx (y_0 \alpha_0 + y_1 \alpha_1) \delta, \quad (3)$$

for every step function G_ε with value (y_0, y_1) .

- (c) Let m be an odd positive integer. A net d_ε is said to be *m'-singular delta function* (m'SD-function, for short) with value (β_0, β_1) if $d_\varepsilon = \beta_0 d_\varepsilon^- + \beta_1 d_\varepsilon^+$, $\beta_0^{m-1} + \beta_1^{m-1} = 1$, $(d_\varepsilon^\pm)^i \approx 0$, $i \in \{1, \dots, m-2, m\}$, $(d_\varepsilon^\pm)^{m-1} \approx \delta$, and

$$d_\varepsilon^{m-1} G_\varepsilon \approx (y_0 \beta_0^{m-1} + y_1 \beta_1^{m-1}) \delta, \quad (4)$$

for every step function G_ε with value (y_0, y_1) .

- (d) Let m be an odd positive integer. A net d_ε is said to be *m-singular delta function* (mSD-function, for short) with value (β_0, β_1) if $d_\varepsilon = \beta_0 d_\varepsilon^- + \beta_1 d_\varepsilon^+$, $\beta_0^m + \beta_1^m = 1$, $(d_\varepsilon^\pm)^i \approx 0$, $i \in \{1, \dots, m-1\}$, $(d_\varepsilon^\pm)^m \approx \delta$, and

$$d_\varepsilon^m G_\varepsilon \approx (y_0 \beta_0^m + y_1 \beta_1^m) \delta, \quad (5)$$

for every step function G_ε with value (y_0, y_1) .

In this paper we shall assume the *compatibility condition* $D_\varepsilon d_\varepsilon \approx 0$, where D_ε is an S δ - and d_ε is an mSD- or m'Sd-function.

Suppose that the initial data are given by

$$u|_{t=T} = \begin{cases} u_0, & x < X \\ u_1, & x > X \end{cases} \quad v|_{t=T} = \begin{cases} v_0, & x < X \\ v_1, & x > X. \end{cases} \quad (6)$$

According to our model, the initial data can be regularized to be

$$u|_{t=T} = G_\varepsilon \text{ and } v|_{t=T} = H_\varepsilon, \quad (7)$$

where G_ε and H_ε are step functions with values (u_0, u_1) and (v_0, v_1) , respectively.

The classical waves in the net interpretation are given in the following way. (One can look in [2] or [7] for the definition of the classical waves.)

Definition 2. *Approximate shock wave* solution to (1) with the initial data (6) is

$$\begin{aligned} u_\varepsilon(x, t) &= G_\varepsilon(x - X - c(t - T)) \\ v_\varepsilon(x, t) &= H_\varepsilon(x - X - c(t - T)), \end{aligned} \quad (8)$$

where c is a speed of the wave, G_ε and H_ε are given by (7). The wave is admissible if it satisfies Lax conditions (with $\lambda_2 > \lambda_1$)

$$\lambda_1(u_0, v_0) > c > \lambda_1(u_1, v_1) \text{ or } \lambda_2(u_0, v_0) > c > \lambda_2(u_1, v_1)$$

provided that the appropriate field is genuinely nonlinear.

An i -th *approximate rarefaction wave* solution to (1) with the initial data (6) is

$$(u_\varepsilon, v_\varepsilon)(x, t) = \begin{cases} (u_0, v_0), & \frac{x-X}{t-T} < \lambda_i(u_0, v_0) - \varepsilon \\ (R_{i,u}, R_{i,v})\left(\frac{x-X}{t-T}\right), & \lambda_i(u_0, v_0) + \varepsilon < \frac{x-X}{t-T} < \lambda_i(u_1, v_1) - \varepsilon \\ (u_1, v_1), & \frac{x-X}{t-T} > \lambda_i(u_1, v_1) + \varepsilon \end{cases} \quad (9)$$

where u_ε and v_ε are of the bounded type, while $(R_{i,u}, R_{i,v})$ are self-similar non-constant solution to (1) (non-constant parts of classical i -th rarefaction wave solution to the system).

One can prove immediately that (8) is an approximate solution to the given system if and only if

$$(u, v)(x, t) = \begin{cases} (u_0, v_0), & x - X < c(t - T) \\ (u_1, v_1), & x - X > c(t - T) \end{cases}$$

satisfy the usual Rankine-Hugoniot conditions for the system. Also, one can see directly that (9) is approximate solution, too.

There are no contact discontinuity used in this paper, but one can define its approximate version in the same way as approximate shock wave.

One will see later that we do not need rarefaction wave to be smooth function but only continuous one. Thus the above definition of approximate rarefaction wave can be replaced by the rarefaction wave itself.

Now, we define the non-standard type of the wave used in the present paper.

Definition 3. *Singular shock wave* (DSSW for short) is an approximate solution to (2) with the initial data (6) of the form

$$\begin{aligned} u_\varepsilon(x, t) &= G_\varepsilon((x - X) - c(t - T)) \\ &\quad + \tilde{s}(t)(\alpha_0 d_\varepsilon^-((x - X) - c(t - T)) + \alpha_1 d_\varepsilon^+((x - X) - c(t - T))) \\ v_\varepsilon(x, t) &= H_\varepsilon((x - X) - c(t - T)) \\ &\quad + s(t)(\beta_0 D_\varepsilon^-((x - X) - c(t - T)) + \beta_1 D_\varepsilon^+((x - X) - c(t - T))) \\ &\quad + \tilde{\tilde{s}}(t)(\gamma_0 d_\varepsilon^-((x - X) - c(t - T)) + \gamma_1 d_\varepsilon^+((x - X) - c(t - T))) \end{aligned} \quad (10)$$

where

- (i) $c \in \mathbb{R}$ is the speed of the wave,
- (ii) $s(t)$, $\tilde{s}(t)$ and $\tilde{\tilde{s}}(t)$ are smooth functions for $t \geq 0$, and equal zero at $t = T$.
- (iii) G_ε and H_ε are step functions with values (u_0, u_1) and (v_0, v_1) respectively,

- (iv) $d_\varepsilon^1 = \alpha_0 d_\varepsilon^- + \beta_1 d_\varepsilon^+$ and $d_\varepsilon^2 = \gamma_0 d_\varepsilon^- + \gamma_1 d_\varepsilon^+$ are m SD- or m' SD-functions,
- (v) $D_\varepsilon = \alpha_0 D_\varepsilon^- + \alpha_1 D_\varepsilon^+$ is an $S\delta$ -function compatible with d^i , $i = 1, 2$.

The *singular part* of the wave is

$$\left[\begin{array}{c} \tilde{s}(t)(\alpha_0 d_\varepsilon^- + \alpha_1 d_\varepsilon^+) \\ s(t)(\beta_0 D_\varepsilon^- + \beta_1 D_\varepsilon^+) + \tilde{s}(t)(\gamma_0 d_\varepsilon^- + \gamma_1 d_\varepsilon^+) \end{array} \right].$$

The wave is overcompressive if its speed is less or equal to the left- and greater or equal to the right-hand side characteristics i.e.

$$\lambda_2(u_0, v_0) > \lambda_1(u_0, v_0) \geq c \geq \lambda_2(u_1, v_1) > \lambda_1(u_1, v_1).$$

In solving of an interaction problem below, the form of (10) will be changed in the following way. The values on the left- and/or the right-hand side of a DSSW can be non-constant functions which would imply that the speed of the DSSW becomes non-constant, $c = c(t)$. If a DSSW is involved in the interaction, then $s(T)$ will equal a strength of incoming DSSWs, not zero as in the case of initial data (6).

Remark 1. (a) In [8] one can find special choice for $S\delta$ - and d is m SD- or m' SD-functions. For example D_ε^\pm are given by the representatives

$$D_\varepsilon^\pm(y) := \frac{1}{\varepsilon} \phi\left(\frac{y - (\pm 2\varepsilon)}{\varepsilon}\right),$$

where $\phi \in \mathcal{C}_0^\infty$ is non-negative and even, $\text{supp } \phi \subset [-1, 1]$. m SD- and m' SD-functions can be chosen in the same manner, with different shifts of their arguments and with $\phi^{1/m}$ or $\phi^{1/(m-1)}$ instead of ϕ .

(b) Compatibility condition for an $S\delta$ -function D and an m SD- or m' SD-function d is automatically fulfilled if

$$\text{supp } d_\varepsilon^+ \cap \text{supp } D_\varepsilon^+ = \text{supp } d_\varepsilon^- \cap \text{supp } D_\varepsilon^- = \emptyset$$

(c) Idea behind the above definition of products (3), (4) and (5) is the following. Starting point is that we know nothing about infinitesimal values of the initial data (carried on by step functions G_ε and H_ε above) around zero, but only that any such unmeasurable influence stops at the points $\pm\varepsilon$. The above mentioned definitions are made in order to get uniqueness of all products where step functions, $S\delta$ -, m SD- and m' SD-functions appear. With an additional information for G_ε and H_ε around zero, one can choose D_ε and d_ε much more freely. For example, in G_ε and H_ε are monotone functions (which is quite natural assumption), relation (3) can be substituted by

$$D_\varepsilon G_\varepsilon \approx \gamma \delta, \quad \gamma \text{ can be any real between } \min\{y_0, y_1\} \text{ and } \max\{y_0, y_1\}.$$

(d) The following result is proved in [6], Theorem 3. Suppose that a solution $(u_\varepsilon, v_\varepsilon)$ to Dafermos-DiPerna viscosity approximation of (2)

$$\begin{aligned} u_t + (u^2 - v)_x &= \varepsilon t u_{xx} \\ v_t + (u^3/3 - u)_x &= \varepsilon t v_{xx} \\ (u, v)|_{t=0} &= \begin{cases} (u_0, v_0), & x < 0 \\ (u_1, v_1), & x > 0 \end{cases} \end{aligned}$$

is unbounded with respect to ε and approaches a constant states as $\varepsilon \rightarrow 0$ except the line $x = ct$. Then (u_0, v_0) and (u_1, v_1) can be joined by an overcompressive DSSW.

That is a motivation to use overcompressiveness as an admissibility condition for DSSW. Note that the same condition is used for delta shock waves, too (see [5] or [10], for example).

(e) The DSSWs are much more peculiar than the delta shock waves. The later can be viewed as elements of certain measure spaces (see [5] or [9], for example) which inserted into a system solves it in an appropriate sense. A DSSW viewed as a net of smooth functions converge to weighted measure spaces defined in [6], but the obtained limits can not be inserted back into the system. That is the reason why we use only nets of smooth functions for DSSWs, shock waves and rarefaction ones.

Definition 4. The set of all points $(u_1, v_1) \in \Omega$ such that there exists an singular shock wave solution (called *corresponding DSSW*) to Cauchy problem (1,6) is called *delta singular locus*. We shall write $(u_1, v_1) \in \text{DSL}(u_0, v_0)$. If the corresponding DSSW is overcompressive, then it is called *overcompressive delta singular locus*. We shall write $(u_1, v_1) \in \text{DSL}^*(u_0, v_0)$.

In the sequel, the term “solution” will always denote the approximate one. Also, we shall drop the subscript ε in the approximate solution. One has to keep in mind that any function depends on ε .

3. THE NEW INITIAL DATA

Suppose that system (1) posses a DSSW solution for some initial data. Assume one of the following.

(i) If an m SD-function is contained in the above DSSW, then assume

$$\deg(g_1) < m - 1, \deg(g_2) < m, \deg(f_2) < m. \quad (11)$$

(ii) If an m' SD-function is contained in the above DSSW, then assume

$$\deg(g_1) < m - 2, \deg(g_2) < m - 1, \deg(f_2) < m - 1. \quad (12)$$

Take the new initial data

$$u|_{t=T} = \begin{cases} u_0, & x < X \\ u_1, & x > X \end{cases}, \quad v|_{t=T} = \begin{cases} v_0, & x < X \\ v_1, & x > X \end{cases} + \zeta \delta_{(X,T)}, \quad (13)$$

for system (1), where ζ is a non-zero real.

Definition 5. The set of all points $(u_1, v_1) \in \Omega$ such that there exists an DSSW solution (called *corresponding DSSW*) to Cauchy problem (1,13) for some ζ is called *second delta singular locus* of initial strength ζ for (u_0, v_0) . We shall write $(u_1, v_1) \in \text{SDSL}_\zeta(u_0, v_0)$. If the the corresponding DSSW is overcompressive, then it is called *overcompressive second delta singular locus*, and write $(u_1, v_1) \in \text{SDSL}_\zeta^*(u_0, v_0)$.

Before the main theorem, let us give a useful lemma.

Lemma 1. Suppose that $(u_1, v_1) \in \text{DSL}(u_0, v_0)$. Then $(u_1, v_1) \in \text{SDSL}_\zeta(u_0, v_0)$ if $\zeta > 0$.

If the corresponding DSSW contains m SD-function, and m is an odd number, then the statement holds true for every real ζ .

Additionally, β_i , $i = 1, 2$, from Definition 3 for the corresponding DSSW do not depend on ζ .

Proof. We shall give the proof for a DSSW containing m SD-function (10). The other case can be proved in the same way.

Inserting functions u and v from (10) into system (1) with initial data (13) and taking into account relations (11) or (12), one gets

$$\begin{aligned}
f_2(u) &\approx f_2(G) \\
g_1(u) &\approx g_1(G) \\
g_2(u) &\approx g_2(G) \\
f_3(u) &\approx f_3(G) + \tilde{s}(t)^{m-1}(u_1\alpha_0^{m-1}(d^-)^{m-1} + u_0\alpha_1^{m-1}(d^+)^{m-1})ma_{3,m} \\
&\quad + \tilde{s}(t)^m(\alpha_0^m(d^-)^m + \alpha_1^m(d^+)^m)a_{3,m} \\
&\quad + \tilde{s}(t)^{m-1}(\alpha_0^{m-1}(d^-)^{m-1} + \alpha_1^{m-1}(d^+)^{m-1})a_{3,m-1} \\
f_4(u) &\approx f_4(G) + \tilde{s}(t)^{m-1}(u_1\alpha_0^{m-1}(d^-)^{m-1} + u_0\alpha_1^{m-1}(d^+)^{m-1})ma_{4,m} \\
&\quad + \tilde{s}(t)^m(\alpha_0^m(d^-)^m + \alpha_1^m(d^+)^m)a_{4,m} \\
&\quad + \tilde{s}(t)^{m-1}(\alpha_0^{m-1}(d^-)^{m-1} + \alpha_1^{m-1}(d^+)^{m-1})a_{4,m-1} \\
g_3(u) &\approx g_3(G) + \tilde{s}(t)^{m-1}(u_1\alpha_0^{m-1}(d^-)^{m-1} + u_0\alpha_1^{m-1}(d^+)^{m-1})mb_{3,m} \\
&\quad + \tilde{s}(t)^m(\alpha_0^m(d^-)^m + \alpha_1^m(d^+)^m)b_{3,m} \\
&\quad + \tilde{s}(t)^{m-1}(\alpha_0^{m-1}(d^-)^{m-1} + \alpha_1^{m-1}(d^+)^{m-1})b_{3,m-1} \\
g_4(u) &\approx g_4(G) + \tilde{s}(t)^{m-1}(u_1\alpha_0^{m-1}(d^-)^{m-1} + u_0\alpha_1^{m-1}(d^+)^{m-1})mb_{4,m} \\
&\quad + \tilde{s}(t)^m(\alpha_0^m(d^-)^m + \alpha_1^m(d^+)^m)b_{4,m} \\
&\quad + \tilde{s}(t)^{m-1}(\alpha_0^{m-1}(d^-)^{m-1} + \alpha_1^{m-1}(d^+)^{m-1})b_{4,m-1}.
\end{aligned}$$

There are two possible cases. Either $\tilde{s} \not\equiv 0$ and $a_{3,m} = b_{3,m} = 0$ (i.e. $\deg(f_3) \leq m-1$ and $\deg(g_3) \leq m-1$), or $\tilde{s} \equiv 0$. In both the cases, the procedure which follows is the same, so take $\tilde{s} \not\equiv 0$ and $a_{3,m} = b_{3,m} = 0$. From the first equation of (1) one gets

$$\begin{aligned}
&(f_2(u))_t + (f_3(u)v + f_4(u))_x \\
&\approx -c([f_2(G)] + [f_3(G)H + f_4(G)])\delta \\
&\quad + \tilde{s}(t)^{m-1}\tilde{s}'(t)(u_1\alpha_0^{m-1}\gamma_0 + u_0\alpha_1^{m-1}\gamma_1)ma_{3,m-1}\delta' \\
&\quad + (f_3(u_0)\beta_0 + f_3(u_1)\beta_1)\delta' + \tilde{s}(t)^m a_{4,m}\delta' \approx 0.
\end{aligned}$$

One immediately gets the speed of DSSW,

$$c = \frac{[f_3(G)H + f_4(G)]}{[f_2(G)]},$$

and the relations

$$\kappa_1 s(t) = \tilde{s}(t)^{m-1}\tilde{s}'(t) \text{ and } \kappa_2 s(t) = \tilde{s}(t)^m,$$

for some reals κ_1 and κ_2 . Finally, one gets

$$\kappa_1(u_1\alpha_0^{m-1}\gamma_0 + u_0\alpha_1^{m-1}\gamma_1)ma_{3,m-1} + f_3(u_0)\beta_0 + f_3(u_1)\beta_1 + \kappa_2 a_{4,m} = 0. \quad (14)$$

Inserting all these relations into the second equation, one gets

$$\begin{aligned} & (g_1(u)v + g_2(u))_t + (g_3(u)v + g_4(u))_x \\ & \approx (-c[g_1(G)H + g_2(G)] + [g_3(G)H + g_4(G)] + s'(t)(g_1(u_0\beta_0 + g_1(u_1)\beta_1))\delta \\ & + s(t)(g_1(u_0)\beta_0 + g_1(u_1)\beta_1 + g_3(u_0)\beta_0 + g_3(u_1)\beta_1 \\ & + \kappa_1(u_1\alpha_0^{m-1}\gamma_0 + u_0\alpha_1^{m-1}\gamma_1)mb_{3,m-1} + \kappa_2b_{4,m})\delta' \approx 0. \end{aligned}$$

The function s must be a linear one, say $s'(t) = \sigma$, and the above functional equation gives the last two equations in \mathbb{R} ,

$$-c[g_1(G)H + g_2(G)] + [g_3(G)H + g_4(G)] + \sigma(g_1(u_0)\beta_0 + g_1(u_1)\beta_1) = 0 \quad (15)$$

and

$$\begin{aligned} & -c((g_1(u_0) + g_3(u_0)\beta_0 + (g_1(u_1) + g_3(u_1)\beta_1) \\ & + \kappa_1(u_1\alpha_0^{m-1}\gamma_0 + u_0\alpha_1^{m-1}\gamma_1)mb_{3,m-1} + \kappa_2b_{4,m}) = 0. \end{aligned} \quad (16)$$

In the above equations, only important fact about s is its derivative. Thus one can safely put $s(t) = \sigma t + \zeta$ and if the above system (14-16) has a solution, then $(u_1, v_1) \in \text{SDSL}_\zeta(u_0, v_0)$ provided that \tilde{s} and $\tilde{\tilde{s}}$ can be recovered. Using the fact that $(u_1, v_1) \in \text{DSL}(u_0, v_0)$, that is certainly the case when $\zeta > 0$. If m is an odd number, then $\tilde{s} = s(t)^{1/m}$ and $\tilde{\tilde{s}} = \tilde{s}$ are always determined.

The second part of the assertion, that β_i , $i = 1, 2$ are independent of ζ is obvious from the above. \square

Remark 2. From the proof of the lemma one can see that it is actually possible for ζ to take some negative values, i.e. it is enough that $\zeta \geq -s(T)$, where T is a time of interaction when new initial data are given.

The following assertion is crucial for the construction of weak solution (a solution in an associated sense) to (1) after an interaction: At an interaction point of a DSSW and some other wave one can consider the new initial value problem which contains delta function.

Suppose that the initial data are given by

$$u(x, 0) = \begin{cases} u_0, & x < a \\ u_1, & a < x < b \\ u_2, & x > b \end{cases} \text{ and } v(x, 0) = \begin{cases} v_0, & x < a \\ v_1, & a < x < b \\ v_2, & x > b \end{cases} \quad (17)$$

such that there exist a DSSW starting from the point $x = a$ and a shock wave (or another DSSW wave) starting from the point $x = b$, $a < b$. They can interact if $c_1 > c_2$, where c_i is the speed of the i -th wave, $i = 1, 2$. For the simplicity we shall assume that $b = 0$.

Let (X, T) be the interaction point of the overcompressive DSSW wave starting at the point $x = a$

$$\begin{aligned} u^1(x, t) &= G^1(x - c_1t - a) + \tilde{s}^1(t) \left(\alpha_0^1 d^-(x - c_1t - a) + \alpha_1^1 d^+(x - c_1t - a) \right) \\ v^1(x, t) &= H^1(x - c_1t - a) + s^1(t) \left(\beta_0^1 D^-(x - c_1t - a) + \beta_1^1 D^+(x - c_1t - a) \right) \\ &+ \tilde{s}^1(t) \left(\gamma_0^1 d^-(x - c_1t - a) + \gamma_1^1 d^+(x - c_1t - a) \right) \end{aligned} \quad (18)$$

and the admissible (singular) shock wave

$$\begin{aligned} u^2(x, t) = & G^2(x - c_2t) + \tilde{s}^2(t) \left(\alpha_0^2 d^-(x - c_2t) + \alpha_1^2 d^+(x - c_2t) \right) \\ v^2(x, t) = & H^2(x - c_2t) + s^2(t) \left(\beta_0^2 D^-(x - c_2t) + \beta_1^2 D^+(x - c_2t) \right) \\ & + \tilde{s}^2(t) \left(\gamma_0^2 d^-(x - c_2t - a) + \gamma_1^2 d^+(x - c_2t - a) \right) \end{aligned} \quad (19)$$

where G^1 , G^2 , H^1 and H^2 are the generalized step functions with values (u_0, u_1) , (u_1, u_2) , (v_0, v_1) and (v_1, v_2) , respectively. Also, $(\alpha_0^i)^{m_1} + (\alpha_1^i)^{m_1} = (\gamma_0^i)^{m_1} + (\gamma_1^i)^{m_1} = \beta_0^i + \beta_1^i = 1$, $i = 1, 2$. Here, $m_1 = m$ if singular part of DSSW is m SD-function and $m_1 = m - 1$ in the case of m' SD-function. If the second wave is a shock one, then one can put $s^2 \equiv \tilde{s}^2 \equiv 0$.

The speed of a DSSW (as well as for a shock wave) can be found using the first equation in (1) because of assumption (11) or (12). For example, in the case of the first DSSW (18) we have

$$\begin{aligned} (f_2(u))_t + (f_3(u)v + f_4(u))_x & \approx (f_2(G))_t + (f_3(G)H + f_4(G))_x + (\text{const } s^1(t)\delta)_x \\ & \approx (-c_1[f_2(G)] + [f_3(G)H + f_4(G)])\delta + \text{const } s^1(t)\delta' \approx 0, \end{aligned}$$

where the term $\text{const } s^1(t)$ is determined, but we shall not write the exact value since it is not needed for the assertion. Missing argument in the above expression is $x - c_1t - a$.

Let $\Gamma_1 = \{x = c_1t + a\}$ and $\Gamma_2 = \{x = c_2t\}$. Then $[\cdot]_{\Gamma_i}$ denotes the jump at the curve Γ_i , $i = 1, 2$. Thus, one can see that the speed of that DSSW has the same value as in the case of shock wave,

$$c_1 = \frac{[f_3(G)H + f_4(G)]_{\Gamma_1}}{[f_2(G)]_{\Gamma_1}}.$$

Also,

$$c_2 = \frac{[f_3(G)H + f_4(G)]_{\Gamma_2}}{[f_2(G)]_{\Gamma_2}}.$$

Finally, one can see that the waves given by (18) and (19) will interact at the point (X, T) if $a < 0$ and $c_1 > c_2$, where

$$\begin{aligned} T &= \frac{a[f_2(G)]_{\Gamma_1}[f_2(G)]_{\Gamma_2}}{[f_3(G)H + f_4(G)]_{\Gamma_2}[f_2(G)]_{\Gamma_1} - [f_3(G)H + f_4(G)]_{\Gamma_1}[f_2(G)]_{\Gamma_2}} \\ X &= \frac{a[f_3(G)H + f_4(G)]_{\Gamma_2}[f_2(G)]_{\Gamma_1}}{[f_3(G)H + f_4(G)]_{\Gamma_2}[f_2(G)]_{\Gamma_1} - [f_3(G)H + f_4(G)]_{\Gamma_1}[f_2(G)]_{\Gamma_2}}. \end{aligned}$$

Denote by $(\tilde{u}(x, t), \tilde{v}(x, t))$ a solution before interaction time $t = T$ consisting of waves (18, 19).

Remark 3. In the case of system (2) one can easily calculate speeds of the above shocks and coordinates of the interaction point. The speeds of DSSW and entropy shock wave are

$$c_1 = \frac{u_1^2 - v_1 - u_0^2 + v_0}{u_1 - u_0} \text{ and } c_2 = \frac{u_2^2 - v_2 - u_1^2 + v_1}{u_2 - u_1}.$$

If $c_1 > c_2$, then one gets

$$X = \frac{-ac_2}{c_2 - c_1} \text{ and } T = \frac{a}{c_2 - c_1}.$$

for the interaction point (X, T) .

Theorem 1. *Let system (1) be given. Suppose that $(u_2, v_2) \in \text{SDSL}_\zeta(u_0, v_0)$, $\zeta = (\zeta_1 + \zeta_2)/(g_1(u_0)\beta_0 + g_1(u_1)\beta_1)$, where the constants ζ_i , $i = 1, 2$, are defined by*

$$\begin{aligned} g_1(u^1)v^1 + g_2(u^1)|_{(t=T)} &\approx \zeta_1\delta_{(X,T)} \\ g_1(u^2)v^2 + g_2(u^2)|_{(t=T)} &\approx \zeta_2\delta_{(X,T)}. \end{aligned}$$

The corresponding DSSW, $(\hat{u}, \hat{v})(x, t)$ is given by

$$\begin{aligned} \hat{u}(x, t) &= G(x - X - c(t - T)) \\ &\quad + \tilde{s}(t) \left(\alpha_0 d^-(x - X - c(t - T)) + \alpha_1 d^+(x - X - c(t - T)) \right) \\ \hat{v}(x, t) &= H(x - X - c(t - T)) \\ &\quad + s(t) \left(\beta_0 D^-(x - X - c(t - T)) + \beta_1 D^+(x - X - c(t - T)) \right) \\ &\quad + \tilde{\tilde{s}}(t) \left(\gamma_0 d^-(x - X - c(t - T)) + \gamma_1 d^+(x - X - c(t - T)) \right) \end{aligned} \quad (20)$$

for $t > T$. By Lemma 1, β_0 and β_1 are determined independently on ζ , so the definition of DSSW makes sense.

Then there exist a solution to (1,17) in the association sense such that it equals $(\tilde{u}, \tilde{v})(x, t)$ for $t < T - \varepsilon$, and it equals $(\hat{u}, \hat{v})(x, t)$ for $t > T + \varepsilon$.

Proof. Take a constant t_0 such that singular parts of the waves $(u^1(x, t), v^1(x, t))$ and $(u^2(x, t), v^2(x, t))$ has disjoint supports (i.e. $c_1 t - a - c_2 t > 4\varepsilon$, for $t < T - t_0\varepsilon$, if one uses the construction of the $S\delta$, $m\text{SD}$ and $m'\text{SD}$ -functions defined above).

Let us denote

$$\begin{aligned} \Delta_\varepsilon &= \{(x, t) : |x - X| \leq t_0\varepsilon + \varepsilon, |t - T| \leq t_0\varepsilon + \varepsilon\}, \\ \tilde{\Delta}_\varepsilon &= \{(x, t) : |x - X| \leq t_0\varepsilon, |t - T| \leq t_0\varepsilon\}, \\ A_\varepsilon &= \{(x, t) : |x - X| \leq t_0\varepsilon + \varepsilon, t = T - t_0\varepsilon - \varepsilon\}, \\ B_\varepsilon &= \{(x, t) : x = X + t_0\varepsilon + \varepsilon, |t - T| \leq t_0\varepsilon + \varepsilon\}, \\ C_\varepsilon &= \{(x, t) : |x - X| \leq t_0\varepsilon + \varepsilon, t = T + t_0\varepsilon + \varepsilon\}, \\ D_\varepsilon &= \{(x, t) : x = X - t_0\varepsilon - \varepsilon, |t - T| \leq t_0\varepsilon + \varepsilon\}. \end{aligned}$$

Define a cut-off function $\xi_\varepsilon(x, t)$ which equals zero for $(x, t) \in \Delta_\varepsilon$ and 1 for $(x, t) \in \mathbb{R}_+^2 \setminus \tilde{\Delta}_\varepsilon$. Let

$$(u_{temp}, v_{temp})(x, t) = \begin{cases} (\tilde{u}(x, t), \tilde{v}(x, t)), & t < T \\ (\hat{u}(x, t), \hat{v}(x, t)), & t > T. \end{cases}$$

We shall prove that

$$u(x, t) = u_{temp}(x, t)\xi_\varepsilon(x, t), \text{ and } v(x, t) = v_{temp}(x, t)\xi_\varepsilon(x, t), \quad x \in \mathbb{R}, \quad t \geq 0 \quad (21)$$

are approximate solutions to (1).

Denote

$$\mathbf{F}(u, v) = \begin{bmatrix} f_2(u) \\ g_1(u)v + g_2(u) \end{bmatrix} \text{ and } \mathbf{G}(u, v) = \begin{bmatrix} f_3(u)v + f_4(u) \\ g_3(u)v + g_4(u) \end{bmatrix}.$$

We have

$$\begin{aligned}
& \iint_{\mathbb{R}_+^2} \mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x dx dt \\
&= \iint_{\tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x dx dt \\
&= \iint_{\mathbb{R}_+^2 \setminus \tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x dx dt,
\end{aligned}$$

for every test function $\Psi = \begin{bmatrix} \psi_1 \\ \psi_2 \end{bmatrix} \in \mathcal{C}_0^\infty(\mathbb{R}_+^2)$.

The measure of the set $\tilde{\Delta}_\varepsilon$ is $\mathcal{O}(\varepsilon^2)$, as $\varepsilon \rightarrow 0$, while

$$\|\mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x\|_{L^\infty(\mathbb{R}_+^2)} \leq \text{const } \varepsilon^{-1+1/m}$$

due to the assumptions in Definition 1. Thus,

$$\iint_{\tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x dx dt \sim \varepsilon^{1-1/m} \rightarrow 0, \text{ as } \varepsilon \rightarrow 0.$$

Using the divergence theorem for the second integral one gets

$$\begin{aligned}
& \iint_{\mathbb{R}_+^2 \setminus \tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi_t + \mathbf{G}(u, v) \Psi_x dx dt \\
&= \int_{\partial \tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi \nu_t + \int \mathbf{G}(u, v) \Psi \nu_x ds \\
&\quad - \iint_{\mathbb{R}_+^2 \setminus \tilde{\Delta}_\varepsilon} \mathbf{F}(u, v)_t \Psi + \mathbf{G}(u, v)_x \Psi dx dt.
\end{aligned}$$

The last integral in the above expression tend to zero as $\varepsilon \rightarrow 0$ since (u, v) solves (1) in $\mathbb{R}_+^2 \setminus \tilde{\Delta}_\varepsilon$ due to the construction. For the other one we have

$$\begin{aligned}
& \int_{\partial \tilde{\Delta}_\varepsilon} \mathbf{F}(u, v) \Psi \nu_t + \int \mathbf{G}(u, v) \Psi \nu_x ds \\
&= \int_{A_\varepsilon} \mathbf{F}(u, v) \Psi dx - \int_{C_\varepsilon} \mathbf{F}(u, v) \Psi dx + \int_{D_\varepsilon} \mathbf{G}(u, v) \Psi dt - \int_{B_\varepsilon} \mathbf{G}(u, v) \Psi dt.
\end{aligned}$$

Functions u and v are L^∞ -bounded uniformly in ε on the sides B_ε and D_ε . Since their lengths are $\mathcal{O}(\varepsilon)$, integrals over them tends to zero as $\varepsilon \rightarrow 0$.

Using the fact that $f_2(d) \approx 0$ one gets

$$\lim_{\varepsilon \rightarrow 0} F(\tilde{u}, \tilde{v})|_{t=T} = \begin{bmatrix} 0 \\ (\zeta_1 + \zeta_2) \delta_{(X, T)} \end{bmatrix},$$

as well as the construction of $S\delta$ - and $m'SD$ (or mSD)-functions, one gets

$$\lim_{\varepsilon \rightarrow 0} \int_{A_\varepsilon} \mathbf{F}(u, v) dx = \begin{bmatrix} 0 \\ \zeta_1 + \zeta_2 \end{bmatrix} \cdot \Psi(X, T).$$

Thus

$$\lim_{\varepsilon \rightarrow 0} \int_{C_\varepsilon} \mathbf{F}(u, v) dx = - \begin{bmatrix} 0 \\ \zeta_1 + \zeta_2 \end{bmatrix} \cdot \Psi(X, T)$$

has to be true. This implies $f_2(\hat{u})|_{(X, T)} \approx 0$ and

$$g_1(\hat{u})\hat{v} + g_2(\hat{u})|_{(X, T)} \approx (\zeta_1 + \zeta_2) \delta_{(X, T)}. \quad (22)$$

Due to conditions (11) or (12) one immediately gets $f_2(\hat{u})|_{(X,T)} \approx 0$. Put $\zeta = (\zeta_1 + \zeta_2)/(g_1(u_0)\beta_0 + g_1(u_1)\beta_1)$. Then

$$g_1(\hat{u})\hat{v} + g_2(\hat{u})|_{t=T} \approx \hat{G}\hat{H} + \hat{G} + s(T)(g_1(u_0)\beta_0 + g_1(u_1)\beta_1)\delta(X)$$

and after another restriction on the point $x = X$,

$$g_1(\hat{u})\hat{v} + g_2(\hat{u})|_{(X,T)} \approx (\zeta_1 + \zeta_2)\delta_{(X,T)}.$$

That concludes the proof. \square

Remark 4. The distributional limit of the result of the interaction is given by

$$u(x, t) = \begin{cases} u_0, & x < c_1 t - a, & t < t \\ u_1, & c_1 t - a < x < c_2 t, & t < T \\ u_2, & x > c_2 t, & t < T \\ u_0, & x < ct + X, & t > T \\ u_2, & x > ct + X, & t > T \end{cases}$$

$$v(x, t) = \begin{cases} v_0, & x < c_1 t - a, & t < t \\ v_1, & c_1 t - a < x < c_2 t, & t < T \\ v_2, & x > c_2 t, & t < T \\ v_0, & x < ct + X, & t > T \\ v_2, & x > ct + X, & t > T \end{cases} + s_1(t)\delta_{S_1} + s_2(t)\delta_{S_2} + s(t)\delta_S,$$

where $S_1 = \{(x, t) : x = c_1 t + a, t \in [0, T]\}$, $S_2 = \{(x, t) : x = c_2 t, t \in [0, T]\}$ and $S = \{(x, t) : x - X = c(t - T), t \in [T, \infty)\}$. If the second wave (19) is a shock one, then $s_2 \equiv 0$.

The above solution is continuous in t with values in $\mathcal{D}'(\mathbb{R})$. This fact can be used in the approach similar to [3] and [4], where the variable t is treated separately, i.e. when system (1) is considered to be in evolution form.

The theorem shows that after an interaction of a DSSW with some shock or another DSSW the problem reduces to solving system (1) with the new initial data (13).

Remark 5. (i) The solution to the interaction problem from Theorem 1 is always associated with a lower association rate than the solution of the original Riemann problem. For specific system it may be possible to make more sophisticated construction in order to improve the rate.

(ii) It appears that d^\pm are unavoidable correction factors even their distributional limit equals zero. The conditions (11) and (12) ensures that the new initial data at intersection point do not depend on m SD- or m' SD-functions in the solution. We have used them because the real nature of m SD- and m' SD-functions is not so clear yet.

The above theorem will be used in the rest of the paper for investigation of interactions between singular shock waves and other types of waves in the special case of system (2).

4. APPLICATIONS

Consider now system (2) which is a special case of (1). The authors of [6] defined and proved existence of DSSW solutions for some Riemann problems of this system. In the present paper, we will investigate interactions of such solutions with the other solutions to Riemann problem for (2). In order to familiarize a reader with the presented results, let us give some basic remarks about such solutions.

For a given Riemann data (u_0, v_0) , (v_1, v_1) , there are three basic solution types:

(a) *Shock waves*

$$u(x, t) = \begin{cases} u_0, & x < ct \\ u_1, & x > ct \end{cases} \quad v(x, t) = \begin{cases} v_0, & x < ct \\ v_1, & x > ct \end{cases} \quad (23)$$

where $c = [u^2 - v]/[u]$ and (u_1, v_1) lies in an admissible part of Hugoniot locus of the point (u_0, v_0) .

(b) *Centered rarefaction waves*

$$u(x, t) = \begin{cases} u_0, & x < (u_0 - 1)t \\ x/t + 1, & (u_0 - 1)t \leq x \leq (u_1 - 1)t \\ u_1, & x > (u_1 - 1)t \end{cases} \quad (24)$$

$$v(x, t) = \begin{cases} v_0, & x < (u_0 - 1)t \\ (x/t)^2/2 + 2x/t + C_1, & (u_0 - 1)t \leq x \leq (u_1 - 1)t \\ v_1, & x > (u_1 - 1)t \end{cases}$$

(1-rarefaction wave), where $C_1 = v_0 - u_0^2/2 - u_0 - 1/2$, and (u_1, v_1) lies in an 1-rarefaction curve starting at the point (u_0, v_0) . Or

$$u(x, t) = \begin{cases} u_0, & x < (u_0 + 1)t \\ x/t - 1, & (u_0 + 1)t \leq x \leq (u_1 + 1)t \\ u_1, & x > (u_1 + 1)t \end{cases} \quad (25)$$

$$v(x, t) = \begin{cases} v_0, & x < (u_0 + 1)t \\ (x/t)^2/2 - 2x/t + C_2, & (u_0 + 1)t \leq x \leq (u_1 + 1)t \\ v_1, & x > (u_1 + 1)t \end{cases}$$

(2-rarefaction wave), where $C_2 = v_0 - u_0^2/2 + u_0 - 1/2$, and (u_1, v_1) lies in an 2-rarefaction curve starting at the point (u_0, v_0) .

(c) *Singular shock waves* (see Definition 10) of 3'SD-type,

$$u(x, t) = \begin{cases} u_0, & x < ct \\ u_1, & x > ct \end{cases} + \tilde{s}(t)(\alpha_0 d_\varepsilon^-(x - ct) + \alpha_1 d_\varepsilon^+(x - ct)) \quad (26)$$

$$v(x, t) = \begin{cases} v_0, & x < ct \\ v_1, & x > ct \end{cases} + s(t)(\beta_0 D_\varepsilon^-(x - ct) + \beta_1 D_\varepsilon^+(x - ct)),$$

where $c = [u^2 - v]/[u]$, and all other terms are as in that definition. That means

$$D_\varepsilon \approx \delta, \quad (d_\varepsilon^\pm)^i \approx 0, \quad i = 1, 3, \quad (d_\varepsilon^\pm)^2 \approx \delta, \quad (27)$$

while (u_1, v_1) lies in a region denoted by Q_7 in [6] of the point (u_0, v_0) (see Figure 1).

For an arbitrary Riemann problem to (2) one can construct a solution by the means of these waves or their combinations ([6]).

While interactions of the first two types can be handled in a usual way, interactions involving DSSW are quite different and far more interesting, so they become a topic of this paper.

The procedure for the DSSW interactions can be also used for systems (1). But a complete after-interaction solution highly depends on a particular system. That is the reason why we treat system (2) only.

In order to simplify notation, we shall substitute the point (X, T) in (13) by $(0, 0)$ and then solve the Cauchy problem (2, 13).

There are no multiplication of v with u in system (2), so in the sequel it will be enough to take $D^- = D^+$, $\alpha_0(t) := \alpha_0 \tilde{s}(t)$, $\alpha_1(t) := \alpha_1 \tilde{s}(t)$ and $\beta(t) := s(t)$, i.e. to look for a solution of the form

$$\begin{aligned} u &= G(x - ct) + (\alpha_0(t)d^-(x - ct) + \alpha_1(t)d^+(x - ct)) \\ v &= H(x - ct) + \beta(t)D(x - ct), \end{aligned} \quad (28)$$

where G and H are generalized step functions, while d is 3'SD- and D is S δ -function and $c \in \mathbb{R}$.

Let us determine SDSL of (2) for some $(u_0, v_0) \in \mathbb{R}^2$.

Substitution of (28) into the first equation of the system gives

$$\begin{aligned} c &= \frac{u_1^2 - v_1 - u_0^2 + v_0}{u_1 - u_0} \\ \alpha_0^2(t) + \alpha_1^2(t) &= \beta(t), \end{aligned} \quad (29)$$

where c is the speed of the wave. After neglecting all terms converging to zero as $\varepsilon \rightarrow 0$, the second equation becomes

$$\begin{aligned} \partial_t H_\varepsilon(x - ct) + \beta'(t)\delta(x - ct) - c\beta(t)\delta'(x - ct) + \partial_x \left(\frac{1}{3}G_\varepsilon^3 - G \right) \\ + (u_1\alpha_0^2(t) + u_0\alpha_1^2(t))\delta'(x - ct) = 0. \end{aligned}$$

Thus, the following relations has to hold.

$$\beta'(t) = c(v_1 - v_0) - \left(\frac{1}{3}u_0^3 - u_0 - \frac{1}{3}u_1^3 + u_1 \right) =: k, \quad (30)$$

i.e.

$$\beta(t) = kt + \zeta, \text{ since } \beta(0) = \zeta$$

and

$$u_1\alpha_0^2(t) + u_0\alpha_1^2(t) = c\beta(t). \quad (31)$$

Like in [6] one can see that the overcompressiveness means

$$u_0 - 1 \geq c \geq u_1 + 1,$$

i.e., v_1 lies between the curves

$$\begin{aligned} D &= \{(u, v) : v = v_0 + u^2 + u - u_0u - u_0\} \\ E &= \{(u, v) : v = v_0 - u + u_0u - u_0^2 + u_0\}, \end{aligned}$$

and $u_0 - u_1 \geq 2$.

Denote by J_1 the union of the parts of admissible Hugoniot locus

$$S_1 = \left\{ (u, v) : v - v_0 = (u - u_0) \left(\frac{u_0 + u}{2} + \sqrt{1 - \frac{(u_0 - u)^2}{12}} \right) \right\},$$




FIGURE 1

and

$$S_2 = \{(u_1, v_1) : v - v_0 = (u - u_0) \left(\frac{u_0 + u}{2} - \sqrt{1 - \frac{(u_0 - u)^2}{12}} \right)\},$$

for $u \in [u_0 - \sqrt{12}, u_0 - 3]$. Note that S_i is not an i th shock curve but only a label.

The points between the curves D and E , and on the left-hand side of J_1 defines the area denoted by Q_7 in [6]. Here, this area is called delta singular locus.

One can easily check that system (29,31) has a solution if and only if $\beta(t) > 0$. Depending on k , defined in (30), there are three possibilities for a resulting wave:

- (i) If $k > 0$, then $\tilde{\beta}'(t) > 0$ and $(u_1, v_1) \in Q_7$. The resulting DSSW has the same properties as before, i.e. its strength increases with the time.
- (ii) If $k = 0$, then $\tilde{\beta} \equiv \text{const} = \zeta > 0$ and the corresponding part of a singular overcompressive locus is J_1 . The result of the interaction is a new kind of DSSW, its strength is a constant with respect to the time.
- (iii) If $k < 0$ (this means that the point (u_1, v_1) is on the left-hand side of J_1), then the resulting DSSW has much more differences from the usual one (with an increasing strength). Its initial strength equals ζ , $\beta(0) = \zeta > 0$, but linearly decreases in time. At some point T_0 the strength of the DSSW equals zero and the singular shock wave does not exist after that. In the rest of the paper we shall see some cases when this happens. The new initial data for time $t = T_0$ are the Riemann ones, and the solution after that time can be find in the usual way, by using the results in [6].

All the above facts are collected in the following theorem.

Theorem 2. *The $SDSL_\zeta$, $\zeta > 0$, for (2,13) is the area bounded by the curves D , E , $S_2 \setminus J_1$ and $S_1 \setminus J_1$. (The area Q_7 is a subset of this one, as known from Lemma 1.) The overcompressive $SDSL_\zeta$, $\zeta > 0$, is a part of the $SDSL$ bounded by the curves D and E such that $u_1 \leq u_0 - 2$.*

4.1. Interaction of a DSSW and an admissible shock wave. Suppose that a DSSW with a speed c_1 and a left- and right-hand values $U_0 = (u_0, v_0)$ and $U_1 = (u_1, v_1)$, respectively, interact with an admissible shock wave with a speed $c_2 < c_1$ having left-hand and right-hand values $U_1 = (u_1, v_1)$ and $U_2 = (u_2, v_2)$, respectively, at a point (X, T) .

Lemma 2. *If the above DSSW and shock wave are admissible, (u_2, v_2) lies between the lines D and E . Thus, the solution after the interaction is a single overcompressive DSSW.*

Proof. Since $u_0 \geq u_1 + 3$ and $u_1 > u_2$ (because of the admissibility conditions for singular and shock wave), we have $u_0 > u_2 + 3$. The point (u_2, v_2) lies on the curve S_1 or S_2 with the origin at the point (u_1, v_1) . Thus

$$v_2 = v_1 + (u_2 - u_1) \left(\frac{u_1 + u_2}{2} \pm \sqrt{1 - \frac{(u_1 - u_2)^2}{12}} \right).$$

The point (u_1, v_1) lies in the area denoted by Q_7 , thus below or at the curve D with the origin at (u_0, v_0) . Therefore

$$v_1 \leq v_0 + u_1^2 + u_1 - u_0 u_1 - u_0.$$

Let the point (u_0, v_0) be the origin. The point (u_2, v_2) will be below the curve D if

$$\begin{aligned} & v_0 + u_1^2 + u_1 - u_0 u_1 - u_0 + (u_2 - u_1) \left(\frac{u_1 + u_2}{2} \pm \sqrt{1 - \frac{(u_1 - u_2)^2}{12}} \right) \\ & \leq v_0 + u_2^2 + u_2 - u_0 u_2 - u_0. \end{aligned}$$

Non-positivity of $u_1 - u_2$ gives

$$\pm \sqrt{1 - \frac{(u_1 - u_2)^2}{12}} \leq \frac{1}{2}(u_0 - u_1) + \frac{1}{2}(u_0 - u_2) - 1.$$

The left-hand side of the above inequality is less than 2, while the right-hand side is greater than 2. Thus, the point (u_2, v_2) really lies below the curve D .

In the same way one can prove that the point (u_2, v_2) lies above the curve E . \square

Remark 6. In the same manner as above, one can prove that the situation is the same when DSSW and shock wave change sides. That is, when an admissible DSSW interacts with an admissible shock wave from the right-hand side, then the solution is again a single admissible DSSW.

4.2. Double DSSW wave interaction. Suppose that an admissible DSSW with a speed c_1 and left- and right-hand side values $U_0 = (u_0, v_0)$ and $U_1 = (u_1, v_1)$, respectively, interacts with another DSSW with a speed $c_2 < c_1$ and left-hand (right-hand) side values $U_1 = (u_1, v_1)$ ($U_2 = (u_2, v_2)$) at the point (X, T) . Since the conditions for the existence of DSSW include $u_0 - u_1 \geq 3$ and $u_1 - u_2 \geq 3$, then $u_0 - u_2 \geq 6$, i.e. the point (u_2, v_2) is on the left-hand side of the line $u = u_0 - \sqrt{12}$. Concerning the position of the point (u_2, v_2) in the plane of wave regions with the origin at (u_0, v_0) there are three possibilities:

- (i) The point (u_2, v_2) is between or at the curves D and E . The result of the interaction is a single DSSW (with increasing strength).
- (ii) The point (u_2, v_2) is above the curve D . The result of the interaction is an 1-rarefaction wave followed with a DSSW.
- (iii) The point (u_2, v_2) is below the curve E . The result of the interaction is a DSSW followed by a 2-rarefaction wave.

SDSL's always have increasing strength in these three cases.

5. INTERACTION OF A DSSW AND A RAREFACTION WAVE

The last possibility of DSSW interaction is with a rarefaction wave. That possibility is omitted from a considerations of the general system (1) due to a richness of possible behaviors. Nevertheless, the most of specific Riemann problems can be treated similarly as system (2) was here, at least up to some point.

For a given point (u_0, v_0) , the rarefaction curves are given by (see [6])

$$\begin{aligned} R_1 &= \left\{ (u, v) : v = v_0 - \frac{1}{2}u_0^2 + \frac{1}{2}u^2 + u - u_0 \right\}. \\ R_2 &= \left\{ (u, v) : v = v_0 - \frac{1}{2}u_0^2 + \frac{1}{2}u^2 - u + u_0 \right\} \end{aligned}$$

Suppose that a DSSW coming from the left-hand side with left- and right-hand side values $U_0 = (u_0, v_0)$ and $U_1 = (u_1, v_1)$, interacts with a rarefaction wave at some point (X, T) . If the rarefaction wave is approximated with a number of small amplitude (non-admissible) shock waves like in wave front tracking algorithm (see [1] for example), intuition given in Theorem 1, indicates that the first task should be to look at the DSSW and the interaction of DSSW and non-admissible shock wave. It is possible to extend Theorem 1 for such a case, providing that a non-admissible shock wave has amplitude small enough (of the rate ε^2 , say). Denote by (u_r, v_r) the end-point in a rarefaction curve. Let us note that the starting point of the curve (u_1, v_1) is in Q_7 .

In what follows, we shall abuse the notation and denote by $(u_1, v_1) \in Q_7$ the left-hand side and by (u_2, v_2) the right-hand side value of a part from the rarefaction curve. These values will represent constant states in an approximated non-admissible shock wave, too. If $(u_2, v_2) \in Q_7$, then the result of the interaction is a single DSSW, with the left-hand side value equals (u_0, v_0) . The speed depends on initial values as in (29). So, one can continue the procedure taking approximate points from the rarefaction curve as the right-hand values of the non-admissible shock wave until it reaches the border of Q_7 .

After looking at the above discrete model we are back in a real situation.

Let us denote by $(c(t), t)$, t belonging to some interval, a path of the resulting DSSW through Q_7 . It is possible to explicitly calculate the above path. For example if a DSSW interacts with a centered 1-rarefaction waves, substituting

$$\begin{aligned} u(x, t) &= \begin{cases} u_1, & x < c(t) \\ \phi_1(x/t), & x > c(t) \end{cases} + \alpha_0(t)d^-(x - c(t)) + \alpha_1(t)d^+(x - c(t)) \\ v(x, t) &= \begin{cases} v_1, & x < c(t) \\ \phi_2(x/t), & x > c(t) \end{cases} + \beta(t)D(x - c(t)) \end{aligned}$$

in system (2), one obtains

$$\begin{aligned}
\tilde{\alpha}_0^2(t) + \tilde{\alpha}_1^2(t) &= \tilde{\beta}(t) \\
c(t) &= \left(t(1 - 2(u_1 - v_0 + v_1 + u_0^2 - u_1^2)) \right. \\
&\quad \left. + T(1 - 2(u_0 - v_1 - u_0 u_1 + u_0^2 - u_1^2)) \right) / (2(u_0 - 1)) \\
\tilde{\beta}'(t) &= c'(t) \left(\frac{1}{2} \left(\frac{c(t)}{t} + 1 \right) + \left(\frac{c(t)}{t} + 1 \right) + v_1 - \frac{1}{2} u_1^2 - u_1 - v_0 \right) \\
&\quad - \left(\frac{1}{3} \left(\frac{c(t)}{t} + 1 \right)^3 - \left(\frac{c(t)}{t} + 1 \right) - \frac{1}{3} u_0^3 + u_0 \right),
\end{aligned}$$

where the initial data for β at the point $t = T$ is the initial strength of the DSSW $\beta(T)$. The above calculations means that a form of the resulting singular shock curve and its strength are uniquely determined through the area Q_7 . If $(u_r, v_r) \in Q_7$, then the analysis is finished. Suppose that this is not true. The main problem is to analyze situation when rarefaction curve intersects the boundary of Q_7 . Let us try to find out what is happening by using a discrete model.

Thus, the first real problem is to find a form of solution when the points from the rarefaction curve satisfy: $(u_1, v_1) \in Q_7$ and $(u_2, v_2) \notin Q_7$.

Denote by \tilde{D} and \tilde{G} the intersection points of the curve J_1 (or the line $u = u_0 - 3$) with the curves E and D , respectively (see Figure 2).

5.1. The first critical case. Denote by J the 1-rarefaction curve starting from the point \tilde{G} and by J_2 the 2-rarefaction curve starting from the point \tilde{D} . The region where (u_2, v_2) can lie consist of five subregions:

- (i) *The rarefaction curve which starts at (u_1, v_1) intersects the curve D out of point \tilde{G} .* The point (u_2, v_2) lies in the region above the curve D and left of the line $u = u_0 - 3$. The final result of the interaction is a 1-rarefaction wave (R_1) followed by a DSSW with increasing strength.
- (ii) *The rarefaction curve which starts at (u_1, v_1) intersects the curve E out of point \tilde{D} .* The point (u_2, v_2) lies in the region below the curve E and on the left-hand side of the line $u = u_0 - 3$. The result of the interaction is a DSSW with increasing strength followed by a 2-rarefaction wave (R_2).
- (iii) *The rarefaction curve which starts at (u_1, v_1) intersects the curve J_1 out of points \tilde{D} and \tilde{G} .* Since an amplitude of a non-admissible shock wave can be as small as necessary, one can assume that the point (u_2, v_2) lies in the second delta singular locus and the resulting DSSW has a negative strength. The strength-function $\tilde{\beta}(t) = \zeta + k(t - T_0)$ of the resulting DSSW is decreasing, so, there could exists a point $T_1 = T - \zeta/k$ such that $\tilde{\beta}(T_1) = \alpha_0 = \alpha_1 = 0$. Let $X_1 = cT_1 + (X - T)$, where c is the speed of the resulting DSSW (space coordinate of the point where strength reaches zero). Therefore, in the time $t = T_1$, we have to solve new Riemann problem

$$u|_{t=T_1} = \begin{cases} u_0, & x < X_1 \\ u_2, & x > X_1 \end{cases}, \quad v|_{t=T_1} = \begin{cases} v_0, & x < X_1 \\ v_2, & x > X_1 \end{cases}.$$

This problem has a unique entropy solution consists from two shock waves, since the point (u_2, v_2) is between the curves S_1 and S_2 , with respect to the origin at the point (u_0, v_0) . This means that the singular shock wave decouples into a pair of admissible shock waves. If the decoupling starts before u reaches the value

$\min\{u_r, u_0 - 2\}$, then the result of the interaction will be known after solving shock – rarefaction wave interactions. In any case there are no DSSWs in the solution after time $t = T_1$. If $u_r \leq u_0 - 2$ and the decoupling does not start before u reaches the value u_r , this pair of the shock waves are the final solution. The case when $u_r > u_0 - 2$ belongs to the following subsection, i.e. the second critical case.

(iv) *The rarefaction curve R_j , $j = 1$ or 2 , which starts at (u_1, v_1) intersects the curve J_1 in the point \tilde{G} .* We can take $\tilde{G} = (u_2, v_2)$ for convenience. The set of such points (u_1, v_1) lies on the inverse rarefaction curve (it starts from the right-hand side values),

$$\tilde{R}_1 = \{(u, v) : v = v_0 + (u_0^2 - u^2)/2 + u_0 - u\}$$

and

$$\tilde{R}_2 = \{(u, v) : v = v_0 + (u_0^2 - u^2)/2 - u_0 + u\}.$$

A straightforward calculation shows that this curve lies in the region Q_7 , thus this situation is possible, as one can see using the inverse rarefaction curves \tilde{R}_1 and \tilde{R}_2 given above.

If $j = 1$, then the point (u_2, v_2) belongs to J and the solution after the interaction is an R_1 -wave followed by a DSSW wave with a constant strength.

If $j = 2$, then the point (u_2, v_2) lies in the area below the curve J . This can be verified by direct calculation, taking into account that the amplitude of a non-admissible shock is small enough, $u_2 < u_0 - 2$. The solution after the interaction is an admissible DSSW with a decreasing strength. Further explanations of a such DSSW is given in the following subsection.

(v) *The rarefaction curve R_j which starts at (u_1, v_1) intersects the curve J_1 in the point \tilde{D} .* Again, let $\tilde{D} = (u_2, v_2)$. Simple calculation, as in the case (iv), shows that this situation is also possible since the inverse rarefaction curves starting from \tilde{D} stay in Q_7 . If $j = 2$, the point (u_2, v_2) belongs to J_2 , and then the solution after the interaction is DSSW with a constant strength followed by an R_2 -wave.

If $j = 1$, then use of the same arguments as above gives that the point (u_2, v_2) lies in the area above the curve J_2 and the result of the interaction is an admissible DSSW with a decreasing strength. Again, one can see the following subsection for the further analysis.

5.2. The second critical case. Now we are dealing with the problem when the rarefaction wave passes through the curves D or E , after passing through J_1 . One can see that this is the continuation of the cases (iii)-(v) from the previous part.

(a) Denote by \hat{D} the area above the curve D , below S_1 and on the left-hand side of the line $u = u_0 - 2$. Also denote by $\hat{\hat{D}}$ the area above the curve E , below S_1 and on the right-hand side of the line $u = u_0 - 2$. If (u_2, v_2) lies in one of these regions, the solution is combination of a rarefaction wave R_1 and an overcompressive DSSW with a decreasing or constant strength.

(b) Denote by \hat{E} the area below the curve E , above S_2 and on the left-hand side of the line $u = u_0 - 2$. Also denote by $\hat{\hat{E}}$ the area below the curve D , above S_2 and on the right-hand side of the line $u = u_0 - 2$. If (u_2, v_2) lies in one of these regions, the solution is then a combination of an overcompressive DSSW with a decreasing or constant strength and a rarefaction wave R_2 . Denote by D_0 the area bounded by the curves D , E , S_1 and S_2 such that $u < u_0 - 2$ in D_0 .




FIGURE 2

One can see that a rarefaction curve cannot enter into D_0 since it has to pass through the intersection point $(u_0 - 2, -2u_0 + v_0 + 2)$ of D and E , but

$$2u - 2u_0 - uu_0 + u^2/2 + u_0^2/2 + 2 > 0 \text{ (ie. } R_1 \text{ is above the curve } E)$$

$$2u_0 - 2u + uu_0 - u^2/2 - u_0^2/2 - 2 < 0 \text{ (ie. } R_2 \text{ is below the curve } D).$$

Therefore, a rarefaction curve which passes through the point $D \cap E$ goes either into \hat{D} or \hat{E} , and these cases are analyzed above.

Thus, we have described all important points of the interactions between DSSW and rarefaction waves. When a result of a single interaction is known, the question about further singular shock path could be answered by a successive use of the above procedures.

Remark 7. One can use the similar analysis of all possible cases when a rarefaction wave which interacts with a DSSW is on the left-hand side of it. Instead of direct rarefaction and singular shock curves, the inverse ones should be used, i.e. (u_2, v_2) is a starting point and one is able to calculate v_0 from formulas of E , D , S_1 , S_2 , R_1 and R_2 .

Remark 8. In the contrast with the case in [9], where interaction can generate some “strange” solution containing unbounded L^1_{loc} function, in the presented system one can find only bounded functions and DSSWs as a result on an interaction.

For a system (1) with $g_1 \neq \text{const}$, or $g_2 \neq 0$, interaction of DSSW and rarefaction waves cannot be treated as easy as here.

Thus, we have proved the following assertion for the interaction in the case of system (2).

Theorem 3. *Suppose that a DSSW interacts with a rarefaction wave at the time T . For some time period $T < t < T_1$ the solution is represented by a DSSW wave supported by a uniquely defined curve (not a line) followed by a new rarefaction wave. Depending on the right-hand value of the primary rarefaction wave, one has the following possible cases for a solution after $t > T_1$.*

- (a) *Single DSSW (supported by a line) with an increasing strength.*
- (b) *1-rarefaction wave followed by DSSW with an increasing strength.*
- (c) *Singular shock wave with an increasing strength followed by 2-rarefaction wave.*
- (d) *Singular shock wave with a decreasing strength prolonged by either a single DSSW with an increasing strength, or a pair of admissible shock waves.*
- (e) *1-rarefaction wave followed by DSSW with a constant strength.*
- (f) *Singular shock wave with a constant strength followed by 2-rarefaction wave.*
- (g) *Singular shock wave with a decreasing strength prolonged by either 1-rarefaction wave followed by DSSW with decreasing or constant strength, or DSSW with decreasing or constant strength followed by 2-rarefaction wave.*

“Prolonged” is the state after strength of DSSW becomes zero. Such wave can also stop with non-zero strength, and then there is obviously no prolongation described above.

REFERENCES

- [1] A. Bressan, Hyperbolic Systems of Conservation Laws, Oxford University Press, New York, 2000.
- [2] C. Dafermos, Hyperbolic Conservation Laws in Continuum Physics, Springer-Verlag, Heidelberg, 2000.
- [3] V. G. Danilov, V. M. Shelkovich, Dynamics of propagation and interaction of shock waves in conservation law systems, J. Differ. Equations, 211, 333-381 (2005).
- [4] V. G. Danilov, V. M. Shelkovich, Delta-shock wave type solution of hyperbolic systems of conservation laws, Q. Appl. Math., 29, 401-427 (2005).
- [5] F. Huang, Weak solution to pressureless type system, Comm. Partial Differential Equations **30**, no. 1-3, 283-304 (2005).
- [6] B. L. Keyfitz, H. C. Kranzer, Spaces of weighted measures for conservation laws with singular shock solutions, J. Differ. Equations **118**,2, 420-451 (1995).
- [7] P. D. Lax, Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves, SIAM, Philadelphia, 1973.
- [8] M. Nedeljkov, Delta and singular delta locus for one dimensional systems of conservation laws, Math. Method Appl. Sci. **27**, 931-955 (2004).
- [9] M. Nedeljkov, M. Oberguggenberger, Delta shock wave and interactions in a simple model case, Submitted.
- [10] Tan, D., Zhang, T. and Zheng, Y., Delta-shock waves as limits of vanishing viscosity for hyperbolic systems of conservation laws, J. Differ. Equations **112**, 1-32 (1994).

DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF NOVI SAD, TRG D. OBRADOVIĆA 4, 21000 NOVI SAD, YUGOSLAVIA

E-mail address: markonne@uns.ns.ac.yu, marko@im.ns.ac.yu