

Subsampled Inexact Newton methods for minimizing large sums of convex functions

Stefania Bellavia,* Nataša Krejić,† Nataša Krklec Jerinkić†

Abstract

This paper deals with the minimization of large sum of convex functions by Inexact Newton (IN) methods employing subsampled functions, gradients and Hessian approximations. The Conjugate Gradient method is used to compute the inexact Newton step and global convergence is enforced by a nonmonotone line search procedure. The aim is to obtain methods with affordable costs and fast convergence. Assuming strongly convex functions, R -linear convergence and worst-case iteration complexity of the procedure are investigated when functions and gradients are approximated with increasing accuracy. A set of rules for the forcing parameters and subsample Hessian sizes are derived that ensure local q -linear/superlinear convergence of the proposed method. The random choice of the Hessian subsample is also considered and convergence in the mean square, both for finite and infinite sums of functions, is proved. Finally, global convergence with asymptotic R -linear rate of IN methods is extended to the case of sum of convex function and strongly convex objective function. Numerical results on well known binary classification problems are also given. Adaptive strategies for selecting forcing terms and Hessian subsample size, streaming out of the theoretical analysis, are employed and the numerical results showed that they yield effective IN methods.

Key words: Inexact Newton, subsampled Hessian, superlinear convergence, global convergence, mean square convergence

1 Introduction

The problem we consider is

$$\min f_{\mathcal{N}}(x) = \frac{1}{N} \sum_{i=1}^N f_i(x) \quad (1)$$

*Department of Industrial Engineering, University of Florence, Viale Morgagni, 40/44, 50134 Florence, Italy, e-mail: stefania.bellavia@unifi.it. Member of the INdAM Research Group GNCS. Research supported by Gruppo Nazionale per il Calcolo Scientifico, (GNCS-INdAM) of Italy.

†Department of Mathematics and Informatics, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 4, 21000 Novi Sad, Serbia, e-mail: natasak@uns.ac.rs, natasa.krklec@dmi.uns.ac.rs. Research supported by Serbian Ministry of Education Science and Technological Development, grant no. 174030.

with $x \in \mathbb{R}^n$, N - very large and all functions $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ convex and $f_{\mathcal{N}}$ strongly convex. We are also interested in the case of large dimension n . There is a number of important problems of this type. To start with, one can be interested in minimizing the objective function stated in the form of mathematical expectation, $f(x) = E[F(x, w)]$, with w being a random variable from some probability space. Given that the analytical expression for mathematical expectation is rarely available, one possibility is to approximate the expectation with Sample Average Approximation, SAA, function. In that case, a sample $\{w^1, \dots, w^N\}$ is generated and the approximate objective function of the form (1) with $f_i(x) = F(x, w^i)$ is minimized. To ensure good approximation of the original objective function in general, one has to take a very large sample and thus calculating $f_{\mathcal{N}}(x)$, its gradient and Hessian is expensive.

Binary and multi-class classification problems, e.g., employing softmax activation function and cross-entropy loss can also be expressed in the form (1). For a given (very large) set of data, we are interested in classifying the data according to a set of rules specified by the data features.

In the framework of classical optimization, (1) is a convex problem that can be solved either by a first order or a second order method. However, the size of N makes classical approaches prohibitively costly and thus calls for specific methods. One possibility is to consider different subsampling schemes which are used to reduce the cost of calculating $f_{\mathcal{N}}$, its gradient and Hessian. There are many approaches in the literature, based on the idea of using a small sample subset at the beginning of iterative process and increasing the sample size as the solution is approached, ranging from simple heuristic approach [20, 9] to elaborate schemes that take into account the progress achieved up to the current iteration, [1, 2, 3, 6, 16, 25, 26, 27, 34, 36].

Whatever scheduling one adopts, the next question to be discussed is the choice of method. First order methods are attractive due to their low cost. One successful example is the stochastic gradient method, that employs a smaller subset of gradient components and thus reduce the cost even further, [20]. On the other hand several papers investigate the use of second order methods in this framework and demonstrate advantages in some important problems if the second order methods are correctly implemented, [4, 7, 5, 10, 11, 19, 37, 35, 40, 41]. For a comprehensive discussion of this issue one can see [8] and references cited therein.

In this paper we focus on subsampled Inexact Newton (IN) methods for (1) wrapped in a nonmonotone linesearch strategy. In Inexact Newton methods [15, 32], the Newton equation is approximately solved and in case of large scale problems an iterative Krylov method is used to compute an approximate solution of the Newton equation. The convexity of the objective function allows us to use CG method [24].

The choice of the nonmonotone strategy is motivated by the fact that the method uses approximate functions, at least initially, before the full sample is reached. Then, enforcing strict decrease in the Armijo rule might require unnecessary small steps. We adopt the nonmonotone line search procedure introduced in [29] and, assuming that each of the functions f_i is strongly convex, we prove R-linear convergence. Also the worst-case iteration complexity is investigated and it is proved that the worst-case complexity bound of this class of non-monotone algorithms, analyzed in [21], is maintained provided that errors in gradient and function also decay with R-rate. Namely, the method requires at most $O(\log(\epsilon^{-1}))$ iterations to reach $f(x^k) - f(x^*) < \epsilon$, where

x^* is the minimizer of (1).

Then, we turn our attention to the local properties of the method to obtain a local convergence rate faster than the R-linear convergence provided by first-order methods. The local convergence rate of IN methods with full Hessian depends on the choice of forcing terms which governs the error in solving each Newton linear system [17]. Here, as the Hessian of the objective function given in (1) might be prohibitively expensive to compute, we concentrate on subsampled Hessian and IN method that employs such Hessian approximations. We point out that in this context, it is pointless to solve the Newton equation exactly as the Hessian is generally approximated with a lower accuracy than function and gradient and the Newton model employed is actually a subsampled Newton model. Therefore, the use of CG method that allows us to control the accuracy in the solution of the Newton equation is advisable even if n is not large [9].

Assuming that the sample size scheduling is given for the objective function and the gradient, i.e. assuming that eventually one reaches the full sample size N , we analyze the local convergence of a subsampled Hessian IN method. The analysis provides bounds on the Hessian accuracy requirements that depend on the employed forcing terms. Adaptive forcing terms, streaming out of the iterative process itself are derived as well. Furthermore, it is shown that the local method combines well with the nonmonotone line search, i.e., the q -linear/ q -superlinear convergence rate of the local method are preserved.

In the second part of the paper we consider a randomized method obtained by relaxing the conditions for Hessian subsampling. Hence we prove the q -linear/superlinear convergence in the mean sense assuming that the Hessian approximation is good enough with high probability. The analysis yields relation between the Hessian subsample size, the forcing term and the (computable) sampled gradient at each iteration. Q -linear convergence in the mean sense is proved for fixed forcing terms with a fixed Hessian subsample size, while superlinear convergence in the mean square sense is obtained for the forcing terms that approach zero and increasing Hessian subsample sizes.

Having in mind the binary classification problem and the fact that the number of training points is enlarged over time in many applications, we also consider the case of unbounded N , i.e., the case where the objective function is defined as the mathematical expectation. For this problem we obtain convergence in the mean square sense as well.

Finally, the strong convexity assumption is relaxed similarly to the problem considered in [37]. A bound on the Hessian sample size, derived in [41] is used to obtain Hessians approximations which is positive definite with some high probability and CG is adapted to deal with possibly singular problems. The convergence in the the mean square is obtained for this problem as well.

From the performed theoretical analysis we derive adaptive rules for selecting both the forcing term sequence and the Hessian sample size. Particularly important feature of the proposed method is that the Hessian sample size is related to the current forcing term and approximated gradient norm, both quantities actually computable. Moreover, when q -superlinear convergence is sought, the Hessian sample size is adaptively chosen along the iterations, low accuracy and smaller Hessian sample size are generally used in the early stage of the method while the accuracy and the Hessian sample size increase in the last stage of the convergence. Finally, we note that the Hessian sample size is also allowed to decrease if too high accuracy is used at the previous iteration.

Numerical results on binary classification problems give numerical evidence of the proposed adaptive choices effectiveness.

The paper is organized as follows. In Section 2 the method is introduced and the global and local analysis is performed using the standard deterministic reasoning for the case of strongly convex functions f_i . Convergence in the mean square is proved in Section 3, considering all three cases - finite number of strongly convex functions f_i , an infinite number of strongly convex functions f_i 's and the last case with relaxed convexity assumptions. Some numerical results are presented in Section 4.

1.1 Related work

Our analysis is strictly related to that developed in [5, 7, 37], where convergence of Inexact subsampled Newton methods is investigated both in probability [37] and expectation [9, 7, 5]. We focus on the choice of the forcing terms, on the nonmonotone line search strategy and on adaptive choices of Hessian sample size.

The local behaviour of subsampled IN methods has been analysed in [7, 37]. Bounds to the accuracy required in the last stage of the procedure have been given, but the issue of developing an automatic transition between the initial stage of the procedure where a low accuracy in the Hessian approximation is enough, and the last stage stage where more accurate Hessian approximations are needed, is not investigated. However, in [37] the analysis is carried out under weaker assumptions than those we used here as the function $f_{\mathcal{N}}$ is supposed to be strongly convex only in a neighborhood of the sought minimizer without any assumptions on convexity of functions f_i . A set of conditions on the gradient and the Hessian sample sizes that ensures local R-superlinear convergence in the expectation under the assumption on the variance of the error norms (Bounded moment of Iterates) is given in [7]. The Hessian sample size is assumed to be increased at each iteration starting from a large enough initial sample size. In [9] an adaptive rule for choosing the gradient sample size is proposed along with an automatic criterion for the forcing term related to a variance estimation of the Hessian accuracy. The Hessian sample size is a fixed fraction of of the used gradient sample size.

Finally, we would like to mention that in [5] the authors perform a local complexity analysis of subsampled Inexact Newton methods and also show that methods that incorporate stochastic second-order information can be far more efficient on badly-scaled or ill-conditioned problems than first-order methods.

2 Inexact subsampled Newton method

We first introduce the notation and give some preliminary results. Throughout the paper $\mathcal{N}_k \subset \{1, 2, \dots, N\}$ denotes the sample used to approximate the objective function and its gradient, N_k denotes its cardinality and the subsampled function and gradient are defined as

$$f_{\mathcal{N}_k}(x) = \frac{1}{N_k} \sum_{i \in \mathcal{N}_k} f_i(x), \quad \nabla f_{\mathcal{N}_k} = \frac{1}{N_k} \sum_{i \in \mathcal{N}_k} \nabla f_i(x).$$

Moreover, $\mathcal{D}_k \subset \{1, 2, \dots, N\}$ is the sample used for Hessian approximation with cardinality D_k , and the subsampled Hessian is given by

$$\nabla^2 f_{\mathcal{D}_k}(x) = \frac{1}{D_k} \sum_{i \in \mathcal{D}_k} \nabla^2 f_i(x). \quad (2)$$

Here, we will consider subsampled Inexact Newton methods, that is iterative processes where at iteration k , given the current iterate x^k , the step s^k used to update the iterate satisfies

$$\nabla^2 f_{\mathcal{D}_k}(x^k)s^k = -\nabla f_{\mathcal{N}_k}(x^k) + r^k, \quad \|r^k\| \leq \eta_k \|\nabla f_{\mathcal{N}_k}(x^k)\|. \quad (3)$$

The term η_k belongs to $(0, 1)$ and it is called forcing term [15, 32]. Here and in the rest of paper $\|\cdot\|$ denotes the 2-norm.

Through the paper we will restrict our attention to convex functions, more precisely we will first consider strongly convex functions and then, in Subsection 3.2 relax the strong convexity to convexity. Let us state this formally.

Assumption A1 The functions f_i , $i = 1, \dots, N$ are twice continuously differentiable and strongly convex, i.e., for some $\lambda_n \geq \lambda_1 > 0$ there holds

$$\lambda_1 I \preceq \nabla^2 f_i(x) \preceq \lambda_n I, \quad \forall x \in \mathbb{R}^n \quad i = 1, \dots, N. \quad (4)$$

Assumption A1 implies a couple of inequalities that will be used further on. First of all, for all $x \in \mathbb{R}^n$ we have

$$\lambda_1 \|x - x^*\| \leq \|\nabla f_{\mathcal{N}}(x)\| \leq \lambda_n \|x - x^*\|, \quad (5)$$

where x^* is the unique minimizer of the function $f_{\mathcal{N}}$. Furthermore, according to [33, Theorem 2.10], for every $x \in \mathbb{R}^n$

$$\frac{\lambda_1}{2} \|x - x^*\|^2 \leq f_{\mathcal{N}}(x) - f_{\mathcal{N}}(x^*) \leq \frac{1}{\lambda_1} \|\nabla f_{\mathcal{N}}(x)\|^2. \quad (6)$$

Since $\nabla^2 f_{\mathcal{D}}(x)$ is positive definite we choose CG as the linear solver for computing s^k in (3). Thus, we assume that CG initialized with zero vector is employed at each Inexact Newton iteration. We will use the following technical Lemma from [18].

Lemma 2.1 [18] *Let $Ax = b$, where A is a symmetric and positive definite matrix. Furthermore, let us assume that CG is applied to this system and it is terminated at the i th iteration. Then if CG is initialized with the null vector the approximate solution x_i satisfies $x_i^T A x_i = x_i^T b$.*

Now, the above Lemma and Assumption A1 clearly imply the following result.

Lemma 2.2 *Assume that s^k satisfying (3) is obtained through CG method initialized with the null vector applied to the linear system*

$$\nabla^2 f_{\mathcal{D}_k}(x^k)s = -\nabla f_{\mathcal{N}_k}(x^k).$$

Then, $\|s^k\| \leq \lambda_1^{-1} \|\nabla f_{\mathcal{N}_k}(x^k)\|$.

2.1 Global convergence

In this section we analyze the behavior of the subsampled IN method and CG as the inner solver wrapped into the nonmonotone line-search strategy given in [29]. Iteration k of this procedure, denoted as Algorithm GIN, is sketched in Algorithm 2.1. The nonmonotone line-search has been chosen as in the first stage of the procedure subsampled functions and gradients are used and thus a strict Armijo type decrease might yield unnecessary small steps without real decrease in the original objective function. An additional freedom in the step size selection is therefore introduced adding to the Armijo condition a positive term ν_k usually denoted as error term [29]. The error sequence $\{\nu_k\}$ has to satisfy the following properties

$$\nu_k > 0, \quad \sum_k \nu_k < \infty. \quad (7)$$

Algorithm 2.1: k -th iteration of Algorithm GIN

Given $x^k \in \mathbb{R}^n$, $c \in (0, 1)$, $\bar{\eta} \in (0, 1)$ and $\{\nu_k\}$ such that (7) holds.

Step 1. Choose $\mathcal{N}_k, \mathcal{D}_k, \eta_k \in (0, \bar{\eta})$.

Step 2. Apply CG method initialized by the null vector to $\nabla^2 f_{\mathcal{D}_k}(x^k)s^k = -\nabla f_{\mathcal{N}_k}(x^k)$ and compute s^k satisfying (3).

Step 3. Find the smallest nonnegative integer j such that for $t_k = 2^{-j}$ there holds

$$f_{\mathcal{N}_k}(x^k + t_k s^k) \leq f_{\mathcal{N}_k}(x^k) + ct_k (s^k)^T \nabla f_{\mathcal{N}_k}(x^k) + \nu_k \quad (8)$$

and set $x^{k+1} = x^k + t_k s^k$, $k = k + 1$.

The Algorithm GIN is stated with an arbitrary scheduling sequences $\{\mathcal{N}_k\}$ and $\{\mathcal{D}_k\}$. The line search rule is defined with $f_{\mathcal{N}_k}$ implying that inexact function values, as well as approximated gradient values, are allowed in Algorithm GIN. Naturally, one expects to save computational effort while working with smaller samples, before reaching the full sample at some iteration. Complexity analysis and global convergence of this algorithm are presented. First, we give iteration complexity result for an arbitrary schedule for the gradient approximation considering the decrease in (possibly inexact) gradient. Then, we will prove R-linear convergence and show that the classical complexity result of $\mathcal{O}(\log(\epsilon^{-1}))$, [21] is obtained for a schedule that eventually ends up with the full sample.

As the search direction s^k is generated by the CG method and $f_{\mathcal{N}_k}$ is strictly convex, we know by Lemma 2.1 that s^k is descent direction for $f_{\mathcal{N}_k}$ at x^k and thus, under Assumption A1, the step size t_k is strictly positive even for the standard Armijo rule in Step 3. The lower bound for t_k is obtained in Lemma 2.4, similarly to [26].

Lemma 2.3 Let s^k be the step generated in Step 2 of Algorithm GIN. Then

$$-(\nabla f_{\mathcal{N}_k}(x^k))^T s^k \geq \frac{\lambda_1}{\lambda_n^2} (1 - \eta_k)^2 \|\nabla f_{\mathcal{N}_k}(x^k)\|^2.$$

Proof. The inexact condition implies

$$\|\nabla f_{\mathcal{N}_k}(x^k) - r^k\| \geq \|\nabla f_{\mathcal{N}_k}(x^k)\| - \|r^k\| \geq (1 - \eta_k) \|\nabla f_{\mathcal{N}_k}(x^k)\|.$$

On the other hand, we have

$$\|\nabla f_{\mathcal{N}_k}(x^k) - r^k\| = \|\nabla^2 f_{\mathcal{D}_k}(x^k) s^k\| \leq \|\nabla^2 f_{\mathcal{D}_k}(x^k)\| \|s^k\| \leq \lambda_n \|s^k\|$$

due to (4). Therefore

$$\|s^k\| \geq \frac{\|\nabla f_{\mathcal{N}_k}(x^k) - r^k\|}{\lambda_n} \geq \frac{1 - \eta_k}{\lambda_n} \|\nabla f_{\mathcal{N}_k}(x^k)\|. \quad (9)$$

Then, Lemma 2.1 and (4) yield

$$-(\nabla f_{\mathcal{N}_k}(x^k))^T s^k = (s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k \geq \lambda_1 \|s^k\|^2 \quad (10)$$

$$\geq \frac{\lambda_1}{\lambda_n^2} (1 - \eta_k)^2 \|\nabla f_{\mathcal{N}_k}(x^k)\|^2. \quad \square \quad (11)$$

Lemma 2.4 Suppose that the assumption A1 holds and let s^k be generated in Step 2 of algorithm GIN. Then (8) holds for $t_k \geq \bar{t} = (1 - c)\lambda_1/\lambda_n$.

Proof. Let k be an arbitrary iteration. If $t_k = 1$ satisfies (8), that is in Step 3 we have $j = 0$, then t_k is greater than \bar{t} , as $\bar{t} \in (0, 1)$. So let us consider the case $t_k < 1$. Then there exists $t'_k = 2t_k$ such that

$$\begin{aligned} f_{\mathcal{N}_k}(x^k + t'_k s^k) &> f_{\mathcal{N}_k}(x^k) + ct'_k (s^k)^T \nabla f_{\mathcal{N}_k}(x^k) + \nu_k \\ &\geq f_{\mathcal{N}_k}(x^k) + ct'_k (s^k)^T \nabla f_{\mathcal{N}_k}(x^k). \end{aligned}$$

On the other hand, Assumption A1 implies, using the standard arguments for functions with bounded Hessians,

$$\begin{aligned} f_{\mathcal{N}_k}(x^k + t'_k s^k) &= f_{\mathcal{N}_k}(x^k) + \int_0^1 (\nabla f_{\mathcal{N}_k}(x^k + yt'_k s^k))^T (t'_k s^k) dy \\ &\leq \frac{\lambda_n}{2} (t'_k)^2 \|s^k\|^2 + f_{\mathcal{N}_k}(x^k) + t'_k (\nabla f_{\mathcal{N}_k}(x^k))^T s^k. \end{aligned}$$

Combining the previous two inequalities we obtain

$$ct'_k (s^k)^T \nabla f_{\mathcal{N}_k}(x^k) \leq \frac{\lambda_n}{2} (t'_k)^2 \|s^k\|^2 + t'_k (\nabla f_{\mathcal{N}_k}(x^k))^T s^k.$$

Dividing by t'_k and using $t_k = t'_k/2$, by rearranging the previous inequality we get

$$t_k \geq \frac{-(1 - c)(\nabla f_{\mathcal{N}_k}(x^k))^T s^k}{\lambda_n \|s^k\|^2}. \quad (12)$$

Now, the result follows from (10) and $\min\{1, (1 - c)\lambda_1/\lambda_n\} = (1 - c)\lambda_1/\lambda_n$. \square

To prove the main results we need the following Lemma from [26].

Lemma 2.5 Assume that $\zeta_k \rightarrow 0$ R -linearly. Then, for every $\rho \in (0, 1)$ the sequence $\{a_k\}$ such that $a_k = \sum_{j=1}^k \rho^{j-1} \zeta_{k-j}$ converges to zero R -linearly.

Let us denote with ξ_k^g and ξ_k^f the inaccuracy in function and gradient,

$$\max_{x \in \{x^k, x^{k+1}\}} |f_{\mathcal{N}_k}(x) - f_{\mathcal{N}}(x)| \leq \xi_k^f, \quad \left| \|\nabla f_{\mathcal{N}_k}(x^k)\|^2 - \|\nabla f_{\mathcal{N}}(x^k)\|^2 \right| \leq \xi_k^g. \quad (13)$$

Following Grapiglia and Sachs [21], we now prove that despite inaccuracy in function and gradient, Algorithm GIN meets the complexity results of nonmonotone line search methods with exact function and gradients.

Theorem 2.1 Suppose that Assumption A1 holds and

$$\sum_{k=0}^{\infty} \xi_k^f < \infty. \quad (14)$$

Then, for a given $\epsilon \in (0, 1)$ Algorithm GIN takes at most

$$\bar{k} = \left\lceil \frac{f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^*) + \sum_{k=0}^{\infty} (2\xi_k^f + \nu_k)}{\kappa_c} \epsilon^{-2} \right\rceil,$$

iterations to ensure $\|\nabla f_{\mathcal{N}_{\bar{k}}}(x^{\bar{k}})\| \leq \epsilon$, where

$$\kappa_c = c(1-c) \frac{\lambda_1^2}{\lambda_n^3} (1-\bar{\eta})^2. \quad (15)$$

Proof. Note that, by Lemma 2.3 there follows

$$(\nabla f_{\mathcal{N}_k}(x^k))^T s^k \leq -\frac{\lambda_1}{\lambda_n^2} (1-\bar{\eta})^2 \|\nabla f_{\mathcal{N}_k}(x^k)\|^2. \quad (16)$$

Then, we can proceed as in the proof of Theorem 1 in [21]. Let \bar{k} be the first iteration such that $\|\nabla f_{\mathcal{N}_{\bar{k}}}(x^{\bar{k}})\| \leq \epsilon$. By (8) we obtain

$$\nu_k + f_{\mathcal{N}_k}(x^k) - f_{\mathcal{N}_k}(x^{k+1}) \geq -ct_k (\nabla f_{\mathcal{N}_k}(x^k))^T s^k, \quad k = 0, 1, \dots, \bar{k} - 1.$$

Moreover, by (16) and Lemma 2.4, there follows

$$-ct_k (\nabla f_{\mathcal{N}_k}(x^k))^T s^k \geq c(1-c) \frac{\lambda_1^2}{\lambda_n^3} (1-\bar{\eta})^2 \|\nabla f_{\mathcal{N}_k}(x^k)\|^2.$$

Then, for $k = 0, 1, \dots, \bar{k} - 1$, there holds

$$\nu_k + f_{\mathcal{N}_k}(x^k) - f_{\mathcal{N}_k}(x^{k+1}) \geq \kappa_c \epsilon^2$$

with κ_c given in (15). Therefore, by (13)

$$\nu_k + 2\xi_k^f + f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^{k+1}) \geq \kappa_c \epsilon^2 \quad k = 0, \dots, \bar{k} - 1.$$

Summing up for $k = 0, \dots, \bar{k} - 1$ we get

$$\sum_{k=0}^{\infty} (2\xi_k^f + \nu_k) + f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^{\bar{k}}) \geq \bar{k}\kappa_c\epsilon^2$$

and the thesis follows. \square

Note that in the previous theorem if $N_{\bar{k}} < N$ then

$$\|\nabla f_{\mathcal{N}}(x^k)\| \leq \epsilon_{\bar{k}} + \xi_k^g. \quad (17)$$

Strengthening the assumption on the accuracy in function and gradients we can prove R -linear convergence and complexity bounds which are far better than $O(\epsilon^{-2})$.

Theorem 2.2 *Assume that Assumption A1 holds and let $\{x^k\}$ be generated by Algorithm GIN. If the error sequences $\{\nu_k\}$, $\{\xi_k^f\}$ and $\{\xi_k^g\}$ converge to zero R -linearly then $\{x^k\}$ converges R -linearly to the solution of (1).*

Proof. Inequalities (8), (13) and (6) and Lemma 2.3 imply

$$\begin{aligned} f_{\mathcal{N}_k}(x^{k+1}) - f_{\mathcal{N}}(x^*) &\leq f_{\mathcal{N}_k}(x^k) - f_{\mathcal{N}}(x^*) + ct_k \nabla f_{\mathcal{N}_k}(x^k)^T s^k + \nu_k \\ &\leq f_{\mathcal{N}_k}(x^k) - f_{\mathcal{N}}(x^*) - ct_k \frac{\lambda_1}{\lambda_n^2} (1 - \eta_k)^2 \|\nabla f_{\mathcal{N}_k}(x^k)\|^2 + \nu_k \\ &\leq f_{\mathcal{N}_k}(x^k) - f_{\mathcal{N}}(x^*) \\ &\quad - ct_k \frac{\lambda_1}{\lambda_n^2} (1 - \eta_k)^2 (\lambda_1 (f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^*)) - \xi_k^g) + \nu_k \end{aligned}$$

Then, using Lemma 2.4 and (13) again, we obtain

$$f_{\mathcal{N}}(x^{k+1}) - f_{\mathcal{N}}(x^*) \leq \rho (f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^*)) + \bar{\xi}_k \quad (18)$$

where $\rho = 1 - ct \frac{\lambda_1^2}{\lambda_n^2} (1 - \bar{\eta})^2 \in (0, 1)$ and $\bar{\xi}_k = \nu_k + 2\xi_k^f + \xi_k^g$. Furthermore, we obtain

$$f_{\mathcal{N}}(x^{k+1}) - f_{\mathcal{N}}(x^*) \leq \rho^{k+1} (f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^*)) + \rho \sum_{j=1}^k \rho^{j-1} \bar{\xi}_{k-j} + \bar{\xi}_k. \quad (19)$$

Thus, Lemma 2.5 yields the statement. \square

Notice that R -linear convergence result obtained in Theorem 2.2 also holds for $\nu_k = 0$, that is, for the Armijo line search. The theorem above also allows us to prove the complexity result below.

Theorem 2.3 *Assume that Assumption A1 holds. If the error sequences $\{\nu_k\}$, $\{\xi_k^f\}$ and $\{\xi_k^g\}$ converge to zero R -linearly then for any $\epsilon \in (0, e^{-1})$, there exist $\hat{\rho} \in (0, 1)$ and $Q > 0$ such that Algorithm GIN takes at most*

$$\bar{k} = \lceil \frac{\log((f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^*) + Q))}{|\log(\hat{\rho})|} \rceil \log(\epsilon^{-1})$$

iterations to ensure $f_{\mathcal{N}}(x^{\bar{k}}) - f_{\mathcal{N}}(x^*) < \epsilon$.

Proof. Assumptions of Theorem 2.2 are satisfied and

$$f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^*) \leq \rho^k (f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^*)) + \sum_{j=0}^{k-1} \rho^j \bar{\xi}_{k-j}. \quad (20)$$

Given that the sequence $\{\sum_{j=0}^k \rho^j \bar{\xi}_{k-j}\}$ converges to zero R-linearly by Lemma 2.5, there exist $\bar{\rho} \in (0, 1)$ and $Q > 0$ such that

$$\sum_{j=0}^{k-1} \rho^j \bar{\xi}_{k-j} \leq Q \bar{\rho}^k.$$

Therefore, for $\hat{\rho} = \max\{\rho, \bar{\rho}\}$, we have from (20)

$$f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^*) \leq \hat{\rho}^k (f_{\mathcal{N}}(x^0) - f_{\mathcal{N}}(x^*) + Q)$$

and the statement follows as in Theorem 6 of [21]. \square

A couple of comment is due here. First of all, Theorems 2.1-2.3 deal with the possibility of infinite sequence of errors in the objective function and the gradient and thus provide a framework for considering unbounded N as well. However, our focus here is on the problems with fixed N . So, the statements of the above theorems apply to any kind of scheduling that ensures reaching the full sample for the objective function. Theorem 2.3 proves the complexity bound of $\log(\epsilon^{-1})$, although we work with cheaper objective function and the gradient whenever $N_k < N$ and thus provides theoretical justification for working with smaller samples. Moreover, the analysis carried out so far does not involve either the forcing term or the accuracy in Hessian approximation. Then, the results we provided hold even if $D_k = 1$ and only one CG-iteration is performed at each iteration of Algorithm GIN.

2.2 Local convergence

In this section we assume that the scheduling of the sample sizes \mathcal{N}_k is given and that eventually we reach $f_{\mathcal{N}}$ and $\nabla f_{\mathcal{N}}$ at some iteration \bar{k} and continue with $\mathcal{N}_k = \mathcal{N}$ for $k \geq \bar{k}$. Then, we may restrict all asymptotic theoretical considerations to the method defined with $\mathcal{N}_k = \mathcal{N}$, that is the step s^k satisfies

$$\nabla^2 f_{\mathcal{D}_k}(x^k) s^k = -\nabla f_{\mathcal{N}}(x^k) + r^k, \quad \|r^k\| \leq \eta_k \|\nabla f_{\mathcal{N}}(x^k)\|. \quad (21)$$

Following the analysis in [17], in this subsection we focus on the local q -linear and q -superlinear convergence of method GIN. Then, we assume that the generated sequence converges to the solution of (1), relying on the global convergence results proved in the previous subsection. The analysis presented here differs from that in [17] as we have to take into account the approximation in the Hessian and enlightens that the accuracy in the Hessian's approximation must be related to the adopted forcing term. This theoretical study enables us to devise an adaptive and computable rule for the Hessian sample size yielding q -linear and q -superlinear convergence. In fact, this is what we should obtain in order to justify the use of a second order method. The analysis is carried out under the following assumption on the subsampled Hessian matrices.

Assumption A2. There exists a constant L such that for any $\mathcal{D} \subseteq \{1, 2, \dots, N\}$ and any x sufficiently close to x^* we have

$$\|\nabla^2 f_{\mathcal{D}}(x) - \nabla^2 f_{\mathcal{D}}(x^*)\| \leq L\|x - x^*\|. \quad (22)$$

The above Assumption holds if there exists a neighborhood of x^* where each Hessian $\nabla^2 f_i$ is Lipschitz continuous. Note also that Assumption A1 implies that all function ∇f_i are Lipschitz continuous with the constant λ_n .

Moreover, we assume that the error in the Hessian approximation is determined only by the subsample size, independently of the sample taken and denote by $h(D, x)$ the norm of the error in the Hessian approximation for a given subsample size D at point x , i.e.

$$h(D, x) := \|\nabla^2 f_{\mathcal{N}}(x) - \nabla^2 f_{\mathcal{D}}(x)\|.$$

We make the following assumption on the subsample size D_k used at each iteration k .

Assumption A3. At each iteration k of Algorithm GIN the subsample size D_k is chosen such that

$$h(D_k, x^k) \leq C\eta_k, \quad (23)$$

with $0 < C < (1/\bar{\eta} - 1)\lambda_1/2$.

In the subsequent analysis for any $\delta > 0$ the ball with center x^* and radius δ will be denoted by $N_\delta(x^*)$, i.e. $N_\delta(x^*) = \{x \in \mathbb{R}^n : \|x - x^*\| \leq \delta\}$. Moreover, we let δ^* sufficiently small such that (22) holds for any x in $N_{\delta^*}(x^*)$ and $\delta^* < 2 \min\{\lambda_1, 1/\lambda_n\}/L$.

The next Theorem states that the full step is taken in Algorithm GIN eventually.

Theorem 2.4 *Assume that the sequence $\{x^k\}$ generated by Algorithm GIN converges to x^* . Let Assumptions A1 and A3 hold and $c \in (0, 1/4)$ in (8). Then, there exists k_0 such that for all $k \geq k_0$ the full step s^k is accepted in Step 3 of Algorithm GIN.*

Proof. Take $\bar{\varepsilon} \in (0, \bar{\eta}\lambda_1/2)$ and $\bar{\delta} > 0$ such that

$$\|\nabla^2 f_{\mathcal{N}}(x^k + \xi s^k) - \nabla^2 f_{\mathcal{N}}(x^k)\| \leq \bar{\varepsilon}, \text{ for all } \|s^k\| \leq \bar{\delta}, \xi \in (0, 1). \quad (24)$$

According to Lemma 2.2, $\lim_{k \rightarrow \infty} \|s^k\| = 0$. Then, there exists k_0 such that (24) holds for $k \geq k_0$. Moreover, the Taylor expansion yields

$$\begin{aligned} f_{\mathcal{N}}(x^k + s^k) &= f_{\mathcal{N}}(x^k) + (\nabla f_{\mathcal{N}}(x^k))^T s^k + \frac{1}{2}(s^k)^T \nabla^2 f_{\mathcal{N}}(\theta^k) s^k \\ &= f_{\mathcal{N}}(x^k) + (\nabla f_{\mathcal{N}}(x^k))^T s^k + \frac{1}{2}(s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k \\ &\quad + \frac{1}{2}(s^k)^T (\nabla^2 f_{\mathcal{N}}(\theta^k) \pm \nabla^2 f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{D}_k}(x^k)) s^k, \end{aligned} \quad (25)$$

with $\theta^k = x^k + \xi s^k$, $\xi \in (0, 1)$. From Assumption A3 we obtain

$$(s^k)^T (\nabla^2 f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{D}_k}(x^k)) s^k \leq h(D_k, x^k) \|s^k\|^2 \leq C\eta_k \|s^k\|^2 \leq C\bar{\eta} \|s^k\|^2. \quad (26)$$

Recall that Lemma 2.1 implies

$$(s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k = -\nabla f_{\mathcal{N}}(x^k)^T s^k. \quad (27)$$

Putting (24), (26) and (27) into (25) we get

$$f_{\mathcal{N}}(x^k + s^k) \leq f_{\mathcal{N}}(x^k) + \frac{1}{2}(\nabla f_{\mathcal{N}}(x^k))^T s^k + \frac{1}{2}(\varepsilon + C\bar{\eta})\|s^k\|^2. \quad (28)$$

From (10), we conclude that

$$f_{\mathcal{N}}(x^k + s^k) \leq f_{\mathcal{N}}(x^k) + \frac{1}{2}\left(1 - \frac{\varepsilon + C\bar{\eta}}{\lambda_1}\right)(\nabla f_{\mathcal{N}}(x^k))^T s^k.$$

Note that from $C < (1/\bar{\eta} - 1)\lambda_1/2$ there follows $\frac{C\bar{\eta}}{\lambda_1} < (1 - \bar{\eta})/2$. Therefore, the choice of ε yields $\varepsilon/\lambda_1 + C\bar{\eta}/\lambda_1 < 1/2$. Then,

$$f_{\mathcal{N}}(x^k + s^k) \leq f_{\mathcal{N}}(x^k) + \frac{1}{4}\nabla f_{\mathcal{N}}^T(x^k)s^k$$

and condition (8) is satisfied with $t_k = 1$ for any $k > k_0$ as $c \in (0, 1/4)$. \square

The following Lemma, whose proofs can be found in the Appendix, is needed in the subsequent convergence analysis.

Lemma 2.6 *Let Assumptions A1-A2 hold. If $x^k \in \mathcal{N}_{\delta^*}(x^*)$, $s^k \in \mathbb{R}^n$ and $\eta \in (0, 1)$ are such that $\|\nabla f_{\mathcal{N}}(x^k) + \nabla^2 f_{\mathcal{D}_k}(x^k)s^k\| \leq \eta\|\nabla f_{\mathcal{N}}(x^k)\|$ and $x^k + s^k \in \mathcal{N}_{\delta^*}(x^*)$ then*

$$\|\nabla f_{\mathcal{N}}(x^k + s^k)\| \leq (\eta + B(x^k))\|\nabla f_{\mathcal{N}}(x^k)\|,$$

with $B(x^k) = \frac{1}{\lambda_1}(\frac{1}{2\lambda_1}L\|\nabla f_{\mathcal{N}}(x^k)\| + h(D, x^k))$.

Lemma 2.7 *Let Assumption A1 holds and $\delta \in (0, \delta^*/(1 + \lambda_1^{-1}\lambda_n))$. If $x^k \in \mathcal{N}_{\delta}(x^*)$ and $\|\nabla f_{\mathcal{N}}(x^k) + \nabla^2 f_{\mathcal{D}_k}(x^k)s^k\| \leq \eta\|\nabla f_{\mathcal{N}}(x^k)\|$ then $x^k + s^k \in \mathcal{N}_{\delta^*}(x^*)$.*

The convergence of $\{x^k\}$ together with (5), implies the following result.

Theorem 2.5 *Assume that the sequence $\{x^k\}$ generated by Algorithm GIN converges to x^* . Let Assumptions A1-A3 hold and $c \in (0, 1/4)$ in (8). If $\eta_k = \bar{\eta}$ at each iteration of Algorithm GIN, then the sequence $\{x^k\}$ converges to x^* q -linearly for $\bar{\eta}$ small enough. Moreover, if $\lim_{k \rightarrow \infty} \eta_k = 0$, the convergence is q -superlinear.*

Proof. Note that, by the choice of C in Assumption A3, $\bar{\eta}(1 + \lambda_1^{-1}C) < (1 + \bar{\eta})/2$. Let $\delta \in (0, \delta^*/(1 + \lambda_1^{-1}\lambda_n))$. Take $\bar{\varepsilon} \in (0, \delta\lambda_1]$ sufficiently small such that $\bar{\eta}(1 + \lambda_1^{-1}C) + \lambda_1^{-2}L\bar{\varepsilon}/2 < \tau < 1$ for some $\tau \in (0, 1)$. Let k_0 be defined as in Theorem 2.4 and $\bar{k} \geq k_0$, such that $x^{\bar{k}} \in \mathcal{N}_{\delta}(x^*)$ sufficiently near to x^* to guarantee $\|\nabla f_{\mathcal{N}}(x^{\bar{k}})\| \leq \bar{\varepsilon}$.

Lemma 2.7 yields $x^{\bar{k}+1} \in \mathcal{N}_{\delta^*}(x^*)$ and by Lemma 2.6 we obtain

$$\|\nabla f_{\mathcal{N}}(x^{\bar{k}+1})\| \leq \tau\|\nabla f_{\mathcal{N}}(x^{\bar{k}})\| \leq \|\nabla f_{\mathcal{N}}(x^{\bar{k}})\| \leq \bar{\varepsilon}.$$

Therefore, using (5),

$$\|x^{\bar{k}+1} - x^*\| \leq \frac{1}{\lambda_1}\|\nabla f_{\mathcal{N}}(x^{\bar{k}+1})\| \leq \frac{1}{\lambda_1}\bar{\varepsilon} \leq \delta,$$

so $x^{\bar{k}+1} \in \mathcal{N}_\delta(x^*)$ and $\|\nabla f_{\mathcal{N}}(x^{\bar{k}+1})\| \leq \bar{\varepsilon}$.

As an inductive hypothesis suppose that for some $k > \bar{k}$ we have $x^k \in \mathcal{N}_\delta(x^*)$ and $\|\nabla f_{\mathcal{N}}(x^k)\| \leq \bar{\varepsilon}$. Then $x^{k+1} = x^k + s^k \in \mathcal{N}_{\delta^*}(x^*)$ by Lemma 2.7, and Lemma 2.6 implies

$$\begin{aligned} \|\nabla f_{\mathcal{N}}(x^{k+1})\| &\leq [(1 + \lambda_1^{-1}C)\eta_k + \lambda_1^{-2}L\varepsilon/2]\|\nabla f_{\mathcal{N}}(x^k)\| \\ &\leq \tau\|\nabla f_{\mathcal{N}}(x^k)\| \leq \|\nabla f_{\mathcal{N}}(x^k)\| \leq \varepsilon. \end{aligned}$$

Again, (5) yields $\|x^{k+1} - x^*\| \leq \delta$ and $x^{k+1} \in \mathcal{N}_\delta(x^*)$. Therefore, proceeding by induction we conclude that $x^k \in \mathcal{N}_{\delta^*}(x^*)$ for any $k \geq \bar{k}$ and by Lemma 2.6,

$$\|\nabla f_{\mathcal{N}}(x^{k+1})\| \leq [(1 + \lambda_1^{-1}C)\eta_k + \lambda_1^{-2}L\|\nabla f_{\mathcal{N}}(x^k)\|/2]\|\nabla f_{\mathcal{N}}(x^k)\|, \quad k \geq \bar{k}. \quad (29)$$

Therefore, as $\|\nabla f_{\mathcal{N}}(x^k)\| \rightarrow 0$, using (5) we obtain that $\{x^k\}$ converges to x^* with q -linear rate provided that

$$\bar{\eta} < \frac{\lambda_1}{\lambda_n} \frac{1}{1 + \lambda_1^{-1}C} < \frac{\lambda_1}{\lambda_n}.$$

Moreover, the q -superlinear convergence follows if $\lim_{k \rightarrow \infty} \eta_k = 0$. \square

The above results are in line with the classical convergence theory of Inexact Newton methods [15] as the local linear convergence require $\eta_k \leq \bar{\eta} < 1$ and the upper bound on $\bar{\eta}$ depends on inverse of the conditioning of the Hessian.

Let us now discuss one possible choice of η_k in order to obtain q -superlinear convergence of the procedure. Following ideas in [17] our choice of η_k depends on the agreement between the function and the subsampled Newton model. If there is a good agreement between these two quantities, even if the quality of the approximation in the Hessian is lower than that in the function, it is reasonable to use a small η in the subsequent iteration. Let us consider the following choice of η_k in Algorithm 2.1,

$$\eta_k = \min\{\bar{\eta}, \frac{|f_{\mathcal{N}_k}(x^k) - m_{k-1}(s^{k-1})|}{\|\nabla f_{\mathcal{N}_{k-1}}(x^{k-1})\|}\}, \quad \bar{\eta} < 1 \quad (30)$$

where

$$m_{k-1}(s) = f_{\mathcal{N}_{k-1}}(x^{k-1}) + \nabla f_{\mathcal{N}_{k-1}}(x^{k-1})^T s + \frac{1}{2} s^T \nabla^2 f_{\mathcal{D}_{k-1}}(x^{k-1}) s. \quad (31)$$

In the following theorem we show that the sequence $\{\eta_k\}$ generated by (30) converges to zero and ensures q -superlinear convergence provided that the full sample is used eventually.

Theorem 2.6 *Let assumptions in Theorem 2.5 hold and η_k given by (30). Then $\{x^k\}$ converges to x^* superlinearly.*

Proof. Let the iteration index $k > \tilde{k} + 1$, i.e. such that $\mathcal{N}_{k-1} = \mathcal{N}_k = \mathcal{N}$. Using the Taylors expansion and (23) we obtain

$$|f_{\mathcal{N}}(x^k) - m_{k-1}(s^{k-1})| \leq \frac{1}{2} \|s^{k-1}\|^2 \left(\frac{L}{2} \|s^{k-1}\| + C\eta_{k-1} \right). \quad (32)$$

Now, by Lemma 2.2 there follows

$$\eta_k \leq \frac{1}{2} \lambda_1^{-2} \|\nabla f_{\mathcal{N}}(x^{k-1})\|^2 \left[\lambda_1^{-1} \frac{L}{2} \|\nabla f_{\mathcal{N}}(x^{k-1})\| + C\bar{\eta} \right]. \quad (33)$$

Then, as $\lim_{k \rightarrow \infty} \|\nabla f_{\mathcal{N}}(x^k)\| = 0$ we have $\lim_{k \rightarrow \infty} \eta_k = 0$ and the superlinear convergence follows by Theorem 2.5. \square

The above result can be proved also choosing η_k as

$$\eta_k = \min\left\{\bar{\eta}, \frac{|f_{\mathcal{N}_k}(x^k) - m_{k-1}(s^{k-1})|}{\omega_k}\right\}, \quad \bar{\eta} < 1 \quad (34)$$

with

$$\omega_k = \mathcal{O}(\|\nabla f_{\mathcal{N}_{k-1}}(x^{k-1})\|).$$

We also underline that Theorem 2.6 in case of full Hessian, i.e. $\mathcal{N}_k = \mathcal{D}_k$, shows that the Inexact Newton method with the choice of forcing terms given by (34) is superlinearly convergent. Such result also follows from the analysis in [28, Theorem 3.10].

3 Mean square convergence

Large part of the previous analysis strongly relies on the Hessian error bound $h(D, x)$ which is not easily accessible. In this section, we are considering randomly chosen D_k and we ask for a good enough Hessian approximation with some probability smaller than 1. Thus, less conservative estimates are feasible. We use a bound similar to that derived in [41] to carry out the analysis and we obtain stochastic convergence results - convergence in a mean square sense (m.s.). The main result is that the q -linear convergence in m.s. can be achieved with a small enough but fixed forcing term η and with large enough but fixed Hessian sample size D . On the other hand, to achieve q -superlinear convergence in m.s. with η_k defined by (34), the Hessian sample size is required to increase as η_k goes to zero. This analysis paths the way to devise adaptive rules for selecting the Hessian sample size such that a small Hessian sample size is used in the early stage of the procedure, when η_k is close to one and linear systems are solved only to a low accuracy, and it is automatically increased when the solution is approached.

In this section we assume that the subsample \mathcal{D} is chosen randomly and uniformly - every $\nabla^2 f_i(x)$ has the same chance to be chosen. Let \mathcal{D} be any subset of \mathcal{N} such that $|\mathcal{D}| = D$. Then one can derive a bound on D such that, given $\gamma > 0$ and $\alpha \in (0, 1)$,

$$P(\|\nabla^2 f_{\mathcal{D}}(x) - \nabla^2 f_{\mathcal{N}}(x)\| \leq \gamma) \geq 1 - \alpha. \quad (35)$$

The corresponding bound is stated in Lemma 3.1 (see the proof in the Appendix), while a similar bound is provided in Lemma 4 of [41]. The result is obtained by using the Bernstein inequality, see [39] and [31] for further references.

Lemma 3.1 *Assume that A1 holds and that the subsample \mathcal{D} is chosen randomly and uniformly from \mathcal{N} . Let $\gamma > 0$ and $\alpha \in (0, 1)$ be given. Then (35) holds at any point x if the subsample size D satisfies*

$$D \geq \frac{2(\ln(2n/\alpha))(\lambda_n^2 + \lambda_n\gamma/3)}{\gamma^2} := \tilde{l}. \quad (36)$$

We use the above results and the analysis of the previous section, to design a globally convergent inexact subsampled Newton method with adaptive choice of the Hessian sample size. In particular we will choose D_k such that the inequality

$$h(D_k, x^k) \leq C \max\{\eta_k, \|\nabla f_{\mathcal{N}_k}(x^k)\|\} \quad (37)$$

holds with probability $1 - \alpha_k$, with $\alpha_k \in (0, 1)$ and $0 < C < (1/\bar{\eta} - 1)\lambda_1/2$. This corresponds to $\gamma = C \max\{\eta_k, \|\nabla f_{\mathcal{N}_k}(x^k)\|\}$ in (35).

3.1 Bounded sample - GIN-R method

In this subsection we are interested in the case of finite N . Let us assume that the full sample is eventually reached for the objective function and the gradient, i.e., $\mathcal{N}_k = \mathcal{N}$, for k sufficiently large, say for all $k \geq \bar{k}$. The procedure we obtain is based on GIN method but with a specific choice of the Hessian subsample \mathcal{D}_k , namely its cardinality is set according to (37) and the sample is randomly chosen. This procedure is denoted as GIN-R to emphasize the specific random choice of the Hessian subsample. We list its generic iteration k in Algorithm 3.1, where we denote the first steps as Step 1.a-1.c to make clear that they correspond to specific choices in Step 1 of Algorithm GIN.

We will analyze the convergence in the m.s. considering two possibilities. If the forcing terms converge to zero, then γ_k given by (38) converges to zero. The other case we consider is $\eta_k = \bar{\eta} < 1$. In this latter case Algorithm GIN-R yields γ_k bounded away from zero and we have $\mathcal{D}_k \subset \mathcal{N}$ with $D_k < N$ during the whole iterative process. We remark that Theorem 2.2 implies the R-linear convergence of the sequence generated by Algorithm GIN for any Hessian subsampling under the condition $\mathcal{N}_k = \mathcal{N}, k \geq \bar{k}$. The key issue in the global convergence analysis is that s^k is a descent search direction with an arbitrary good or poor Hessian approximation, i.e., regardless of the subsample used in GIN algorithm. The line search globalization strategy makes the algorithm R-linearly convergent for any choice of the subsample and the following statement holds.

Theorem 3.1 *Assume that A1 holds and let $\{x^k\}$ be generated by Algorithm GIN-R. If $\{\nu_k\}$ converges to zero R-linearly then $\{x^k\}$ converges R-linearly to x^* .*

Next we show another important intermediate result.

Algorithm 3.1: k -th iteration of Method GIN-R

Given $x^k \in \mathbb{R}^n$, $\bar{\eta} \in (0, 1)$, $c \in (0, 1)$, $C > 0$, ν_k , $\alpha_k \in (0, 1)$.

Step 1.a Choose $\mathcal{N}_k, \eta_k \in (0, \bar{\eta})$.

Step 1.b Compute

$$\gamma_k = C \max\{\eta_k, \|\nabla f_{\mathcal{N}_k}(x^k)\|\}. \quad (38)$$

Step 1.c Set $\alpha = \alpha_k$ and $\gamma = \gamma_k$ and compute D such that (36) holds. If $D \geq N_k$ set $\mathcal{D}_k = \mathcal{N}_k$. Else, choose the sample \mathcal{D}_k randomly and uniformly from \mathcal{N}_k such that $D_k \geq D$.

Step 2 Apply CG method initialized by the null vector to

$$\nabla^2 f_{\mathcal{D}_k}(x^k) s^k = -\nabla f_{\mathcal{N}_k}(x^k) \text{ and compute } s^k \text{ satisfying (3).}$$

Step 3. Find the smallest nonnegative integer j such that (8) holds for $t_k = 2^{-j}$ and set $x^{k+1} = x^k + t_k s^k$.

Lemma 3.2 *Suppose that the assumptions of Theorem 3.1 are satisfied and let $\{x^k\}$ be a sequence generated by Algorithm GIN-R. If η_k is defined by (30) then there exist positive constants B_1 and B_2 and $\tau \in (0, 1)$ such that*

$$\eta_k \leq B_1 \tau^k \quad \text{and} \quad \gamma_k \leq B_2 \tau^k$$

for all k large enough. If $\eta_k = \bar{\eta}$ then $\gamma_k = C\bar{\eta}$ for all k large enough.

Proof. First, Theorem 3.1 implies that the sequence of iterates x^k converges to the unique solutions R-linearly - for all k large enough we have

$$\|x^k - x^*\| \leq B\tau^k \quad (39)$$

where $B > 0$ and $\tau \in (0, 1)$. Now, using the Taylor's expansion, Lemma 2.2 and (4) we obtain for some $\theta_k \in [x^{k-1}, x^k]$

$$\begin{aligned} & |f_{\mathcal{N}}(x^k) - m_{k-1}(s^{k-1})| \\ &= |f_{\mathcal{N}}(x^{k-1}) + t_k \nabla f_{\mathcal{N}}(x^{k-1})^T s^{k-1} + \frac{1}{2} t_k^2 (s^{k-1})^T \nabla^2 f_{\mathcal{N}}(\theta_k) s^{k-1} \\ &\quad - f_{\mathcal{N}}(x^{k-1}) - \nabla f_{\mathcal{N}}(x^{k-1})^T s^{k-1} - \frac{1}{2} (s^{k-1})^T \nabla^2 f_{\mathcal{D}_{k-1}}(x^{k-1}) s^{k-1}| \\ &\leq \frac{1-t_k}{\lambda_1} \|\nabla f_{\mathcal{N}}(x^{k-1})\|^2 + \lambda_n \|s^{k-1}\|^2 \leq \frac{1}{\lambda_1} \|\nabla f_{\mathcal{N}}(x^{k-1})\|^2 + \frac{\lambda_n}{\lambda_1^2} \|\nabla f_{\mathcal{N}}(x^{k-1})\|^2. \end{aligned}$$

Therefore, using (5), for η_k given by (30) we obtain

$$\eta_k \leq \frac{|f_{\mathcal{N}}(x^k) - m_{k-1}(s^{k-1})|}{\|\nabla f_{\mathcal{N}}(x^{k-1})\|} \leq \left(\frac{1}{\lambda_1} + \frac{\lambda_n}{\lambda_1^2}\right) \lambda_n \|x^{k-1} - x^*\| \leq B_1 \tau^k \quad (40)$$

where $B_1 = (\frac{1}{\lambda_1} + \frac{\lambda_n}{\lambda_1^2})\lambda_n B/\tau$. Moreover, for γ_k given by (38) we get

$$\gamma_k \leq C \max\{B_1\tau^k, \lambda_n B\tau^k\} = C \max\{B_1, \lambda_n B\}\tau^k := B_2\tau^k.$$

Considering the case with $\eta_k = \bar{\eta}$, since the gradient converges to zero, for all k large enough we have $\|\nabla f_{\mathcal{N}}(x^k)\| < \bar{\eta}$ and therefore $\gamma_k = C\bar{\eta}$. \square

Now, we are ready to show the main result of this subsection.

Theorem 3.2 *Suppose that the assumptions of Theorem 3.1 and Assumption A2 are satisfied. Moreover, assume $c \in (0, 1/4)$ in (8) and $0 < C < (1/\bar{\eta} - 1)\lambda_1/2$ in (37). Let $\{x^k\}$ be a sequence generated by Algorithm GIN-R. Then, there are positive constants V_1, V_2, C_1, C_2 and $\tau \in (0, 1)$ such that for all k sufficiently large*

a) if η_k is defined by (30) then

$$E(\|x^{k+1} - x^*\|^2) \leq (V_1\tau^{2k} + V_2\alpha_k) E(\|x^k - x^*\|^2);$$

b) if $\eta_k = \bar{\eta}$ is sufficiently small then

$$E(\|x^{k+1} - x^*\|^2) \leq ((C_1\tau^k + C_2\bar{\eta})^2 + V_2\alpha_k) E(\|x^k - x^*\|^2).$$

Proof. Since the assumptions of Theorem 3.1 are satisfied, the sequence $\{x^k\}$ converges to the solution R-linearly, independently of D_k and for $\eta_k \in (0, 1)$. Thus, there exist constants $B > 0$ and $\tau \in (0, 1)$ such that $\|x^k - x^*\| \leq B\tau^k$. Moreover, $N_k = N$ for all k sufficiently large. So, without loss of generality, we assume that the full sample is used for the gradient and the function and $\|\nabla f_{\mathcal{N}}(x^k)\| < \bar{\eta}$.

Employing (6) and Lemma 2.2, we have the following estimate

$$\begin{aligned} \|x^k + t_k s^k - x^*\|^2 &\leq \frac{2}{\lambda_1} (f_{\mathcal{N}}(x^k + t_k s^k) - f_{\mathcal{N}}(x^*)) \\ &\leq \frac{2}{\lambda_1} (f_{\mathcal{N}}(x^k) - f_{\mathcal{N}}(x^*) + t_k (\nabla f_{\mathcal{N}}(x^k))^T s^k + \frac{\lambda_n}{2} \|t_k s^k\|^2) \\ &\leq \frac{2}{\lambda_1} \left(\frac{1}{\lambda_1} \|\nabla f_{\mathcal{N}}(x^k)\|^2 + \frac{\lambda_n}{2} \frac{1}{\lambda_1^2} \|\nabla f_{\mathcal{N}_k}(x^k)\|^2 \right) \\ &\leq \frac{2}{\lambda_1} \left(\frac{1}{\lambda_1} + \frac{\lambda_n}{2\lambda_1^2} \right) \lambda_n^2 \|x^k - x^*\|^2 := V_2 \|x^k - x^*\|^2. \end{aligned} \quad (41)$$

Let us denote by A_k the event $\|\nabla^2 f_{\mathcal{D}_k}(x^k) - \nabla^2 f_{\mathcal{N}}(x^k)\| \leq \gamma_k$. Due to Step 1.c of algorithm GIN-R it follows that $P(A_k) \geq 1 - \alpha_k$, i.e., $P(\bar{A}_k) \leq \alpha_k$. Notice that (41) holds in both cases but in the case of A_k we can derive better estimate.

Assume that A_k happens. Then

$$\begin{aligned} &\|\nabla^2 f_{\mathcal{N}}(x^k) s^k + \nabla f_{\mathcal{N}}(x^k)\| \\ &\leq \|(\nabla^2 f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{D}_k}(x^k)) s^k\| + \|\nabla^2 f_{\mathcal{D}_k}(x^k) s^k + \nabla f_{\mathcal{N}}(x^k)\| \\ &\leq \gamma_k \|s^k\| + \eta_k \|\nabla f_{\mathcal{N}}(x^k)\| \leq (\gamma_k/\lambda_1 + \eta_k) \|\nabla f_{\mathcal{N}}(x^k)\|. \end{aligned}$$

Moreover, as $\gamma_k < C\bar{\eta}$ we can repeat the reasoning used in the proof of Theorem 2.4 to conclude that the full step is accepted for k sufficiently large. As the standard assumptions for (Inexact) Newton method are satisfied one can prove

$$\|x^k + s^k - x^*\| \leq c_1(\|x^k - x^*\| + \gamma_k/\lambda_1 + \eta_k)\|x^k - x^*\| \quad (42)$$

for some positive constant c_1 and for k sufficiently large. Indeed, denoting $\tilde{r}_k = \nabla^2 f_{\mathcal{N}}(x^k)s^k + \nabla f_{\mathcal{N}}(x^k)$, it holds $\|\tilde{r}_k\| \leq (\gamma_k/\lambda_1 + \eta_k)\|\nabla f_{\mathcal{N}}(x^k)\|$ and

$$\|x^k + s^k - x^*\| = \|x^k + (\nabla^2 f_{\mathcal{N}}(x^k))^{-1}\tilde{r}_k - (\nabla^2 f_{\mathcal{N}}(x^k))^{-1}\nabla f_{\mathcal{N}}(x^k) - x^*\|.$$

Since the Newton's method converges quadratically there exists $\kappa > 0$ such that

$$\|x^k - (\nabla^2 f_{\mathcal{N}}(x^k))^{-1}\nabla f_{\mathcal{N}}(x^k) - x^*\| \leq \kappa\|x^k - x^*\|^2.$$

Now, using $\|(\nabla^2 f_{\mathcal{N}}(x^k))^{-1}\| \leq 1/\lambda_1$, $\|\nabla f_{\mathcal{N}}(x^k)\| \leq \lambda_n\|x^k - x^*\|$ and defining $c_1 = \max\{\kappa, \lambda_n/\lambda_1\}$ we obtain (42). Using inequality (39) and squaring inequality (42) we obtain

$$\|x^k + s^k - x^*\|^2 \leq 2c_1^2(B^2\tau^{2k} + (\gamma_k/\lambda_1 + \eta_k)^2)\|x^k - x^*\|^2. \quad (43)$$

Now, we distinguish two cases depending on η_k . Using the result of Lemma 3.2 and assuming that k is sufficiently large we obtain the following.

a) If η_k is defined by (30), for $V_1 = 2c_1^2(B^2 + 2B_1^2 + 2(B_2/\lambda_1)^2)$ we get

$$\begin{aligned} & E(\|x^{k+1} - x^*\|^2) \\ &= P(A_k)E(\|x^{k+1} - x^*\|^2|A_k) + P(\bar{A}_k)E(\|x^{k+1} - x^*\|^2|\bar{A}_k) \\ &\leq (V_1\tau^{2k} + \alpha_k V_2)E(\|x^k - x^*\|^2). \end{aligned} \quad (44)$$

b) Considering $\eta_k = \bar{\eta}$, for $C_1 = 2c_1^2 B^2$ and $C_2 = 2c_1^2(C/\lambda_1 + 1)^2$ we get

$$E(\|x^{k+1} - x^*\|^2) \leq (C_1\tau^{2k} + C_2\bar{\eta}^2 + V_2\alpha_k)E(\|x^k - x^*\|^2). \quad \square \quad (45)$$

We conclude the analysis with the following corollary.

Corollary 3.1 *Assume that the conditions of Theorem 3.2 hold and let $\{x^k\}$ be a sequence generated with Algorithm GIN-R. Then the sequence $\{x^k\}$ converges to x^* in the mean square sense:*

- a) linearly if $\eta_k = \bar{\eta}$ is sufficiently small and $\alpha_k < \frac{1-C_2\bar{\eta}^2}{V_2}$;
- b) superlinearly if η_k is defined by (30) and $\lim_{k \rightarrow \infty} \alpha_k = 0$.

Note that the requirement on the probability α_k in order to get the q -linear convergence influences only the logarithmic factor in (36).

3.2 Unbounded sample - GIN method

In many applications, the number of training points is enlarged over time so the cardinality of the sample set N is actually unbounded. This motivated us to consider the following problem as well

$$\min_{x \in \mathbb{R}^n} f(x) = E(F(x, \xi)), \quad (46)$$

where ξ is a random variable defined on a probability space $(\mathcal{A}, \mathcal{F}, P)$ and F is twice differentiable function with respect to x . Let us denote $f_i(x) := F(x, \xi_i)$ where ξ_i , $i = 1, 2, \dots$ is an i.i.d. sequence of variables following the same distribution as ξ . For example, ξ_i can represent the pair of input-output variables in machine learning problems. Then, we can use the same notation as in the previous sections to define SAA approximation of the objective function and its derivatives: $f_{\mathcal{N}}, \nabla f_{\mathcal{N}}, \nabla^2 f_{\mathcal{D}}$. We will prove that, under appropriate assumptions, GIN converges in m.s. towards the solution of problem (46).

Assumption B1 ξ_i , $i = 1, 2, \dots$ is an i.i.d. sequence of variables.

Next we assume that the sequence of iterates $\{x^k\}$ belongs to a bounded set. This assumption is stronger than the *assumption of bounded moments of iterates* used in [7]. However, [7] employs a fixed step length, assuming the knowledge of the maximum eigenvalue λ_n , while here we employ a line-search with approximate function and derivative values and the assumption is needed to cope with such inexactness in the line-search. The analysis of properties of $F(x, \xi)$ that guaranty this assumption is beyond the scope of this paper.

Assumption B2 There exists a compact set Ω such that $\{x^k\}_{k \in \mathbb{N}} \subseteq \Omega$.

Assumption B3 $F(\cdot, \xi) \in C^2(\mathbb{R}^n)$ for every ξ . F and ∇F are dominated by an integrable functions $M_f(\xi)$ and $M_g(\xi)$, respectively, on an open set containing Ω .

Assumption B1 implies that $E(f_{\mathcal{N}}(x)) = f(x)$. Moreover, B1 and B3 imply that $\nabla f(x) = E(\nabla F(x, \xi))$ and therefore $E(\nabla f_{\mathcal{N}}(x)) = \nabla f(x)$. Furthermore, the Uniform law of large numbers implies that $f_{\mathcal{N}}$ and $\nabla f_{\mathcal{N}}$ almost surely (a.s.) converge to $f(x)$ and $\nabla f(x)$, respectively, uniformly on Ω when N tends to infinity. Denote

$$e_{\mathcal{N}} = \max_{x \in \Omega} |f_{\mathcal{N}}(x) - f(x)|, \quad \tilde{e}_{\mathcal{N}} = \max_{x \in \Omega} \|\nabla f_{\mathcal{N}}(x) - \nabla f(x)\|. \quad (47)$$

Then, $\lim_{N \rightarrow \infty} e_{\mathcal{N}} = 0$ and $\lim_{N \rightarrow \infty} \tilde{e}_{\mathcal{N}} = 0$, a.s. and using the Lebesgue Dominated Convergence Theorem (see Theorem 7.31 of [38]) we obtain

$$\lim_{N \rightarrow \infty} E(e_{\mathcal{N}}) = 0, \quad \lim_{N \rightarrow \infty} E(\tilde{e}_{\mathcal{N}}) = 0. \quad (48)$$

Assuming the strong convexity of f_i as in assumption A1, it is easy to show that $f_{\mathcal{N}}$ is also strongly convex with the same constants λ_1 and λ_n for any \mathcal{N} . Moreover, assuming B3, f also remains strongly convex with the constant λ_1 . Indeed, for an arbitrary i and x, y there holds

$$f_i(y) \geq f_i(x) + \nabla^T f_i(x)(y - x) + \frac{\lambda_1}{2} \|x - y\|^2.$$

Taking the expectation and using that $E(\nabla f_i(x)) = \nabla f(x)$ we obtain the strong convexity of f . Therefore, problem (46) has a unique solution x^* .

Theorem 3.3 *Suppose that the assumptions A1, B1-B3 hold and that $N_k \rightarrow \infty$. Then any sequence $\{x^k\}_{k \in \mathbb{N}}$ generated by GIN converges towards the solution of the problem (46) in the mean square sense.*

Proof. First, notice that assumptions B1-B3 imply that $|f_{\mathcal{N}}(x^k) - f(x^k)| \leq e_{\mathcal{N}}$ for every k , so following the reasoning as in the proof of Theorem 2.2 we obtain

$$f(x^{k+1}) - f(x^*) \leq f_{\mathcal{N}_k}(x^k) - c\bar{t}q\|\nabla f_{\mathcal{N}_k}(x^k)\|^2 + e_{\mathcal{N}_k} - f(x^*) + \nu_k, \quad (49)$$

where $\bar{t} = (1 - c)\lambda_1/\lambda_n$ and $q = (\lambda_1(1 - \bar{\eta})^2)/(\lambda_n^2)$. Now,

$$\begin{aligned} \|\nabla f_{\mathcal{N}_k}(x^k)\|^2 &= \|\nabla f_{\mathcal{N}_k}(x^k) - \nabla f(x^k) + \nabla f(x^k)\|^2 \\ &= \|\nabla f(x^k)\|^2 + 2(\nabla f(x^k))^T(\nabla f_{\mathcal{N}_k}(x^k) - \nabla f(x^k)) \\ &\quad + \|\nabla f_{\mathcal{N}_k}(x^k) - \nabla f(x^k)\|^2 \\ &\geq \|\nabla f(x^k)\|^2 - 2\|\nabla f(x^k)\|\|\nabla f_{\mathcal{N}_k}(x^k) - \nabla f(x^k)\| \\ &\geq \|\nabla f(x^k)\|^2 - 2M_g\tilde{e}_{\mathcal{N}_k}, \end{aligned} \quad (50)$$

where the last inequality follows from (47), continuity of ∇f , Assumption B2 and $M_g = \max_{x \in \Omega} \|\nabla f(x)\|$. On the other hand, strong convexity of f implies that $-\|\nabla f(x^k)\|^2 \leq -\lambda_1(f(x^k) - f(x^*))$. Putting all together into (49) we obtain

$$f(x^{k+1}) - f(x^*) \leq (f(x^k) - f(x^*))(1 - \omega) + 2e_{\mathcal{N}_k} + 2c\bar{t}qM_g\tilde{e}_{\mathcal{N}_k} + \nu_k,$$

where $\omega = c\bar{t}q\lambda_1 \in (0, 1)$. Applying expectation we get

$$E(f(x^{k+1}) - f(x^*)) \leq E(f(x^k) - f(x^*))(1 - \omega) + a_k,$$

where $a_k = 2E(e_{\mathcal{N}_k}) + 2c\bar{t}qM_gE(\tilde{e}_{\mathcal{N}_k}) + \nu_k$. Now, (48), (7) and the assumption that $N_k \rightarrow \infty$ together imply that $\lim_{k \rightarrow \infty} a_k = 0$. Therefore, it follows (see [23]) that

$$\lim_{k \rightarrow \infty} E(f(x^k) - f(x^*)) = 0.$$

Finally, strong convexity implies $\|x^k - x^*\|^2 \leq (f(x^k) - f(x^*))2/\lambda_1$ thus

$$\lim_{k \rightarrow \infty} E(\|x^k - x^*\|^2) = 0. \quad \square$$

3.3 Relaxing the strong convexity - method GIN-M

Let us now consider a relaxation of the strong convexity assumption (4) by letting $\nabla^2 f_i(x)$ be only positive semidefinite while the final objective function remains strongly convex. Similar assumptions are stated in [37]. Notice that Theorem 3.3 does not impose any assumption on the size of the Hessian subsample. On the contrary, when the strong convexity assumption is relaxed, a sufficiently large sample is needed to ensure a positive definite Hessian with some prefixed probability. We use the bound on the Hessian sample size provided in [37] but using a different approach. First, we prove mean square convergence to the solution of (46) under appropriate conditions given in the sequel, while in [37] convergence with some (high) probability for finite sum

problems like (1) is considered. Second, we continue to use CG as the inner solver - although modified in this case to cope with possibly singular matrix.

Assumption C1 The functions f and $F(\cdot, \xi)$ are twice continuously differentiable and there exist $0 < \lambda_1 < 1$ and $\lambda_n > 0$ such that for every x, ξ

$$0 \preceq \nabla_x^2 F(x, \xi) \preceq \lambda_n I \quad \text{and} \quad \lambda_1 I \preceq \nabla^2 f(x) \preceq \lambda_n I.$$

This assumption ensures that the unique solution of the original problem still exists. Moreover, we assume that the Hessian approximations are unbiased.

Assumption C2 For every $x \in \mathbb{R}^n$ and every $i \in \mathbb{N}$ there holds $E(\nabla^2 f_i(x)) = \nabla^2 f(x)$.

This assumption allows us to use the Matrix Chernoff result (see [39]) and to obtain the bound presented in Lemma 1 of [37]. Although we observe unbounded sample, the same result holds. More precisely, under the assumptions C1 and C2 we obtain that, given $\mu \in (0, 1 - \lambda_1)$ the following holds

$$P(\lambda_{\min}(\nabla^2 f_{\mathcal{D}}(x)) \geq \mu) \geq 1 - \alpha \text{ if } D \geq \frac{2\lambda_n(1 - \lambda_1)^2 \ln(n/\alpha)}{\mu^2 \lambda_1} := \bar{D}(\alpha). \quad (51)$$

Given that the subsampled Hessian $\nabla^2 f_{\mathcal{D}_k}$ might be singular, for the computation of the step s^k at Step 2 of Algorithm GIN-R, we proceed as follows. If at iteration j of CG we have $(s_j^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s_j^k = 0$, CG is stopped and $s^k = s_{j-1}^k$ is set. Notice that s^k is still a descent direction as $(s_{j-1}^k)^T \nabla f(x^k) = -(s_{j-1}^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s_{j-1}^k < 0$. The algorithm we use is again GIN-R where the sample size is selected such that subsampled Hessian is positive definite with probability $1 - \alpha$. The modified CG, as explained above, is used. We refer to the obtained procedure as Algorithm GINR-M and its iteration k is detailed in Algorithm 3.2.

Algorithm 3.2: k -th iteration of Method GINR-M

Given $x^k \in \mathbb{R}^n$, $c \in (0, 1)$, $\bar{\eta} \in (0, 1)$, $C > 0$, $\{\nu_k\}$, $\alpha \in (0, 1)$ $\mu \in (0, 1 - \lambda_1)$.

Step 1.a Choose $\mathcal{N}_k, \eta_k \in (0, \bar{\eta})$.

Step 1.b If $N_k \leq \bar{D}(\alpha)$ given in (51), set $\mathcal{D}_k = \mathcal{N}_k$. Else, choose $D_k \geq \bar{D}(\alpha)$ and the subsample \mathcal{D}_k randomly and uniformly from \mathcal{N}_k .

Step 2. Determine s^k with modified CG: if $(s_j^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s_j^k = 0$ for some inner iteration j , set $s^k = s_{j-1}^k$. Otherwise, find the step s^k such that (3) holds.

Step 3. Find the smallest nonnegative integer j such that (8) holds for $t_k = 2^{-j}$ and set $x^{k+1} = x^k + t_k s^k$.

Relaxing the strong convexity results in losing the usual relation between the step and the gradient stated in Lemma 2.2. Therefore, we need the following assumption.

Assumption C3 There exists a constant $M_s > 0$ such that the step generated by GINR-M satisfies $\|s^k\| \leq M_s$ for every k .

A comment is due with respect to the above Assumption. Assume that the step s^k computed at Step 4 of GINR-M has been generated at iteration j of CG. Then it belongs to the Krylov subspace

$$\mathcal{K}_j = \text{span}\{\nabla f_{\mathcal{N}_k}(x^k), (\nabla^2 f_{\mathcal{D}_k}(x^k))\nabla f_{\mathcal{N}_k}(x^k), \dots, (\nabla^2 f_{\mathcal{D}_k}(x^k))^{j-1}\nabla f_{\mathcal{N}_k}(x^k)\}.$$

Let V be an orthonormal basis of \mathcal{K}_j , then $s^k = Vy$, where $y \in \mathbb{R}^j$. Therefore

$$(s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k = y^T V^T \nabla^2 f_{\mathcal{D}_k}(x^k) V y \geq \lambda_{\min}(V^T \nabla^2 f_{\mathcal{D}_k}(x^k) V) \|y\|^2.$$

Therefore, noting that $\|s\| = \|y\|$ as V is orthonormal, from

$$(s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k = -(s^k)^T \nabla f_{\mathcal{N}_k}(x^k)$$

and the Cauchy-Schwartz inequality, there follows

$$\|s^k\| \leq \frac{1}{\lambda_{\min}(V^T \nabla^2 f_{\mathcal{D}_k}(x^k) V)} \|\nabla f_{\mathcal{N}_k}(x^k)\|.$$

Then, Assumption C3 is satisfied if the minimal eigenvalue of the projected subsampled Hessian $V^T \nabla^2 f_{\mathcal{D}_k}(x^k) V$ is bounded away from zero and Assumption B2 holds.

Theorem 3.4 *Suppose that the assumptions C1-C3, B1-B3 hold and that N_k tends to infinity. Then there exist α small enough such that any sequence $\{x^k\}_{k \in \mathbb{N}}$ generated by GINR-M converges towards the solution of (46) in m.s.*

Proof. Let us denote by A_k the event $\lambda_{\min}(\nabla^2 f_{\mathcal{D}_k}(x^k)) \geq \mu$ and note that since N_k tends to infinity, $N_k \geq \bar{D}(\alpha)$ will be satisfied for all k large enough ($k \geq k(\alpha)$). Since we are interested in asymptotic result, without loss of generality we assume that $k \geq k(\alpha)$. Then, in Step 1.b D_k is chosen such that (51) holds and this implies $P(\bar{A}_k) \leq \alpha$.

Assume that A_k happens. Then, we can proceed as in the proof of Theorem 3.3 as f is strongly convex and $\lambda_{\min}(\nabla^2 f_{\mathcal{D}}(x^k)) \geq \mu$ yields $(\nabla f_{\mathcal{N}_k}(x^k))^T s^k \leq -\mu \|s^k\|^2$. We obtain

$$f(x^{k+1}) - f(x^*) \leq (f(x^k) - f(x^*))(1 - \omega) + 2e_{\mathcal{N}_k} + \theta \tilde{e}_{\mathcal{N}_k} + \nu_k,$$

where $\omega = c\bar{t}q\lambda_1 \in (0, 1)$, $\bar{t} = (1 - c)\mu/\lambda_n$, $q = \mu(\frac{1-\bar{q}}{\lambda_n})^2$, $\theta = 2c\bar{t}qM_g$ and $M_g = \max_{x \in \Omega} \|\nabla f(x)\|$.

On the other hand, assume that \bar{A}_k happens. Then by the Taylor expansion and assumption C1 we obtain

$$f(x^{k+1}) - f(x^*) \leq f(x^k) - f(x^*) + t_k (\nabla f(x^k))^T s^k + \frac{1}{2} \lambda_n \|t_k s^k\|^2. \quad (52)$$

Again, using the strong convexity of f we get

$$\begin{aligned} \|t_k s^k\|^2 &= \|x^{k+1} - x^k\|^2 \leq 2(\|x^{k+1} - x^*\|^2 + \|x^k - x^*\|^2) \\ &\leq \frac{4}{\lambda_1} (f(x^{k+1}) - f(x^*) + f(x^k) - f(x^*)). \end{aligned} \quad (53)$$

Moreover,

$$\begin{aligned}
(s^k)^T \nabla f(x^k) &= (s^k)^T (\nabla f(x^k) \pm \nabla f_{\mathcal{N}_k}(x^k)) \\
&\leq \|s^k\| \|\nabla f(x^k) - \nabla f_{\mathcal{N}_k}(x^k)\| + (\nabla f_{\mathcal{N}_k}(x^k))^T s^k \\
&= \|s^k\| \|\nabla f(x^k) - \nabla f_{\mathcal{N}_k}(x^k)\| - (s^k)^T \nabla^2 f_{\mathcal{D}_k}(x^k) s^k \\
&\leq M_s \tilde{e}_{\mathcal{N}_k}, \tag{54}
\end{aligned}$$

where the last inequality follows from the fact that $\nabla^2 f_{\mathcal{D}_k}(x^k)$ is positive semidefinite. Putting (54) into (52), together with (53) we obtain

$$f(x^{k+1}) - f(x^*) \leq (f(x^k) - f(x^*)) (1 + 2\lambda_n/\lambda_1) + 2\lambda_n/\lambda_1 (f(x^{k+1}) - f(x^*)) + M_s \tilde{e}_{\mathcal{N}_k}.$$

Combining all together we get

$$\begin{aligned}
&E(f(x^{k+1}) - f(x^*)) \\
&= P(A_k) E(f(x^{k+1}) - f(x^*) | A_k) + P(\bar{A}_k) E(f(x^{k+1}) - f(x^*) | \bar{A}_k) \\
&\leq E(f(x^k) - f(x^*)) (1 - \omega) + 2E(e_{\mathcal{N}_k}) + \theta E(\tilde{e}_{\mathcal{N}_k}) + \nu_k \\
&+ \alpha (E(f(x^k) - f(x^*)) (1 + 2\lambda_n/\lambda_1) + 2\lambda_n/\lambda_1 E(f(x^{k+1}) - f(x^*))) \\
&+ \alpha M_s E(\tilde{e}_{\mathcal{N}_k}).
\end{aligned}$$

Rearranging the previous inequality and assuming $\alpha < \frac{\lambda_1}{2\lambda_n}$ we obtain

$$E(f(x^{k+1}) - f(x^*)) \leq \tau E(f(x^k) - f(x^*)) + a_k,$$

where

$$\tau = \frac{1 - \omega + \alpha(1 + 2\lambda_n/\lambda_1)}{u}, \quad u = 1 - \alpha 2\lambda_n/\lambda_1$$

and

$$a_k = \frac{1}{u} (\nu_k + 2E(e_{\mathcal{N}_k}) + (\theta + \alpha M_s) E(\tilde{e}_{\mathcal{N}_k})).$$

Notice that $\tau \in (0, 1)$ provided that α is small enough. Moreover, as discussed in the previous proof, a_k tends to zero and the result follows. \square

4 Numerical results

In this section we report on our numerical experience with subsampled IN approaches. The experiments were performed in Matlab R2017a, on a Intel Core i5-6600K CPU 3.50 GHz x 4 16GB RAM. For the approximate solution of the linear systems we used the CG method implemented in the Matlab function `pcg`. No preconditioner is employed and the CG is used in a matrix-free manner. Then, only products of $\nabla f_{\mathcal{D}_k}^2$ with vectors are needed. The aim of this section it to provide numerical evidence of the benefits deriving by the employment of adaptive rules, streaming out from the presented theory, for choosing forcing terms and Hessian sample size. Full gradients and functions, i.e. $N_k = N$ for $k > 0$ are used and we compare the full Inexact Newton (FIN) method with $\eta = 10^{-4}$, the subsampled Hessian (SIN) method with $\eta_k = 10^{-4}$

and $D_k = 0.3N$ for all k 's, the subsampled inexact method with adaptive choices of η_k 's and $D_k = 0.3N$ for all k 's (SINA_FT) and the subsampled inexact method with adaptive choices of η_k 's and D_k 's (SINA_FT_Dk). We also consider a subsampled method with constant $D_k = 0.3N$ and a maximum number of five iterations allowed to pCG (SIN_cg5).

In SINA_FT and SINA_FT_Dk, η_k is chosen as follows:

$$\begin{cases} \eta_k = \min\{0.1, \max\{|f(x_k) - m_{k-1}(s^k)|/\|\nabla f(x_{k-1})\|, 10^{-3}\}\} \\ \eta_0 = 0.1. \end{cases}$$

with $m_{k-1}(s^k)$ given in (31) This choice is made according to (30). Finally, we choose the sample size D_k in SINA_FT_Dk as:

$$D_k = \lceil \max\{c_0 D_0, \min\{c_1 \min\{\frac{1}{\eta_k^2}, \frac{1}{\|\nabla f(x_k)\|^2}\}, N\} \rceil \quad c_0, c_1 > 0$$

The above rule is based on inequalities (36) and (38), so D_k is chosen inversely proportional to γ_k^2 given in (38) as suggested by the bounds in (36). The choice of constants c_0 and c_1 depends on the convergence behaviour of CG method. In fact, if CG converges fast large values of D_k should be used, as the lost in the convergence rate due to less accurate second order information is not compensated by the reduced cost of Hessian-vector products. On the other hand, when CG is slower, smaller values of D_k must be used as a large number of matrix-vector products are needed. Then, c_0 and c_1 are chosen according to the following strategy. We set $c_0 = 1$, $c_1 = 0.05$ in case at the previous iteration CG needed more than 20 iterations. Otherwise we set $c_0 = 2$ and $c_1 = 1$. Moreover, D_0 is set to $0.1N$. This choice is motivated by the fact that we allow D_k to change and increase, then we can start with a small D_k leaving the method free to adaptively modify it.

In all the subsampled methods the set \mathcal{D}_k is chosen randomly using the Matlab function `randperm`. All the methods under comparison are in the framework of Algorithm GIN-R, i.e. the nonmonotone linesearch (8), with $c = 10^{-4}$ and $\nu_k = \max(1, f(x^0))/k^{1.1}$ is applied. This latter choice allows even larger step sizes than specified by the global convergence conditions (see also [14], [26]).

The problem we consider is the binary classification problem. We suppose to have at disposal a training set composed of pairs $\{(a_i, b_i)\}$ with $a_i \in \mathbf{R}^n$, $b_i \in \{-1, +1\}$ and $i = 1, \dots, N$, where b_i denotes the correct sample classification. We perform a logistic regression, then we consider as a training objective function the logistic loss with ℓ_2 regularization, [7], i.e. in problem (1) we have

$$f_i(x) = \log c(x, \xi_i) + \lambda \|x\|^2, \quad c(x, \xi_i) = \frac{1}{1 + e^{-b_i a_i^T x}} \quad (55)$$

where $\xi_i = (a_i, b_i)$. Furthermore, the gradients and the Hessians have special forms,

$$\nabla f_i(x) = \frac{(1 - c(x, \xi_i))}{c(x, \xi_i)} b_i a_i + 2\lambda x, \quad \nabla^2 f_i(x) = -\frac{1 - c(x, \xi_i)}{c^2(x, \xi_i)} a_i a_i^T + 2\lambda I. \quad (56)$$

Note that the evaluation of the full function f requires the evaluation of the quantities $(c(x, \xi_i) - 1)b_i a_i$, for $i = 1, \dots, n$ and once these quantities have been computed

they can be used for evaluating $\nabla f_i(x)$ and $\nabla^2 f_i(x)$, $i = 1, \dots, n$. Then, due to the form of gradient and Hessian of each f_i given in (56), the evaluation of $\nabla f_i(x)$ comes for free and the evaluation of $\nabla^2 f_i$ times a vector is as expensive as evaluating $f_i(x)$. Then, we evaluate the performance of the methods under comparison in terms of full function evaluations (FEV). We underline that one `pcg` iteration costs as $\frac{D_k}{N}$ FEV. Finally, in (55) we set $\lambda = 1/N$.

We used the following three datasets:

- `CINA0` [13], $N = 16033$, $n = 132$;
- `Mushrooms` [30], $N = 5000$, $n = 112$;
- `Gisette` [30], $N = 6000$, $n = 5000$.

We stopped the methods under comparison when $\|\nabla f(x_k)\| \leq 10^{-4}$ or when a maximum number of 50 nonlinear iterations is reached. In order to compute the training error we compute the minimizer x^* with the full Newton method to a tight accuracy, i.e we run it until $\|\nabla f(x_k)\|$ is less than 10^{-8} .

We begin by reporting our result with the `mushrooms` dataset. We first underline that the linear algebra phase for this test is not demanding and the average number of required CG iteration is small, as an example it is around 10 when the adaptive choice of the forcing term is used. In Figure 1 we plot $f_k - f(x^*)$ (training error) versus iterations (left) and versus function evaluations FEV (right). We can observe that as expected the full Inexact Newton (FIN) method is the fastest procedure. Moreover, the adaptive choice of the forcing terms seems to speed up the subsampled inexact procedure. Finally, remarkably, the procedure employing both the adaptive choice of η_k and D_k seems to work quite well. Indeed it is slower than `SINA_FT` in the first stage of the convergence history as it uses a smaller Hessian sample set. In the last stage of the procedure it becomes faster as the sample size increases (see Figure 2). On the other hand, if we look to the computational cost of the procedures we can observe that `SINA_FT_Dk` outperforms all the procedures under comparison. We also note that `SINA_FT` outperforms `SIN`. Overall these results show the efficiency of the proposed adaptive strategies.

To give more insight on our adaptive choices we plot in Figure 2 the values of the forcing terms η_k 's (left) and the value of D_k (right) versus iterations using the `SINA_FT_Dk`. We can observe that whenever η_k becomes smaller D_k increases and the approximation of the hessian improves. Moreover, we note that the adaptive procedure allows the Hessian sample size to decrease when the model does not approximate sufficiently well the function and correspondingly the forcing term is increased.

We also compare, in Figure 3, our adaptive procedure `SINA_FT_Dk` with `SIN_cg5`. It is interesting to note that the adaptive procedure is faster than `SIN_cg5` and greater accuracy can be reached. In terms of FEV `SIN_cg5` is slightly better than `SINA_FT_Dk` till a training error's value of the order 10^{-4} . If more accuracy is needed the adaptive procedure is clearly preferable.

Finally, in Figure 4 we compare the behaviour of testing errors versus iterations (left) and FEVs (right) along the sequences generated by full Newton method `FIN`, the adaptive procedure `SINA_FT_Dk` and `SIN_cg5` method. We evaluated the testing error as follows. Let x^k be the approximation computed by a method using the




Figure 1: Mushrooms, training error versus iteration (left) and versus FEV (right) .




Figure 2: Mushrooms, SINA_FT_Dk, η_k 's values (right) and D_k 's (left) versus iterations.

data in the training set. Then, x^k is used to classify the samples in the testing set made up of $\bar{N} = 3124$ instances $z_i, i = 1, \dots, \bar{N}$ and corresponding b_i 's. The classification error at iteration k is defined as $\frac{1}{\bar{N}} \sum_{i=1}^{\bar{N}} \log(1 + \exp(-b_i z_i^T x^k))$. We can observe that the three methods reached almost the same testing errors, the subsampled approaches require a lower computational cost and again the adaptive procedure outperforms SIN_cg5. We also observe that the testing error quickly steadies. This enlightens that the stopping tolerance we used is tight enough and going on with the iterations would not decrease further the testing error.

Let us consider the CINA0 dataset. In Figure 5 we plot the training error versus iterations (left) and versus F-evaluations (right). The full Inexact Newton method FIN is the fastest one, as expected. On the other hand, also in this case, it is the most expensive. The behaviour of the subsampled procedures is that desirable, with the adaptive procedures that outperforms the SINA_FT in terms of FEV. Note that SINA_FT_Dk is slower than the other two subsampled procedures, but less costly, as expected. In fact this is a problem where the linear algebra phase is more demanding, the average




Figure 3: Mushrooms, training error versus iteration (left) and versus FEV (right) .




Figure 4: Mushrooms, testing error versus iteration (left) and versus FEV (right).

number of CG iteration using `SINA_FT` is about 70, and small values of D_k are used to limit the overall computational cost (see Figure 6, left). Since the convergence of CG is slow `SIN_cg5` is not able to converge with a reasonable rate and it can only provide a very rough accuracy (see Figure 6, right)

We finally show the results obtained with the `Gisette` dataset, where n is larger than in the previous tests. In Figure 7, we compare `SINA_FT_DK`, `SIN` and `SIN_FT` in terms of function evaluations FEV (left) and we also report the behaviour of the sample size along the iterations (right). We can observe that `SIN_FT` and `SINA_FT_DK` are less expensive and more accurate than `SIN`, however within 50 nonlinear iterations `SINA_FT_DK` is not able to produce an approximation as accurate as that provided by `SIN_FT`. In fact, the convergence is slow due to the small size of the Hessian subsample set. However, note that if a training error of the order of 10^{-2} is enough, `SINA_FT_DK` is the method of choice as it is less expensive. In case a greater accuracy is needed $D_0 = 0.3N$ should be used in `SINA_FT_DK`. Figure 8 refers to this choice of D_0 . In this case `SINA_FT_DK` outperforms both `SIN` and `SIN_FT`. We also report the values of D_k , η_k and the number of cg-iterations at each Newton iteration.




Figure 5: CINA0, training error versus iteration (left) and versus FEV (right).




Figure 6: CINA0, SINA_FT_DK, D_k 's (left) and training error (right) versus iterations.

Plots show that we use small values of D_k and large values of η_k till the last stage of convergence. Since we are solving linear systems with a rough accuracy, the number of cg-iterations is reasonable (except for a few occurrences) considering that $n = 5000$ and we are not employing a preconditioner.

As a final comment we observe that in SINA_FT and SIN different choices of the Hessian sample size can be adopted and those corresponding to $D_k < 0.3N$ are of particular interest. Then, we performed runs also with $D_k = 0.1N$ for all k 's. SIN method was not competitive, showing that adaptivity of the forcing term is crucial. The results obtained with SINA_FT are strongly dependent on the test problem; below we provide some statistics referring to the average number of FEV employed over 50 runs. In the solution of Gisette test, SINA_FT with $D_k = 0.1N$ is not able to reach the same level of accuracy obtained with $D_k = 0.3N$ as the Hessian subsample is too small. In the solution of the Mushrooms test it is 23% more expensive than SINA_FT with $D_k = 0.3N$ and 30% more expensive than SINA_FT_DK, showing that a fixed sample size larger than $D_k = 0.1N$ is preferable and the adaptivity makes the method




Figure 7: Gisette, training error versus FEV (left), D_k 's versus iterations (right).




Figure 8: Gisette (SINA_FT_DK with $D_0 = 0.3N$), training error versus FEV (top-left), D_k 's versus iterations (top-right), η_k 's versus iterations (bottom-left) and cg-iterations versus Newton iterations. (bottom-right)

able to recovery from the non-ideal choice $D_0 = 0.1N$ providing an overall faster solution. When applied to the CINA0 test, it is 40% less expensive than SINA_FT with $D_k = 0.3N$ and 11% less expensive than SINA_FT_DK. Then, in this latter case the smaller fixed sample size $D_k = 0.1N$ is the best choice, but the adaptive rule provide reasonable values of the parameters avoiding tuning for each data set.

Acknowledgements

The authors are grateful to the anonymous referees for their suggestions, which lead to significant improvement of the manuscript.

References

- [1] Bastin F., Trust-Region Algorithms for Nonlinear Stochastic Programming and Mixed Logit Models, *PhD thesis, University of Namur, Belgium, 2004.*
- [2] Bastin F., Cirillo C., Toint P.L., An adaptive Monte Carlo algorithm for computing mixed logit estimators, *Computational Management Science* 3(1), 55-79, 2006.
- [3] Bastin F., Cirillo C., Toint P.L., Convergence theory for nonconvex stochastic programming with an application to mixed logit, *Mathematical Programming*, 108, 207-234, 2006.
- [4] Bellavia, S., Gurioli G., Morini B., Theoretical study of an adaptive cubic regularization method with dynamic inexact Hessian information, *arXiv preprint arXiv:1808.06239*, 2018.
- [5] Berahas A. S., Bollapragada R., Nocedal J., An Investigation of Newton-Sketch and Subsampled Newton Methods, *arXiv preprint arXiv:1705.06211v3*, 2018.
- [6] Birgin, E.G., Krejić, N., Martínez, J.M., On the employment of Inexact Restoration for the minimization of functions whose evaluation is subject to programming errors, *Mathematics of Computation (to appear)*.
- [7] Bollapragada, R., Byrd, R., Nocedal, J., Exact and Inexact Subsampled Newton Methods for Optimization, *MA Journal Numerical Analysis*, 2018.
- [8] Bottou, L., Curtis F.C., Nocedal, J. Optimization Methods for Large-Scale Machine Learning, *arXiv preprint arXiv:1606.04838v2 [stat.ML]*, 2017.
- [9] Byrd R.H., Chin G.M., Nocedal J., Wu Y., Sample size selection in optimization methods for machine learning, *Mathematical Programming*, 134(1), 127-155, 2012.
- [10] Byrd R.H., Chin G.M., Neveitt W., Nocedal J., On the Use of Stochastic Hessian Information in Optimization Methods for Machine Learning, *SIAM Journal on Optimization*, 21 (3), 977-995, 2011.
- [11] Byrd R.H., Hansen S.L., Nocedal J., Singer Y., A Stochastic Quasi-Newton Method for Large-Scale Optimization, *SIAM Journal on Optimization*, 26(2), 1008-1021, 2016.

- [12] Chernoff H., A measure of the asymptotic efficiency for tests of a hypothesis based on the sum of observations, *Annals of Mathematical Statistics* 23(4), 493-507, 1952.
- [13] Causality workbench team, A marketing dataset, <http://www.causality.inf.ethz.ch/data/CINA.html>, 2008.
- [14] M.A. Diniz-Ehrhardt, J.M. Martínez, M. Raydan, A derivative-free nonmonotone line-search technique for unconstrained optimization, *Journal of Computational and Applied Mathematics*, 219 (2008) 383 – 397.
- [15] Dembo, R.S. Eisenstat, S.C. Steihaug, T., Inexact Newton method, *SIAM Journal on Numerical Analysis* 19(2), 400-409, 1982.
- [16] Deng G., Ferris, M. C., Variable-Number Sample Path Optimization, *Mathematical Programming*, 117 (1-2), 81-109, 2009.
- [17] Eisenstat, S.C., Walker, H.F., Choosing the Forcing Terms in an Inexact Newton Method, *SIAM Journal on Scientific Computing*, 17(1), 16-32, 1996.
- [18] Fountoulakis, K., Godzio, J., A second order method for strongly convex ℓ_1 -regularization problems, *Mathematical Programming*, 156, 189-219, 2016.
- [19] Erdogdu M. A., Montanari A., Convergence rates of sub-sampled Newton methods, *NIPS'15 Proceedings of the 28th International Conference on Neural Information Processing Systems*, 2, 3052-3060, 2015.
- [20] Friedlander M.P., Schmidt M., Hybrid deterministic-stochastic methods for data fitting, *SIAM Journal on Scientific Computing*, 34(3), 1380-1405, 2012.
- [21] Grapiglia G.N., Sachs E.W., On the worst-case evaluation complexity of non-monotone line search algorithms, *Computational Optimization and Applications*, 68(3), 555-577, 2017.
- [22] Hoeffding, W., Probability inequalities for sums of bounded random variables, *American Statistical Association Journal*, 58(103), 13-30, 1963.
- [23] Jakovetic D., Xavier J., Moura J.M.F. Fast Distributed Gradient Methods, *IEEE Trans. Autom. Contr.*, 59(5), 1131-1146, 2014.
- [24] Kelley, C.T., *Iterative Methods for Linear and Nonlinear Equations*, SIAM, 1995.
- [25] Krejić N., Krklec N., Line search methods with variable sample size for unconstrained optimization, *Journal of Computational and Applied Mathematics* 245, 213-231, 2013.
- [26] Krejić N., Krklec Jerinkić N., Nonmonotone line search methods with variable sample size, *Numerical Algorithms* 68(4), 711-739, 2015.
- [27] Krejić, N., Martínez, J.M., Inexact Restoration approach for minimization with inexact evaluation of the objective function, *Mathematics of Computation*, 85 (2016) 1775-1791. .

- [28] Lee, Jason, D., Yuekai Sun, Michael A. Sandres, Proximal Newton type methods for minimizing composite functions, *SIAM Journal on Optimization* 24, 3 (2014), 1420-1443.
- [29] Li D.H., Fukushima M., A derivative-free line search and global convergence of Broyden-like method for nonlinear equations, *Optimization Methods and Software* 13, 181-201, 2000.
- [30] Lichman M., UCI machine learning repository, <https://archive.ics.uci.edu/ml/index.php>, 2013.
- [31] Minsker S., On some extensions of Bernstein’s inequality for self-adjoint operators, *arXiv:1112.5448 [math.PR]*, 2017.
- [32] Nash S.G., A survey of truncated-Newton methods, *Journal of Computational and Applied Mathematics* 124(1-2), 45-59, 2000.
- [33] Nesterov, Y., Introductory Lectures on Convex Optimization: A Basic Course, Springer Science and Media, Vol. 87, 2013.
- [34] Pasupathy R., On Choosing Parameters in Retrospective-Approximation Algorithms for Stochastic Root Finding and Simulation Optimization, *Operations Research* 58(4), 889-901, 2010.
- [35] Pilanci M., Wainwright M. J., Newton sketch: A near linear-time optimization algorithm with linear-quadratic convergence, *SIAM Journal on Optimization* 27(1), 205-245, 2017.
- [36] Polak E., Royset J.O., Efficient sample sizes in stochastic nonlinear programming, *Journal of Computational and Applied Mathematics* 217(2), 301-310, 2008.
- [37] F. Roosta-Khorasani, M.W. Mahoney, Sub-Sampled Newton Methods, *Mathematical Programming*, 174, 293–326, 2019.
- [38] A. Shapiro, D. Dentcheva, A. Ruszczyński, Lectures on stochastic programming: Modeling and theory, *MPS/SIAM Series on Optimization* 9, 2009.
- [39] Tropp, J.A., User-Friendly Tail Bounds for Sums of Random Matrices, *Foundations of Computational Mathematics*, 12 (2012), 389-434.
- [40] Xu P., Yang J., Roosta-Khorasani F., Ré C., and Mahoney M.W., Sub-sampled Newton methods with non-uniform sampling, *Advances in Neural Information Processing Systems 30 (NIPS)*, 2530-2538, 2016.
- [41] Xu P., Roosta-Khorasani F., and Mahoney M.W., Newton-Type Methods for Non-Convex Optimization Under Inexact Hessian Information, *arXiv preprint, arxiv:1708.07164v3 [math.OC]*, 2018.
- [42] Zhang H., Hager W., A nonmonotone line search technique and its application to unconstrained optimization *SIAM Journal on Optimization*, 4, 1043-1056, 2004.

5 Appendix

Proofs of Lemmas 2.6, 2.7 and 3.1

Proof of Lemma 2.6. Using the Mean Value Theorem we get the following inequality:

$$\|\nabla f_{\mathcal{N}}(x^k + s^k) - \nabla f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{N}}(x^k)s^k\| \leq \frac{L}{2}\|s^k\|^2. \text{ Then,}$$

$$\begin{aligned} \|\nabla f_{\mathcal{N}}(x^k + s^k) - \nabla f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{D}}(x^k)s^k\| &\leq \|\nabla f_{\mathcal{N}}(x^k + s^k) - \nabla f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{N}}(x^k)s^k\| \\ &\quad + \|\nabla^2 f_{\mathcal{D}_k}(x^k) - \nabla^2 f_{\mathcal{N}}(x^k)\|\|s^k\| \\ &\leq \|s^k\|(h(D_k, x^k) + \frac{1}{2}L\|s^k\|). \end{aligned}$$

Therefore,

$$\begin{aligned} \|\nabla f_{\mathcal{N}}(x^k + s^k)\| &\leq \|\nabla f_{\mathcal{N}}(x^k) + \nabla^2 f_{\mathcal{D}_k}(x^k)s^k\| + \|\nabla f_{\mathcal{N}}(x^k + s^k) - \nabla f_{\mathcal{N}}(x^k) - \nabla^2 f_{\mathcal{D}_k}(x^k)s^k\| \\ &\leq \eta\|\nabla f_{\mathcal{N}}(x^k)\| + \|s^k\|(\frac{1}{2}L\|s^k\| + h(D_k, x^k)) \end{aligned}$$

and, by Lemma 2.2,

$$\|\nabla f_{\mathcal{N}}(x^k + s^k)\| \leq \eta\|\nabla f_{\mathcal{N}}(x^k)\| + \frac{1}{\lambda_1}\|\nabla f_{\mathcal{N}}(x^k)\|(\frac{1}{2\lambda_1}L\|\nabla f_{\mathcal{N}}(x^k)\| + h(D_k, x^k)). \quad \square$$

Proof of Lemma 2.7. By Lemma 2.2 and (5) we have

$$\begin{aligned} \|x^k + s^k - x^*\| &\leq \|x^k - x^*\| + \|s^k\| \leq \|x^k - x^*\| + \lambda_1^{-1}\|\nabla f_{\mathcal{N}}(x^k)\| \\ &\leq \|x^k - x^*\| + \lambda_1^{-1}\lambda_n\|x^k - x^*\| \leq (1 + \lambda_1^{-1}\lambda_n)\|x^k - x^*\| \leq \delta^*. \end{aligned}$$

□

Proof of Lemma 3.1. Let us define $Y_i(x) = (H_i(x) - \nabla^2 f_{\mathcal{N}}(x))/D$ where $H_i(x)$ is a randomly chosen Hessian. Then,

$$\sum_{i=1}^D Y_i(x) = \nabla^2 f_{\mathcal{D}}(x) - \nabla^2 f_{\mathcal{N}}(x).$$

Also, notice that $E(Y_i(x)) = 0$ and that the Weyl's inequality yields $\lambda_{max}(Y_i(x)) \leq \lambda_n/D$, $\lambda_{min}(Y_i(x)) \geq -\lambda_n/D$. Then, $\|Y_i(x)\| \leq \lambda_n/D$ and

$$\tilde{\sigma}^2(x) := \left\| \sum_{i=1}^D E(Y_i^2(x)) \right\| \leq \sum_{i=1}^D E(\|Y_i(x)\|^2) \leq \frac{\lambda_n^2}{D}.$$

Then, using the Bernstein's inequality (see [39, Theorem 1.6]) we derive

$$P(\|\nabla^2 f_{\mathcal{D}}(x) - \nabla^2 f_{\mathcal{N}}(x)\| \geq \gamma) \leq 2n \exp\left(-\frac{\gamma^2/2}{\lambda_n^2/D + (\lambda_n/D)\gamma/3}\right).$$

This yields the bound. □