

Free idempotent generated semigroups: maximal subgroups and the word problem

Igor Dolinka

dockie@dmi.uns.ac.rs

Department of Mathematics and Informatics, University of Novi Sad

The 17th NBSAN Meeting
Norwich, UK, April 14, 2014



Man is condemned to be free.

Jean-Paul Sartre

Idempotent generated semigroups

Many natural semigroups are idempotent-generated ($S = \langle E(S) \rangle$):

- ▶ The semigroup $\mathcal{T}_n \setminus \mathcal{S}_n$ of singular (non-invertible) transformations on a finite set (Howie, 1966);
- ▶ The singular part of $\mathcal{M}_n(\mathbb{F})$, the semigroup of all $n \times n$ matrices over a field \mathbb{F} (Erdos (not Paul!), 1967);
- ▶ Classification of linear algebraic monoids that are idempotent-generated (Putcha, 2006);
- ▶ The singular part of \mathcal{P}_n , the singular part of the partition monoid on a finite set (East, FitzGerald, 2012);

Hence:

*What can we say about the structure of the free-est idempotent-generated (IG) semigroup with a **fixed structure/configuration of idempotents** ???*

Biordered sets of idempotents

'Configuration of idempotents' = **biordered sets**

Basic pair $\{e, f\}$ of idempotents:

$$\{e, f\} \cap \{ef, fe\} \neq \emptyset$$

(If e.g. $ef \in \{e, f\}$, then $(fe)^2 = fe$, and conversely.)

Biordered set of a semigroup S = the partial algebra on $E(S)$ obtained by retaining the products of basic pairs (in S).

Alternatively, biordered sets can be (abstractly) described as **relational structures** $(E(S), \leq^{(l)}, \leq^{(r)})$ with two quasi-orders and several simple rules/axioms (Easdown, Nambooripad, '80s).

Free IG semigroups: idea

- ▶ To every semigroup S with idempotents E associate the free-est semigroup $IG(E)$ whose idempotents form the same biordered set as in S .
- ▶ To every regular semigroup S with idempotents E associate the free-est **regular** semigroup $RIG(E)$ in whose idempotents form the same biordered set as in S .

Free IG semigroups: formal definitions

Let E be the biordered set of idempotents of a semigroup S .


$$\text{IG}(E) := \langle E \mid e \cdot f = ef \text{ where } \{e, f\} \text{ is a basic pair} \rangle.$$

Suppose now S is regular. We define the **sandwich sets**:


$$S(e, f) = \{h \in E : ehf = ef, fhe = h\} \neq \emptyset$$

$$\text{RIG}(E) := \langle E \mid \text{IG}, ehf = ef \text{ (} e, f \in E, h \in S(e, f)) \rangle.$$

Example 1: V-semilattice



$$\text{IG}(S) = \langle e, f, z \mid e^2 = e, f^2 = f, z^2 = z, ez = ze = fz = zf = z \rangle:$$



$$\text{RIG}(S) = \langle e, f, z \mid \text{IG}, ef = fe = z \rangle = S.$$

Example 2: 2×2 rectangular band

$$S = \langle e_{ij} \mid e_{ij}e_{kl} = e_{il} \ (i, j, k, l \in \{1, 2\}) \rangle:$$

e_{11}	e_{12}
e_{21}	e_{22}

$$\text{IG}(S) = \langle e_{ij} \mid e_{ij}e_{kl} = e_{il} \ (i, j, k, l \in \{1, 2\}, i = k \text{ or } j = l) \rangle:$$

$(e_{11}e_{22})^i e_{11}$	$(e_{12}e_{21})^i e_{12}$
$(e_{12}e_{21})^i$	$(e_{11}e_{22})^i$
$(e_{21}e_{12})^i e_{21}$	$(e_{22}e_{11})^i e_{22}$
$(e_{22}e_{11})^i$	$(e_{21}e_{12})^i$

$$\text{RIG}(S) = \text{IG}(S).$$

Relationships between $S = \langle E \rangle$, $IG(E)$, and $RIG(E)$


- ▶ (Easdown, 1985) There exists a surjective homomorphism $\phi : IG(E) \rightarrow S$ such that
 - ▶ the restriction of ϕ to E is an **isomorphism** of biordered sets;
 - ▶ the maximal subgroup H_e in S is the **ϕ -image** of its counterpart in $IG(E)$ (which is in turn isomorphic to its counterpart in $RIG(E)$).
- ▶ The 'eggbox picture' of the \mathcal{D} -class of e has the **same dimensions** in all three.
- ▶ $IG(E)$ may contain other, **non-regular** \mathcal{D} -classes.

So, understanding $IG(E)$ is essential in understanding the structure of arbitrary IG semigroups.


Question

Which groups arise as maximal subgroups of $IG(E)$ (and thus of $RIG(E)$)?

The big picture



Let's zoom in



Presentation for a max. subgroup of $IG(E)$: Generators

Fact


G is generated by a set in 1-1 correspondence with $D \cap E(S)$.

D				generators of G			
$e_{(11)}$	e_{12}	e_{13}		f_{11}	f_{12}	f_{13}	
	e_{22}		e_{24}		f_{22}		f_{24}
e_{31}	e_{32}	e_{33}	e_{34}	f_{31}	f_{32}	f_{33}	f_{34}

$$G = \langle f_{ij} (e_{ij} \in D \cap E) \mid ??? \rangle$$

Typical relations: $f_{ij}^{-1}f_{il} = f_{kj}^{-1}f_{kl}$

• $h = h^2$



Singular square $\begin{bmatrix} e_{ij} & e_{il} \\ e_{kj} & e_{kl} \end{bmatrix} \Rightarrow \text{relation } f_{ij}^{-1}f_{il} = f_{kj}^{-1}f_{kl}.$

Presentation – Approach #1

Theorem (Nambooripad 1979; Gray, Ruškuc 2012)

The maximal subgroup G of $e \in E$ in $IG(E)$ or $RIG(E)$ is defined by the presentation:

$$\begin{aligned} \langle f_{ij} \mid & f_{i,\pi(i)} = 1 \quad (i \in I), \\ & f_{ij} = f_{il} \quad (\text{if } r_j e_{il} = r_l \text{ is a Schreier rep.}), \\ & f_{ij}^{-1} f_{il} = f_{kj}^{-1} f_{kl} \left(\begin{bmatrix} e_{ij} & e_{il} \\ e_{kj} & e_{kl} \end{bmatrix} \text{ sing. sq.} \right) \rangle. \end{aligned}$$

Proof: Reidemeister–Schreier rewriting process followed by Tietze transformations.


Graham-Houghton complex

Let S be an idempotent generated regular semigroup.

$GH(S)$: a 2-complex whose connected components are in a 1-1 correspondence with \mathcal{D} -classes of S .

D

$e_{(11)}$	e_{12}	e_{13}	
	e_{22}		e_{24}
e_{31}	e_{32}	e_{33}	e_{34}



Presentation – Approach #2

Theorem (Brittenham, Margolis, Meakin, 2009)

The fundamental group of $GH(S)$ at any point of its connected component C_e containing the edge $e \cong$ the maximal subgroup of $RIG(E(S))$ (and thus of $IG(E(S))$) containing e .

So,...

... let \mathcal{T} be an arbitrary spanning tree of C_e . Then the maximal subgroup G of $e \in E$ in $IG(E)$ (or $RIG(E)$) is defined by the presentation:

$$\langle f_{ij} \mid f_{ij} = 1 \quad ((i, j) \in \mathcal{T}), \\ f_{ij} f_{kj}^{-1} f_{kl} f_{il}^{-1} = 1 \quad ((i, j, k, l) \text{ is a 2-cell}) \rangle.$$

Obviously, a clever choice of \mathcal{T} may speed up the computation.

Remarks (1)

- ▶ Two types of relations:
 - ▶ Initial conditions: declaring some generators equal to 1 (or to each other in approach #1);
 - ▶ Main relations: one per singular square.
- ▶ All relations of length ≤ 4 .
- ▶ What can be defined by relations $f_{ij}^{-1}f_{il} = f_{kj}^{-1}f_{kl}$?
- ▶ $\begin{bmatrix} 1 & b \\ a & c \end{bmatrix} \Rightarrow ab = c.$

Remarks (2)

- ▶ But: Every semigroup can be defined by relations of the form $ab = c$.
- ▶ Even better: Every finitely presented semigroup can be defined by finitely many relations of the form $ab = c$.
- ▶ Some more special squares ...
- ▶ $\begin{bmatrix} a & a \\ b & c \end{bmatrix} \Rightarrow b = c.$
- ▶ $\begin{bmatrix} 1 & 1 \\ 1 & a \end{bmatrix} \Rightarrow a = 1.$

The freeness conjecture

Question

Which groups arise as maximal subgroups of $IG(E)$ (and thus of $RIG(E)$)?

- ▶ Work of Pastijn and Nambooripad ('70s and '80s) and McElwee (2002) led to the belief that these maximal subgroups must always be **free groups** (of a suitable rank).
- ▶ This conjecture was proved false by **Brittenham, Margolis, and Meakin** in 2009 who obtained the groups $\mathbb{Z} \oplus \mathbb{Z}$ (from a particular 73-element semigroup) and \mathbb{F}^* for an arbitrary field \mathbb{F} .
- ▶ Finally, Gray and Ruškuc (2012) proved that **every** group arises as a maximal subgroup of some free idempotent generated semigroup (!!!).

Theorem

*Every group is a maximal subgroup of some free idempotent generated semigroup (over a **regular** semigroup).*

Theorem

*Every **finitely presented** group is a maximal subgroup of some free idempotent generated semigroup arising from a **finite** semigroup.*

Remark

Maximal subgroups of free idempotent generated semigroups arising from finite semigroups have to be finitely presented by Reidemeister–Schreier.

Remaining Question

Is every **finitely presented** group a maximal subgroup in some free idempotent generated semigroup over a **finite regular** semigroup?

Calculating the groups for natural examples of S

Some or all maximal subgroups in $IG(E(S))$ have been calculated for the following S :

- ▶ Full transformation monoids: Gray, Ruškuc (symmetric groups, provided $\text{rank} \leq n - 2$);
- ▶ Partial transformation monoids: ID (symmetric groups again);
- ▶ Full matrix monoid over a skew field: Brittenham, Margolis, Meakin ($\text{rank} = 1$); ID, Gray ($\text{rank} < n/3$ – general linear groups);
- ▶ Endomorphism monoid of a free G -act: Dolinka, Gould, Yang (wreath products of G by symmetric groups).

Bands

Theorem (ID)

For every left- or right seminormal band B , all maximal subgroups of $IG(B)$ are free. For every variety \mathbf{V} not contained in $\mathbf{LSNB} \cup \mathbf{RSNB}$ there exists $B \in \mathbf{V}$ such that $IG(B)$ contains a non-free maximal subgroup.

Remaining Question

Which groups arise as maximal subgroups of $IG(B)$, B a (finite) band?

Answer (ID, Ruškuc, 2013): All of them! (Resp. all finitely presented ones.)

New construction (ID & Ruškuc): set-up

Suppose we want to obtain


$$G = \langle a, b, c, \dots \mid ab = c, \dots \rangle$$

as a maximal subgroup of $IG(B)$ for a band B .

- ▶ $I = \{0, a, b, c, \dots, 0', a', b', c', \dots\}$;
- ▶ $J = \{0, a, b, c, \dots, \infty\}$;
- ▶ $\mathcal{T} = \mathcal{T}_I^* \times \mathcal{T}_J$;
- ▶ the minimal ideal: $K = \{(\sigma, \tau) : \sigma, \tau \text{ constants}\}$;
- ▶ K is an $I \times J$ rectangular band.

New construction: set-up

- ▶ $B = K \cup L$, where L is a left zero semigroup.
- ▶ We ensure this by virtue of every $(\sigma, \tau) \in L$ satisfying:
 - ▶ $\sigma^2 = \sigma, \tau^2 = \tau$;
 - ▶ $\ker(\sigma) = \{\{0, a, b, c\}, \{0', a', b', c'\}\}$;
 - ▶ thus σ is determined by its image $\{x, y\}$ transversing its kernel;
 - ▶ $\text{im}(\tau) = \{0, a, b, c\}$;
 - ▶ thus τ is specified by $(\infty)\tau$.



New construction: the action of L on K

Notation	Indexing	$\text{im}(\sigma)$	$(\infty)\tau$
(σ_0, τ_0)	–	$\{0, 0'\}$	0
(σ_a, τ_a)	$a \in A$	$\{0, a'\}$	a
$(\bar{\sigma}_a, \bar{\tau}_a)$	$a \in A$	$\{a, a'\}$	0
$(\sigma_{\mathbf{r}}, \tau_{\mathbf{r}})$	$\mathbf{r} = (ab, c) \in R$	$\{b, c'\}$	a

New construction: the endgame

	0	a	b	c	∞
0	f_{00}	f_{0a}	f_{0b}	f_{0c}	$f_{0\infty}$
a	f_{a0}	f_{aa}	f_{ab}	f_{ac}	$f_{a\infty}$
b	f_{b0}	f_{ba}	f_{bb}	f_{bc}	$f_{b\infty}$
c	f_{c0}	f_{ca}	f_{cb}	f_{cc}	$f_{c\infty}$
0'	$f_{0'0}$	$f_{0'a}$	$f_{0'b}$	$f_{0'c}$	$f_{0'\infty}$
a'	$f_{a'0}$	$f_{a'a}$	$f_{a'b}$	$f_{a'c}$	$f_{a'\infty}$
b'	$f_{b'0}$	$f_{b'a}$	$f_{b'b}$	$f_{b'c}$	$f_{b'\infty}$
c'	$f_{c'0}$	$f_{c'a}$	$f_{c'b}$	$f_{c'c}$	$f_{c'\infty}$

(σ_0, τ_0)
 (σ_a, τ_a)
 $(\bar{\sigma}_a, \bar{\tau}_a)$
 \longrightarrow

	0	a	b	c	∞
0	1	1	1	1	1
a	1	1	1	1	a
b	1	1	1	1	b
c	1	1	1	1	c
0'	1	a	b	c	1
a'	1	a	b	c	a
b'	1	a	b	c	b
c'	1	a	b	c	c

$$(\sigma_r, \tau_r) \quad r : ab = c$$

The word problem

(ongoing joint work with R.D.Gray and N.Ruškcuc)

Let S be a semigroup with a finite biordered set $E = E(S)$.

Theorem

There exists an algorithm deciding whether $w \in E^+$ represents a regular element of $IG(E)$.

Key tool: generalised sandwich sets.

Theorem

If every maximal subgroup of $IG(E)$ has a solvable word problem, then there is an algorithm which, given $u, v \in E^+$ such that u represents a regular element, decides whether $u = v$ holds in $IG(E)$.

Corollary

The word problem for $RIG(E)$ is solvable iff the word problem for each of its maximal subgroups is solvable.

The word problem(?)

So, the following question naturally arises:

Question

Is it true that the word problem for $IG(E)$ (where $E = E(S)$ is finite) is solvable iff the word problem for each of its maximal subgroups is solvable?

ID + RG + NR (2013/14): **NO!**

Theorem

There exists a finite band B such that all the maximal subgroups of $IG(E(B))$ are free, but the word problem of $IG(E(B))$ is still undecidable.

The band $B_{G,H}$

Let G be a finitely presented group and H its finitely generated group. The band $B_{G,H}$ has 5 \mathcal{D} -classes, forming 3 'floors', from top to bottom:

- ▶ a left zero band L ,
- ▶ an 'intermediate' rectangular band K_1 ,
- ▶ a 0-direct union of two copies K' , K'' of the rectangular band K from the ID-NR construction,
- ▶ the action of L on K' and K'' is exactly the same as in the ID-NR construction.

The band $B_{G,H}$

Properties of $B_{G,H}$:

- ▶ any maximal subgroup of $\text{IG}(E(B_{G,H}))$ is either trivial, or isomorphic to G ,
- ▶ as known from the properties of $\text{IG}(E)$, to the rectangular bands K' and K'' correspond two \mathcal{D} -classes M' and M'' of $\text{IG}(E)$ which are completely simple subsemigroups, with typical elements

$$(i', g_1, j') \quad \text{and} \quad (i'', g_2, j''),$$

$$g_1, g_2 \in G.$$

Proposition

$(1', 1, 1')(1'', 1, 1'') = (1', 1, 1')(1'', g, 1'')$ if and only if $g \in H$.

The Mikhailova construction

Hence, it suffices to take G with a solvable word problem and its finitely generated subgroup H with an undecidable membership problem!

If $G = F_2 \times F_2$ and W is a f.p. 2-generated group with an undecidable problem, then taking H to be the fibre product w.r.t. the natural homomorphism $\pi : F_2 \rightarrow W$ (i.e. $H = \ker \pi$) suffices.

Open Problem

Is it at least true that the word problem for $IG(E)$ is solvable when E is the border of a finite normal band?

THANK YOU!

Questions and comments to:

dockie@dmu.ac.uk

Further information may be found at:

<http://people.dmu.ac.uk/~dockie>